

LIALDA- mesalamine tablet, delayed release
Shire US Manufacturing Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use LIALDA safely and effectively. See full prescribing information for LIALDA.

LIALDA[®] (mesalamine) delayed-release tablets, for oral use

Initial U.S. Approval: 1987

----- **INDICATIONS AND USAGE** -----

LIALDA is a locally acting 5-aminosalicylic acid (5-ASA) indicated for the induction of remission in adults with active, mild to moderate ulcerative colitis and for the maintenance of remission of ulcerative colitis. (1)

----- **DOSAGE AND ADMINISTRATION** -----

For induction of remission of active, mild to moderate ulcerative colitis, two to four 1.2 g tablets taken once daily with food. (1, 2)

For maintenance of remission of ulcerative colitis, two 1.2 g tablets taken once daily with food. (1, 2)

----- **DOSAGE FORMS AND STRENGTHS** -----

Delayed-Release Tablets: 1.2 g (3)

----- **CONTRAINDICATIONS** -----

Patients with known hypersensitivity to salicylates or aminosalicylates or to any of the ingredients of LIALDA tablets. (4)

----- **WARNINGS AND PRECAUTIONS** -----

- Renal impairment may occur. Assess renal function at the beginning of treatment and periodically during treatment. (5.1)
- Mesalamine-induced acute intolerance syndrome has been reported. Observe patients closely for worsening of these symptoms while on treatment. (5.2)
- Use caution when treating patients who are hypersensitive to sulfasalazine. (5.3)
- Mesalamine-induced cardiac hypersensitivity reactions (myocarditis and pericarditis) have been reported. (5.3)
- Hepatic failure has been reported in patients with pre-existing liver disease. Use caution when treating patients with liver disease. (5.4)
- Upper GI tract obstruction may delay onset of action. (5.5)
- Use of mesalamine may lead to spuriously elevated test results when measuring urinary normetanephrine by liquid chromatography with electrochemical detection. (5.6)

----- **ADVERSE REACTIONS** -----

- The most common adverse reactions (incidence \geq 2%) are ulcerative colitis, headache, flatulence, liver function test abnormality, and abdominal pain. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Shire US Inc. at 1-800-828-2088 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

----- **DRUG INTERACTIONS** -----

- Nephrotoxic agents including NSAIDs: renal reactions have been reported. (7.1)
- Azathioprine or 6-mercaptopurine: blood disorders have been reported. (7.2)

----- **USE IN SPECIFIC POPULATIONS** -----

- Renal impairment: Use LIALDA with caution in patients with a history of renal disease. (5.1, 7.1, 8.5, 13.2)
- Nursing Women: Caution should be exercised when administered to a nursing woman. (8.3)
- Geriatric Patients: Monitor blood cell counts in geriatric patients. (8.5)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 11/2015

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Renal Impairment
- 5.2 Mesalamine-Induced Acute Intolerance Syndrome
- 5.3 Hypersensitivity Reactions
- 5.4 Hepatic Impairment
- 5.5 Upper GI Tract Obstruction
- 5.6 Interference with Laboratory Tests

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Nephrotoxic Agents, Including Non-Steroidal Anti-Inflammatory Drugs
- 7.2 Azathioprine or 6-mercaptopurine

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 13.2 Animal Toxicology and/or Pharmacology

14 CLINICAL STUDIES

- 14.1 Active, Mild to Moderate Ulcerative Colitis
- 14.2 Maintenance of Remission in Patients with Ulcerative Colitis

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

LIALDA is indicated for the induction of remission in patients with active, mild to moderate ulcerative colitis and for the maintenance of remission of ulcerative colitis.

2 DOSAGE AND ADMINISTRATION

The recommended dosage for the induction of remission in adult patients with active, mild to moderate ulcerative colitis is two to four 1.2 g tablets taken once daily with a meal for a total daily dose of 2.4 g

or 4.8 g. The recommended dosage for the maintenance of remission is two 1.2 g tablets taken once daily with a meal for a total daily dose of 2.4 g.

3 DOSAGE FORMS AND STRENGTHS

The red-brown ellipsoidal delayed-release tablet containing 1.2 g mesalamine is debossed on one side and imprinted with S476.

4 CONTRAINDICATIONS

LIALDA is contraindicated in patients with known hypersensitivity to salicylates or aminosalicylates or to any of the ingredients of LIALDA [see *Warnings and Precautions (5.3)*, *Description (11)*, *Adverse Reactions (6.2)*].

5 WARNINGS AND PRECAUTIONS

5.1 Renal Impairment

Renal impairment, including minimal change nephropathy, acute and chronic interstitial nephritis, and, rarely, renal failure, has been reported in patients given products such as LIALDA that contain mesalamine or are converted to mesalamine.

It is recommended that patients have an evaluation of renal function prior to initiation of LIALDA therapy and periodically while on therapy. Exercise caution when using LIALDA in patients with known renal dysfunction or a history of renal disease.

In animal studies, the kidney was the principal organ for toxicity. [See *Drug Interactions (7.1)* and *Nonclinical Toxicology (13.2)*]

5.2 Mesalamine-Induced Acute Intolerance Syndrome

Mesalamine has been associated with an acute intolerance syndrome that may be difficult to distinguish from an exacerbation of ulcerative colitis. Although the exact frequency of occurrence has not been determined, it has occurred in 3% of patients in controlled clinical trials of mesalamine or sulfasalazine. Symptoms include cramping, acute abdominal pain and bloody diarrhea, and sometimes fever, headache, and rash. Observe patients closely for worsening of these symptoms while on treatment. If acute intolerance syndrome is suspected, promptly discontinue treatment with LIALDA.

5.3 Hypersensitivity Reactions

Some patients who have experienced a hypersensitivity reaction to sulfasalazine may have a similar reaction to LIALDA tablets or to other compounds that contain or are converted to mesalamine.

Mesalamine-induced cardiac hypersensitivity reactions (myocarditis and pericarditis) have been reported with LIALDA and other mesalamine medications. Caution should be taken in prescribing this medicine to patients with conditions predisposing them to the development of myocarditis or pericarditis.

5.4 Hepatic Impairment

There have been reports of hepatic failure in patients with pre-existing liver disease who have been administered mesalamine. Caution should be exercised when administering LIALDA to patients with liver disease.

5.5 Upper GI Tract Obstruction

Pyloric stenosis or other organic or functional obstruction in the upper gastrointestinal tract may cause prolonged gastric retention of LIALDA which would delay mesalamine release in the colon.

5.6 Interference with Laboratory Tests

Use of mesalamine may lead to spuriously elevated test results when measuring urinary normetanephrine by liquid chromatography with electrochemical detection because of the similarity in the chromatograms of normetanephrine and mesalamine's main metabolite, N-acetylaminosalicylic acid (N-Ac-5-ASA). An alternative, selective assay for normetanephrine should be considered.

6 ADVERSE REACTIONS

The most serious adverse reactions seen in Lialda clinical trials or with other products that contain or are metabolized to mesalamine are:

- Renal impairment, including renal failure [See Warnings and Precautions (5.1)]
- Mesalamine-induced acute intolerance syndrome [See Warnings and Precautions (5.2)]
- Hypersensitivity reactions [See Warnings and Precautions (5.3)]
- Hepatic impairment, including hepatic failure [See Warnings and Precautions (5.4)]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

LIALDA has been evaluated in 1368 ulcerative colitis patients in controlled and open-label trials.

Induction of Remission

In two 8-week placebo-controlled clinical trials involving 535 ulcerative colitis patients, 356 received 2.4 g/day or 4.8 g/day LIALDA tablets and 179 received placebo. The most frequent adverse reaction leading to discontinuation from LIALDA therapy was exacerbation of ulcerative colitis (0.8%). Pancreatitis occurred in less than 1% of patients during clinical trials and resulted in discontinuation of therapy with LIALDA in patients experiencing this event.

Adverse reactions occurring in LIALDA or placebo groups at a frequency of at least 1% in two 8-week, double blind, placebo-controlled trials are listed in Table 1. The most common adverse reactions with LIALDA 2.4 g/day and 4.8 g/day were headache (5.6% and 3.4%, respectively) and flatulence (4% and 2.8%, respectively).

Table 1: Adverse Reactions in Two Eight-Week Placebo-Controlled Trials Experienced by at Least 1% of the LIALDA Group and at a Rate Greater than Placebo ^a

| Adverse Reaction | LIALDA 2.4 g/day (n = 177) | LIALDA 4.8 g/day (n = 179) | Placebo (n = 179) |
|------------------------------|----------------------------|----------------------------|-------------------|
| Headache | 10 (5.6%) | 6 (3.4%) | 1 (0.6%) |
| Flatulence | 7 (4%) | 5 (2.8%) | 5 (2.8%) |
| Liver Function Test Abnormal | 1 (0.6%) | 4 (2.2%) | 2 (1.1%) |
| Alopecia | 0 | 2 (1.1%) | 0 |
| Pruritus | 1 (0.6%) | 2 (1.1%) | 2 (1.1%) |

a: Adverse reactions for which the placebo rate equalled or exceeded the rate for at least one of the LIALDA treatment groups were abdominal pain, dizziness, dyspepsia, and nausea.

The following adverse reactions, presented by body system, were reported infrequently (less than 1%) by LIALDA-treated ulcerative colitis patients in the two controlled trials.

Cardiac Disorder: tachycardia

Vascular Disorders: hypertension, hypotension

Skin and Subcutaneous Tissue Disorders: acne, prurigo, rash, urticaria

Gastrointestinal Disorders: abdominal distention, colitis, diarrhea, pancreatitis, rectal polyp, vomiting

Investigations: decreased platelet count

Musculoskeletal and Connective Tissue Disorders: arthralgia, back pain

Nervous System Disorders: somnolence, tremor

Respiratory, Thoracic and Mediastinal Disorders: pharyngolaryngeal pain

General Disorders and Administrative Site Disorders: asthenia, face edema, fatigue, pyrexia

Ear and Labyrinth Disorders: ear pain

Maintenance of Remission of Ulcerative Colitis

The dose evaluated in three studies of LIALDA given for the maintenance of remission in patients with ulcerative colitis was 1.2 g twice daily or 2.4 g/once daily. One of these studies was a 6-month double-blind comparator study while two were 12- to 14-month open-label studies.

The most common adverse reactions with LIALDA in the maintenance arms of long-term trials were colitis ulcerative (5.8%), headache (2.9%), liver function test abnormal (2.3%), and abdominal pain (2.2%). Of the 1082 subjects in the all maintenance studies pooled, 1.9% had severe adverse reactions. The most common severe adverse reactions were gastrointestinal disorders; these were mainly symptoms associated with ulcerative colitis.

Table 2: Adverse Reactions in Three Maintenance Trials Experienced by at Least 1% of the LIALDA Group (maintenance phases of trials)

| Adverse Reaction | All LIALDA (n=1082) | |
|------------------------------|------------------------|--------|
| | n | % |
| Colitis ulcerative | 63 | (5.8%) |
| Headache | 31 | (2.9%) |
| Liver function test abnormal | 25 | (2.3%) |
| Abdominal pain | 24 | (2.2%) |
| Diarrhea | 18 | (1.7%) |
| Abdominal distension | 14 | (1.3%) |
| Abdominal pain upper | 13 | (1.2%) |
| Dyspepsia | 13 | (1.2%) |
| Back pain | 13 | (1.2%) |
| Rash | 13 | (1.2%) |
| Arthralgia | 12 | (1.1%) |
| Fatigue | 11 | (1.0%) |
| Hypertension | 10 | (1.0%) |

The following adverse reactions, presented by body system, were reported infrequently (less than 1%) by LIALDA-treated ulcerative colitis patients in the three long-term maintenance trials (maintenance phases of these trials):

Cardiac Disorder: tachycardia

Skin and Subcutaneous Tissue Disorders: acne, alopecia, pruritis, urticaria

Gastrointestinal Disorders: colitis, flatulence, nausea, pancreatitis, rectal polyp, vomiting

Nervous System Disorders: dizziness

Respiratory, Thoracic and Mediastinal Disorders: pharyngolaryngeal pain

General Disorders and Administrative Site Disorders: asthenia, pyrexia

Ear and Labyrinth Disorders: ear pain

6.2 Postmarketing Experience

In addition to the adverse reactions reported above in clinical trials involving LIALDA, the adverse reactions listed below have been identified during post-approval use of LIALDA and other mesalamine-containing products. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Body as a Whole: lupus-like syndrome, drug fever

Cardiac Disorders: pericarditis, pericardial effusion, myocarditis

Gastrointestinal: pancreatitis, cholecystitis, gastritis, gastroenteritis, gastrointestinal bleeding, perforated peptic ulcer

Hepatic: jaundice, cholestatic jaundice, hepatitis, liver necrosis, liver failure, Kawasaki-like syndrome including changes in liver enzymes

Hematologic: agranulocytosis, aplastic anemia

Immune System Disorders: anaphylactic reaction, angioedema, Stevens-Johnson syndrome (SJS), drug reaction with eosinophilia and systemic symptoms (DRESS)

Musculoskeletal and Connective Tissue Disorders: myalgia

Neurological/Psychiatric: peripheral neuropathy, Guillain-Barre syndrome, transverse myelitis

Renal Disorders: interstitial nephritis

Respiratory, Thoracic and Mediastinal Disorders: hypersensitivity pneumonitis (including interstitial pneumonitis, allergic alveolitis, eosinophilic pneumonitis)

Skin: psoriasis, pyoderma gangrenosum, erythema nodosum

Urogenital: reversible oligospermia

7 DRUG INTERACTIONS

No investigations of interaction between LIALDA and other drugs except for certain antibiotics have been performed [see *Pharmacokinetics (12.3)*]. However, the following drug-drug interactions have been reported for products containing mesalamine:

7.1 Nephrotoxic Agents, Including Non-Steroidal Anti-Inflammatory Drugs

The concurrent use of mesalamine with known nephrotoxic agents, including non-steroidal anti-inflammatory drugs (NSAIDs) may increase the risk of renal reactions.

7.2 Azathioprine or 6-mercaptopurine

The concurrent use of mesalamine with azathioprine or 6-mercaptopurine may increase the risk for blood disorders.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category B. Reproduction studies with mesalamine have been performed in rats at doses up to 1000 mg/kg/day (1.8 times the maximum recommended human dose based on a body surface area comparison) and rabbits at doses up to 800 mg/kg/day (2.9 times the maximum recommended human dose based on a body surface area comparison) and have revealed no evidence of impaired fertility or harm to the fetus due to mesalamine. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Mesalamine is known to cross the placental barrier.

8.3 Nursing Mothers

Low concentrations of mesalamine and higher concentrations of its N-acetyl metabolite have been detected in human breast milk. The clinical significance of this has not been determined and there is limited experience of nursing women using mesalamine. Caution should be exercised if LIALDA is administered to a nursing woman.

8.4 Pediatric Use

Safety and effectiveness of LIALDA in pediatric patients have not been established.

8.5 Geriatric Use

Reports from uncontrolled clinical studies and postmarketing reporting systems suggested a higher incidence of blood dyscrasias, i.e., neutropenia and pancytopenia in patients who were 65 years or older who were taking mesalamine-containing products such as LIALDA. Caution should be taken to closely monitor blood cell counts during mesalamine therapy.

Clinical trials of LIALDA did not include sufficient numbers of patients aged 65 and over to determine whether they respond differently from younger patients. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. Systemic exposures are increased in elderly subjects. [see *Clinical Pharmacology (12.3)*]. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concurrent disease or other drug therapy in elderly patients.


10 OVERDOSAGE

LIALDA is an aminosalicylate, and symptoms of salicylate toxicity may include tinnitus, vertigo, headache, confusion, drowsiness, sweating, seizures, hyperventilation, dyspnea, vomiting, and diarrhea. Severe intoxication may lead to disruption of electrolyte balance and blood-pH, hyperthermia, dehydration, and end organ damage.

There is no specific known antidote for mesalamine overdose; however, conventional therapy for salicylate toxicity may be beneficial in the event of acute overdosage. Fluid and electrolyte imbalance should be corrected by the administration of appropriate intravenous therapy. Adequate renal function should be maintained.

11 DESCRIPTION

Each LIALDA delayed-release tablet for oral administration contains 1.2 g 5-aminosalicylic acid (5-ASA; mesalamine), an anti-inflammatory agent. Mesalamine also has the chemical name 5-amino-2-hydroxybenzoic acid and its structural formula is:



Molecular formula: C₇H₇NO₃

Molecular weight: 153.14

The tablet is coated with a pH dependent polymer film, which breaks down at or above pH 6.8, normally in the terminal ileum where mesalamine then begins to be released from the tablet core. The tablet core contains mesalamine with hydrophilic and lipophilic excipients and provides for extended release of mesalamine.

The inactive ingredients of LIALDA are sodium carboxymethylcellulose, carnauba wax, stearic acid, silica (colloidal hydrated), sodium starch glycolate (type A), talc, magnesium stearate, methacrylic acid copolymer types A and B, triethylcitrate, titanium dioxide, red ferric oxide and polyethylene glycol 6000.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The mechanism of action of mesalamine is not fully understood, but appears to have a topical anti-inflammatory effect on the colonic epithelial cells. Mucosal production of arachidonic acid metabolites, both through the cyclooxygenase and lipoxygenase pathways, is increased in patients with chronic inflammatory bowel disease, and it is possible that mesalamine diminishes inflammation by blocking cyclooxygenase and inhibiting prostaglandin production in the colon.

Mesalamine has the potential to inhibit the activation of nuclear factor kappa B (NFκB) and consequently the production of key pro-inflammatory cytokines. It has been proposed that reduced expression of PPARγ nuclear receptors (γ-form of peroxisome proliferator-activated receptors) may be implicated in ulcerative colitis. There is evidence that mesalamine produces pharmacodynamic effects through direct activation of PPARγ receptors in the colonic/rectal epithelium.

12.2 Pharmacodynamics

The pharmacodynamic actions of mesalamine occur in the colonic/rectal mucosae local to the delivery of drug from LIALDA into the lumen. There is information suggesting that severity of colonic inflammation in ulcerative colitis patients treated with mesalamine is inversely correlated with mucosal concentrations of mesalamine. Plasma concentrations representing systemically absorbed mesalamine are not believed to contribute extensively to efficacy.

12.3 Pharmacokinetics

Absorption

The total absorption of mesalamine from LIALDA 2.4 g or 4.8 g given once daily for 14 days to healthy volunteers was found to be approximately 21-22% of the administered dose.

Gamma-scintigraphy studies have shown that a single dose of LIALDA 1.2 g (one tablet) passed intact through the upper gastrointestinal tract of fasted healthy volunteers. Scintigraphic images showed a trail of radio-labeled tracer in the colon, suggesting that mesalamine had distributed through this region of the gastrointestinal tract.

In a single dose study, LIALDA 1.2 g, 2.4 g and 4.8 g were administered in the fasted state to healthy

subjects. Plasma concentrations of mesalamine were detectable after 2 hours and reached a maximum by 9-12 hours on average for the doses studied. The pharmacokinetic parameters are highly variable among subjects (Table 3). Mesalamine systemic exposure in terms of area under the plasma concentration-time curve (AUC) was slightly more than dose proportional between 1.2 g and 4.8 g LIALDA. Maximum plasma concentrations (C_{max}) of mesalamine increased approximately dose proportionately between 1.2 g and 2.4 g and sub-proportionately between 2.4 g and 4.8 g LIALDA, with the dose normalized value at 4.8 g representing, on average, 74% of that at 2.4 g based on geometric means.

Table 3: Mean (SD) PK Parameters for Mesalamine Following Single Dose Administration of LIALDA Under Fasting Conditions

| Parameter ¹ of Mesalamine | LIALDA 1.2 g (N=47) | LIALDA 2.4 g (N=48) | LIALDA 4.8 g (N=48) |
|--------------------------------------|------------------------------|--------------------------|----------------------------|
| AUC _{0-t} (ng.h/mL) | 9039 ⁺ (5054) | 20538 (12980) | 41434 (26640) |
| AUC _{0-∞} (ng.h/mL) | 9578 [*] (5214) | 21084 (13185) | 44775 [#] (30302) |
| C_{max} (ng/mL) | 857 (638) | 1595 (1484) | 2154 (1140) |
| T_{max} [*] (h) | 9.0 ^{**} (4.0-32.1) | 12.0 (4.0-34.1) | 12.0 (4.0-34.0) |
| T_{lag} [*] (h) | 2.0 ^{**} (0-8.0) | 2.0 (1.0-4.0) | 2.0 (1.0-4.0) |
| $T_{1/2}$ (h) (Terminal Phase) | 8.56 [*] (6.38) | 7.05 [§] (5.54) | 7.25 [#] (8.32) |

¹ Arithmetic mean of parameter values are presented except for T_{max} and T_{lag} .
^{*}Median (min, max); ⁺N=43, ^{*}N=27, [§]N=33, [#]N=36, ^{**}N=46

Administration of a single dose of LIALDA 4.8 g with a high fat meal resulted in further delay in absorption, and plasma concentrations of mesalamine were detectable 4 hours following dosing. However, a high fat meal increased systemic exposure of mesalamine (mean C_{max} : ↑ 91%; mean AUC: ↑ 16%) compared to results in the fasted state. LIALDA was administered with food in the controlled clinical trials that supported its approval.

In a single and multiple dose pharmacokinetic study of LIALDA, 2.4 g or 4.8 g was administered once daily with standard meals to 28 healthy volunteers per dose group. Plasma concentrations of mesalamine were detectable after 4 hours and were maximal by 8 hours after the single dose. Steady state was achieved generally by 2 days after dosing. Mean AUC at steady state was only modestly greater (1.1- to 1.4-fold) than predictable from single dose pharmacokinetics.

In a single dose pharmacokinetic study of LIALDA, 4.8 g was administered in the fasted state to 71 healthy male and female volunteers (28 young (18-35yrs); 28 elderly (65-75yrs); 15 elderly (>75yrs)). Increased age resulted in increased systemic exposure (approximately 2-fold in C_{max}), to mesalamine and its metabolite N-acetyl-5-aminosalicylic acid. Increased age resulted in a slower apparent elimination of mesalamine, though there was high between-subject variability. Systemic exposures in individual subjects were inversely correlated with renal function as assessed by estimated creatinine clearance.

Table 4: Mean (SD) PK Parameters for Mesalamine Following Single Dose Administration of LIALDA 4.8 g under Fasting Conditions to Young and Elderly Subjects

| Parameter of 5-ASA | Young Subjects (18-35 yrs) (N=28) | Elderly Subjects (65-75 yrs) (N=28) | Elderly Subjects (>75 yrs) (N=15) |
|------------------------------|---|---|---|
| AUC _{0-t} (ng.h/mL) | 51570 (23870) | 73001 (42608) | 65820 (25283) |
| AUC _{0-∞} (ng.h/mL) | 58057 ^b (22429) | 89612 ^c (40596) | 63067 ^d (22531) |
| C_{max} (ng/mL) | 2243 (1410) | 4999 (4381) | 4832 (4383) |
| t_{max} ^a (h) | 22.0 (5.98 - 48.0) | 12.5 (4.00 - 36.0) | 16.0 (4.00 - 26.0) |

| | | | |
|--|--------------------------|--------------------------|--------------------------|
| t _{lag} ^a (h) | 2.00 (1.00 - 6.00) | 2.00 (1.00 - 4.00) | 2.00 (2.00 - 4.00) |
| t _{1/2} (h), terminal phase | 5.68 ^b (2.83) | 9.68 ^c (7.47) | 8.67 ^d (5.84) |
| Renal clearance (L/h) | 2.05 (1.33) | 2.04 (1.16) | 2.13 (1.20) |
| Arithmetic mean (SD) data are presented, N = Number of subjects | | | |
| ^a Median (min-max), ^b N=15, ^c N=16, ^d N=13 | | | |

Distribution

Mesalamine is approximately 43% bound to plasma proteins at the concentration of 2.5 µg/mL.

Metabolism

The only major metabolite of mesalamine (5-aminosalicylic acid) is N-acetyl-5-aminosalicylic acid. Its formation is brought about by N-acetyltransferase (NAT) activity in the liver and intestinal mucosa cells, principally by NAT-1.

Elimination

Elimination of mesalamine is mainly via the renal route following metabolism to N-acetyl-5-aminosalicylic acid (acetylation). However, there is also limited excretion of the parent drug in urine. Of the approximately 21-22% of the dose absorbed, less than 8% of the dose was excreted unchanged in the urine after 24 hours, compared with greater than 13% for N-acetyl-5-aminosalicylic acid. The apparent terminal half-lives for mesalamine and its major metabolite after administration of LIALDA 2.4 g and 4.8 g were, on average, 7-9 hours and 8-12 hours, respectively.

Drug Interactions:

The potential effect of Lialda (4.8 g given once daily) on the pharmacokinetics of four commonly used antibiotics were evaluated in healthy subjects. The four antibiotics studied and their dosing regimens were as follows: amoxicillin (single 500 mg dose), ciprofloxacin XR (single 500 mg dose), metronidazole (750 mg twice daily for 3.5 days), and sulfamethoxazole/trimethoprim (800 mg/160 mg twice daily for 3.5 days). Coadministration of Lialda did not result in clinically significant changes in the pharmacokinetics of any of the four antibiotics. The change in C_{max} and AUC of amoxicillin, ciprofloxacin and metronidazole when they were co-administered with Lialda were all ≤ 3%. There was an increase of 12% in C_{max} and an increase of 15% in AUC of sulfamethoxazole when sulfamethoxazole/trimethoprim was coadministered with Lialda [see *Drug Interactions (7)*].

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

In a 104-week dietary carcinogenicity study in CD-1 mice, mesalamine at doses up to 2500 mg/kg/day was not tumorigenic. This dose is 2.2 times the maximum recommended human dose (based on a body surface area comparison) of LIALDA. Furthermore, in a 104-week dietary carcinogenicity study in Wistar rats, mesalamine up to a dose of 800 mg/kg/day was not tumorigenic. This dose is 1.4 times the recommended human dose (based on a body surface area comparison) of LIALDA.

Mutagenesis

No evidence of mutagenicity was observed in an *in vitro* Ames test or an *in vivo* mouse micronucleus test.

Impairment of Fertility

No effects on fertility or reproductive performance were observed in male or female rats at oral doses of mesalamine up to 400 mg/kg/day (0.7 times the maximum recommended human dose based on a body surface area comparison).

13.2 Animal Toxicology and/or Pharmacology

In animal studies with mesalamine, a 13-week oral toxicity study in mice and 13-week and 52-week oral toxicity studies in rats and cynomolgus monkeys have shown the kidney to be the major target organ of mesalamine toxicity. Oral daily doses of 2400 mg/kg in mice and 1150 mg/kg in rats produced renal lesions including granular and hyaline casts, tubular degeneration, tubular dilation, renal infarct, papillary necrosis, tubular necrosis, and interstitial nephritis. In cynomolgus monkeys, oral daily doses of 250 mg/kg or higher produced nephrosis, papillary edema, and interstitial fibrosis.

14 CLINICAL STUDIES

14.1 Active, Mild to Moderate Ulcerative Colitis

Two similarly designed, randomized, double blind, placebo-controlled trials were conducted in 517 adult patients with active, mild to moderate ulcerative colitis. The study population was primarily Caucasian (80%), had a mean age of 42 years (6% age 65 years or older), and was approximately 50% male. Both studies used LIALDA doses of 2.4 g/day and 4.8 g/day administered once daily for 8 weeks except for the 2.4 g/day group in Study 1, which was given in two divided doses (1.2 g twice daily). The primary efficacy end-point in both trials was to compare the percentage of patients in remission after 8 weeks of treatment for the LIALDA treatment groups versus placebo. Remission was defined as an Ulcerative Colitis Disease Activity Index (UC-DAI) of ≤ 1 , with scores of zero for rectal bleeding and for stool frequency, and a sigmoidoscopy score reduction of 1 point or more from baseline.

In both studies, the LIALDA doses of 2.4 g/day and 4.8 g/day demonstrated superiority over placebo in the primary efficacy endpoint (Table 5). Both LIALDA doses also provided consistent benefit in secondary efficacy parameters, including clinical improvement, treatment failure, clinical remission, and sigmoidoscopic improvement. LIALDA 2.4 g/day and 4.8 g/day had similar efficacy profiles.

Table 5: Patients in Remission at Week 8

| Dose | Study 1 (n=262) n/N (%) | Study 2 (n=255) n/N (%) |
|------------------|-------------------------------|-------------------------------|
| LIALDA 2.4 g/day | 30/88 (34.1) | 34/84 (40.5) |
| LIALDA 4.8 g/day | 26/89 (29.2) | 35/85 (41.2) |
| Placebo | 11/85 (12.9) | 19/86 (22.1) |

14.2 Maintenance of Remission in Patients with Ulcerative Colitis

A multicenter, randomized, double-blind, active comparator study was conducted in a total of 826 adult patients in remission from ulcerative colitis. The study population had a mean age of 45 years (8% age 65 years or older), were 52% male, and were primarily Caucasian (64%).

Maintenance of remission was assessed using a modified Ulcerative Colitis Disease Activity Index (UC-DAI). For this trial, maintenance of remission was based on maintaining endoscopic remission defined as a modified UC-DAI endoscopy subscore of ≤ 1 . An endoscopy subscore of 0 represented normal mucosal appearance with intact vascular pattern and no friability or granulation. For this trial the endoscopy score definition of 1 (mild disease) was modified such that it could include erythema, decreased vascular pattern, and minimal granularity; however, it could not include friability.

Subjects were randomized in a 1:1 ratio to receive either LIALDA 2.4 g/day administered once daily or mesalamine delayed release 1.6 g/day administered as 0.8 g twice daily. The proportion of patients who maintained remission at Month 6 in this study using LIALDA 2.4 g once daily (83.7%) was similar to that seen using the comparator (mesalamine delayed release) 1.6 g/day (81.5%).

16 HOW SUPPLIED/STORAGE AND HANDLING

LIALDA is available as red-brown ellipsoidal film coated delayed-release tablets containing 1.2 g mesalamine, and debossed on one side imprinted with S476.

NDC 54092-476-12 HDPE Bottle with a child-resistant closure of 120 delayed-release tablets.

Store at room temperature 15 °C to 25 °C (59 °F to 77 °F); excursions permitted to 30 °C (86 °F).

See USP Controlled Room Temperature.

17 PATIENT COUNSELING INFORMATION

- Instruct patients not to take LIALDA if they have hypersensitivity to salicylates (e.g., aspirin) or other mesalamines.
- Inform patients to let their physicians know all medications they are taking and if they:
 - are allergic to sulfasalazine, salicylates or mesalamine;
 - are taking non-steroidal anti-inflammatory drugs (NSAIDs) or other nephrotoxic agents;
 - are taking azathioprine, or 6-mercaptopurine;
 - experience cramping, abdominal pain, bloody diarrhea, fever, headache or rash;
 - have a history of myocarditis or pericarditis;
 - have kidney or liver disease;
 - have a history of stomach blockage;
 - are pregnant, intend to become pregnant or are breast-feeding.
- Patients should be instructed to swallow LIALDA delayed-release tablets whole, taking care not to break the outer coating.

Manufactured for Shire US Inc., 300 Shire Way, Lexington, MA 02421, USA by Cosmo S.p.A., Milan, Italy. By license of Nogra Pharma Limited, Dublin, Ireland.

U.S. Patent No. 6,773,720.

© 2015 Shire US Inc.

NDC 54092-476-12 - 1.2g 120 Count Bottle

ONCE-DAILY

NDC 54092-476-12

Lialda[®]

(mesalamine)

delayed release
tablets

1.2 g per tablet

Rx Only

120 Tablets

Shire



NDC 54092-476-08 Display Carton and Sample Carton with Blister Pack

ONCE-DAILY

Lialda®
(mesalamine)
delayed release
tablets
1.2 g per tablet

Rx only

Patient Sample–Not for Sale

3 Cartons of 8 Tablets Each

Shire

NDC 54092-476-08

ONCE-DAILY



Lialda[®]
(mesalamine)
delayed release
tablets
1.2 g per tablet

Rx only

Patient Sample—Not for Sale

Lialda[®]
(mesalamine)
delayed release
tablets

3 Cartons of 8 Tablets Each

Shire
NDC 54092-476-08

Shire

tablets

delayed release
(mesalamine)
Lialda[®]



Active Ingredient: Each tablet contains 1.2 g of mesalamine.

Inactive Ingredients: Carnauba wax, magnesium stearate, methacrylic acid copolymer types A and B, polyethylene glycol 6000, red ferric oxide, silica (colloidal hydrated), sodium carboxymethylcellulose, sodium starch glycolate (type A), stearic acid, talc, titanium dioxide, and triethylcitrate.

SEE PACKAGE INSERT FOR DOSAGE INFORMATION.

KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.

Store at room temperature 15°C to 25°C (59°F to 77°F);
excursions permitted to 30°C (86°F).
See USP Controlled Room Temperature.

www.shire.com



Manufactured for Shire US Inc.
300 Shire Way, Lexington,
MA 02421, USA
© 2015 Shire US Inc.
By license of Nogra Pharma Limited,
Dublin, Ireland
Product of Germany
1610v3

313091


Lialda[®]
(mesalamine)
delayed release
tablets

Shire

PATIENT INSTRUCTIONS

Active Ingredient: Each tablet contains 1.2 g of mesalamine.

Inactive Ingredients: Carnauba wax, magnesium stearate, methacrylic acid copolymer types A and B, polyethylene glycol 6000, red ferric oxide, silica (colloidal hydrated), sodium carboxymethylcellulose, sodium starch glycolate (type A), stearic acid, talc, titanium dioxide, and triethylcitrate.

SEE PACKAGE INSERT FOR DOSAGE INFORMATION.

KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.

Store at room temperature 15°C to 25°C (59°F to 77°F);
excursions permitted to 30°C (86°F).
See USP Controlled Room Temperature.



N 3

54092-476-08

www.shire.com

Manufactured for Shire US Inc.
300 Shire Way, Lexington, MA 02421, USA
© 2015 Shire US Inc.
By license of Nogra Pharma Limited,
Dublin, Ireland
Product of Germany

Shire

ONCE-DAILY



Lialda[®]
(mesalamine)
delayed release
tablets
1.2 g per tablet

Rx Only

Patient Sample—Not for Sale

Lialda[®]
(mesalamine)
delayed release
tablets


Lialda[®]
(mesalamine)
delayed release
tablets

1 blister card containing 8 tablets

Shire

NDC 54092-476-08

Lialda[®]
(mesalamine)
delayed release
tablets



NDC 54092-476-01 - 1.2g 120 Count Patient Sample Bottle

ONCE-DAILY

NDC 54092-476-01

Lialda[®]

(mesalamine)

delayed release

tablets

1.2 g per tablet


Rx Only

120 Tablets

Patient Sample–Not for Sale

Shire

ONCE-DAILY NDC 54092-476-01



Lialda[®]

 (mesalamine)


 delayed release

 tablets


 1.2 g per tablet

Rx Only **120 Tablets**

Patient Sample—Not for Sale



Manufactured for **Shire US Inc.**
 300 Shire Way, Lexington, MA 02421, USA
 © 2015 Shire US Inc.
 By license of Nogra Pharma Limited, Dublin, Ireland
 Product of Germany




**KEEP THIS AND ALL MEDICATION
 OUT OF THE REACH OF CHILDREN.**

Each tablet contains: mesalamine 1.2 g

Pharmacist: Dispense in tight, light-resistant container
 as defined in USP. Store at room temperature
 15°C to 25°C (59°F to 77°F); excursions permitted to
 30°C (86°F). See USP Controlled Room Temperature.

**SEE PACKAGE INSERT FOR
 DOSAGE INFORMATION.**

9395v4 116333



 LOT:
 EXP:

NDC 54092-476-02 - 1.2g Carton of 12 bottles containing 120 tablets each

ONCE-DAILY

NDC 54092-476-02

Lialda[®]

(mesalamine)

delayed release tablets

1.2 g per tablet

Rx only

1 carton of 12 bottles containing 120 tablets each

Shire

ONCE-DAILY

NDC 54092-476-02

ONCE-DAILY


Lialda[®]
 (mesalamine)
 delayed release tablets
 1.2 g per tablet

Rx only


Lialda[®]
 (mesalamine)
 delayed release tablets
 1.2 g per tablet

1 carton of 12 bottles containing 120 tablets each

 Shire

 Shire


Lialda[®]
 (mesalamine)
 delayed release tablets
 1.2 g per tablet

ONCE-DAILY

Active Ingredient: Each tablet contains 1.2 g of mesalamine.

Inactive Ingredients: Carnauba wax, magnesium stearate, methacrylic acid copolymer types A and B, polyethylene glycol 6000, red ferric oxide, silica (colloidal hydrated), sodium carboxymethylcellulose, sodium starch glycolate (type A), stearic acid, talc, titanium dioxide, and triethylcitrate.

SEE PACKAGE INSERT FOR DOSAGE INFORMATION.

KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.

Store at room temperature 15°C to 25°C (59°F to 77°F); excursions permitted to 30°C (86°F).

Manufactured for Shire US Inc.
300 Shire Way, Lexington, MA

ONCE-DAILY


Lialda[®]

See USP Controlled Room Temperature.
www.shire.com
22511v3 313094



MA 02421, USA
© 2015 Shire US Inc.
By license of Nogra Pharma Limited,
Dublin, Ireland
Product of Germany
22511v3

(mesalamine)
delayed release tablets
1.2 g per tablet



Shire

Shire

LIALDA

mesalamine tablet, delayed release

Product Information

| | | | |
|-------------------------|-------------------------|--------------------|---------------|
| Product Type | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:54092-476 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|----------|
| MESALAMINE (UNII: 4Q81I59GXC) (MESALAMINE - UNII:4Q81I59GXC) | MESALAMINE | 1.2 g |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311) | |
| CARNAUBA WAX (UNII: R12CBM0EIZ) | |
| STEARIC ACID (UNII: 4ELV7Z65AP) | |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2) | |
| TALC (UNII: 7SEV7J4R1U) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13) | |
| TRIETHYL CITRATE (UNII: 8Z96QXD6UM) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| FERRIC OXIDE RED (UNII: 1K09F3G675) | |
| POLYETHYLENE GLYCOL 6000 (UNII: 30IQX730WE) | |
| METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:2) (UNII: 5KY68S2577) | |

Product Characteristics

| | | | |
|-------|---------------------------|-------|----------|
| Color | red (red) , brown (brown) | Score | no score |
|-------|---------------------------|-------|----------|

| | | | |
|------------------------------|---|--|---|
| Shape | OVAL (OVAL) | Size | 20mm |
| Flavor | | Imprint Code | S476 |
| Contains | | | |
| Packaging | | | |
| # | Item Code | Package Description | Marketing Start Date Marketing End Date |
| 1 | NDC:54092-476-08 | 3 in 1 CARTON | |
| 1 | | 8 in 1 CARTON | |
| 1 | | 1 in 1 BLISTER PACK; Type 0: Not a Combination Product | |
| 2 | NDC:54092-476-01 | 120 in 1 BOTTLE; Type 0: Not a Combination Product | |
| 3 | NDC:54092-476-12 | 120 in 1 BOTTLE; Type 0: Not a Combination Product | |
| 4 | NDC:54092-476-02 | 12 in 1 CARTON | |
| 4 | NDC:54092-476-01 | 120 in 1 BOTTLE; Type 0: Not a Combination Product | |
| Marketing Information | | | |
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| NDA | NDA022000 | 01/16/2007 | |

Labeler - Shire US Manufacturing Inc. (964907406)

Establishment

| Name | Address | ID/FEI | Business Operations |
|------------------------------|---------|-----------|------------------------------------|
| Sharp Packaging Systems, Inc | | 002346625 | label(54092-476) , pack(54092-476) |

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------------|---------|-----------|------------------------------------|
| Sharp Packaging Systems, Inc. | | 143696495 | label(54092-476) , pack(54092-476) |

Establishment

| Name | Address | ID/FEI | Business Operations |
|-----------------|---------|-----------|--|
| PharmaZell GmbH | | 506639652 | analysis(54092-476) , api manufacture(54092-476) |

Establishment

| Name | Address | ID/FEI | Business Operations |
|-----------|---------|-----------|--|
| Cosmo SpA | | 630431955 | analysis(54092-476) , manufacture(54092-476) , pack(54092-476) |

Establishment

| Name | Address | ID/FEI | Business Operations |
|---------------------------------|---------|-----------|--|
| PharmaZell (Vizag) Private Ltd. | | 915041183 | analysis(54092-476) , api manufacture(54092-476) |