

NAPROXEN SODIUM- naproxen sodium tablet, coated
Kroger Company

NAPROXEN SODIUM
TABLETS USP, 220 mg

PAIN RELIEVER / FEVER REDUCER (NSAID)

Drug Facts

Active ingredient (in each tablet/caplet)

Naproxen sodium USP, 220 mg
(naproxen USP, 200 mg) (NSAID)¹

¹ nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAID's (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech

- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

Adults and children 12 years and older	<ul style="list-style-type: none"> ▪ take 1 tablet/caplet every 8 to 12 hours while symptoms last ▪ for the first dose you may take 2 tablets/caplets within the first hour ▪ do not exceed 2 tablets/caplets in any 8 to 12 hour period ▪ do not exceed 3 tablets/caplets in a 24-hour period
Children under 12 years	<ul style="list-style-type: none"> ▪ ask a doctor

Other information

- **each tablet/caplet contains:** sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

FD &C Blue # 2, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, talc, titanium dioxide

Questions or comments?

call toll-free weekdays (8 AM to 10 PM EST) at **1-888-375-3784**

Distributed by:

The Kroger Co.

Cincinnati, OH 45202

Made in India

PACKAGE LABEL, PRINCIPAL DISPLAY PANEL SECTION

Caplets

COMPARE TO the active ingredient of ALEVE® CAPLETS See side panel**
 NDC 30142-954-20
Kroger
EASY OPEN
Strength to Last 12 Hours
Naproxen Sodium Tablets USP, 220 mg (NSAID)
Pain Reliever/ Fever Reducer
200 CAPLETS (capsule-shaped tablets)
PACKAGE NOT CHILD-RESISTANT

TAMPER EVIDENT: DO NOT USE IF FOIL SEAL UNDER CAP PRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING. IMPORTANT: READ THE DIRECTIONS AND WARNINGS BEFORE USE.

Drug Facts
Active ingredient (in each caplet) Purpose
 Naproxen sodium USP, 220 mg (naproxen USP, 200 mg) (NSAID) Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug

Uses
 Temporarily relieves minor aches and pains due to:
 ■ minor pain of arthritis ■ muscular aches
 ■ backache ■ menstrual cramps
 ■ headache ■ toothache
 ■ the common cold
 ■ temporarily reduces fever

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin.
 Symptoms may include:
 ■ hives ■ asthma (wheezing) ■ facial swelling
 ■ skin redness ■ shock ■ rash ■ dizziness
 ■ difficulty breathing
 If an allergic reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinner (anticoagulant) or steroid drug
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 ■ have 3 or more alcoholic drinks every day while using this product
 ■ take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use
 ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
 ■ right before or after heart surgery

Ask a doctor before use if
 ■ the stomach bleeding warning applies to you
 ■ you have a history of stomach problems, such as heartburn
 ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
 ■ you are taking a diuretic
 ■ you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are
 ■ under a doctor's care for any serious condition
 ■ taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
 ■ taking any other drug

When using this product
 ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if
 ■ you experience any of the following signs of stomach bleeding:
 ■ feel faint
 ■ vomit blood
 ■ have bloody or black stools
 ■ have stomach pain that does not get better

0 41260 01321 8
 DISTRIBUTED BY THE KROGER CO., CINCINNATI, OHIO 45202
For More Product Information, Scan UPC Using Your Kroger App or Call 800-632-6900
 *Aleve® is a registered trademark of Bayer HealthCare LLC, Pittsburgh, PA, 15205. Bayer HealthCare LLC is not affiliated with The Kroger Co. or this product.
 OUR BRANDS. OUR GUARANTEE. LOVE IT OR YOUR MONEY BACK. WWW.KROGER.COM
 LOT: 150084302
 EXP:
 REV 04/21

NAPROXEN SODIUM

naproxen sodium tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:30142-954
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Naproxen Sodium (UNII: 9TN87S3A3C) (Naproxen - UNII:57Y76R9ATQ)	Naproxen Sodium	220 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
hypromelloses (UNII: 3NXW29V3WO)	
magnesium stearate (UNII: 70097M6I30)	
cellulose, microcrystalline (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
povidone (UNII: FZ989GH94E)	
talc (UNII: 7SEV7J4R1U)	
titanium dioxide (UNII: 15FIX9V2JP)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	

Product Characteristics

Color	WHITE	Score	no score
Shape	CAPSULE	Size	12mm
Flavor		Imprint Code	R;273
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30142-954-10	1 in 1 CARTON	08/04/2021	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:30142-954-20	1 in 1 CARTON	08/04/2021	
2		200 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:30142-954-50	1 in 1 CARTON	08/04/2021	
3		500 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075168	08/04/2021	

Labeler - Kroger Company (006999528)

