

GREEN GUARD MAXIMUM NON-ASPIRIN- acetaminophen tablet, film coated
Unifirst First Aid Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Green Guard® Maximum Strength Non-Aspirin

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

For the temporary relief of minor aches and pains associated with

- headache
- toothache
- minor arthritis pain
- muscular aches
- common cold
- menstrual cramps

For the reduction of fever.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if you have

- liver disease

Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin

Stop using and ask a doctor if

- symptoms do not improve
- pain or fever persists or gets worse
- new symptoms occur
- redness or swelling is present

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**

Adults and children: (12 years and older)

Take 2 tablets every 4 to 6 hours as needed. Do not take more than 8 tablets in 24 hours.

Children under 12 years:

Do not give this adult strength product to children under 12 years of age;

this will provide more than the recommended dose (overdose) and may cause liver damage.

Other information

- store at room temperature 59°-86°F (15°-30°C)
- tamper-evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

corn starch, hypromellose, maltodextrin*, microcrystalline cellulose*, polyethylene glycol, povidone*, pregelatinized starch*, sodium starch glycolate*, stearic acid, titanium dioxide*

* may contain

Questions or comments? 1-800-869-6970

Green Guard Max Non-Aspirin Label

Maximum Strength

Non-Aspirin

relief

Compare Active Ingredient To Extra Strength Tylenol®

Green Guard®

Pull Out

Tire Para Abrir

Acetaminophen 500 mg/Acetaminofeno 500 mg

Pain Reliever/Fever Reducer

Alivia el Dolor/Reduce la Fiebre

Tylenol® is a Registered Trademark of McNeil Consumer Products

125 Packets of 2 Tablets

250 Tablets

Order #2222

Distributed by Green Guard® St Louis, MO 63045

An FDA Regulated Facility

<p>Información del medicamento</p> <p>Ingredientes activos (en cada tableta) Acetaminofeno 500mg</p> <p>Propiedad Alivia el dolor/Reduce la Fiebre</p> <p>Usos Para aliviar temporalmente leve dolor y molestias asociadas con: <ul style="list-style-type: none"> dolor de cabeza dolor de muelas dolor de los dientes dolor de la menstruación </p> <p>Advertencias Advertencia con respecto al hígado: Este producto contiene acetaminofeno. Daño grave al hígado pudiera ocurrir si usted consume: <ul style="list-style-type: none"> más de 8 tabletas en 24 horas, la cual es la máxima dosis recomendada este producto con cualquier otro medicamento que contenga acetaminofeno 3 o más bebidas alcohólicas diariamente mientras toma este producto </p> <p>Alerta a las alergias: El Acetaminofeno puede causar severas reacciones alérgicas en la piel. Las cuales pueden incluir: <ul style="list-style-type: none"> enrojecimiento de la piel erupción enrojecido </p> <p>No se administre <ul style="list-style-type: none"> con ningún otro producto que contenga acetaminofeno (recetado o sin receta), si no está seguro si su medicamento contiene acetaminofeno consulte con su médico o farmacéutico por más de 10 días para el dolor o fiebre o por más de 3 días para la fiebre a menos que le indique el médico </p> <p>Consulte con un médico antes de administrarle si: <ul style="list-style-type: none"> está enfermo del hígado está embarazada o en periodo de lactancia, consulte la opinión de un médico antes de consumir este producto </p> <p>Deje de tomarlo y consulte al médico si: <ul style="list-style-type: none"> no mejoran los síntomas aparecen nuevos síntomas el dolor o la fiebre persisten o empeoran se produce hinchazón o enrojecimiento </p> <p>Mantenga esto y todos los medicamentos fuera del alcance de los niños. En caso de ocurrir una sobredosis, comuníquese de inmediato con un médico o llame a un centro de control de envenenamiento. La inmediata atención médica es crítica tanto para los adultos como para los niños sin importar cuál fuere la cantidad o síntomas.</p> <p>Instrucciones no consuma más de lo que se indica</p> <p>Adultos y niños: Tome 2 tabletas cada 4 a 6 horas como sea necesario. No consuma más de 8 tabletas en 24 horas. (de 12 años y mayores)</p> <p>Niños menores de 12 años: No se administre este producto, cuya potencia es para adultos, a niños menores de 12 años de edad, esto le suministraría más de la dosis recomendada (sobredosis) y pudiera causarle daño al hígado.</p> <p>Otra información <ul style="list-style-type: none"> almacenamiento: conserve a temperatura ambiente 59°-86°F (15°-30°C) empacado: con sellado evidente no use ningún paquete abierto o roto </p> <p>Ingredientes Inactivos: almidón de maíz, hipromelosa, metilcelulosa, celulosa microcristalina, polidimetilsiloxano, polidimetilsiloxano, amilón glicosilado, ácido estearico, dióxido de titanio, trióxido de hierro</p> <p>¿Tiene alguna pregunta o comentario? Llame al 1-800-899-6870</p>	<p>MAXIMUM STRENGTH</p> <p>Non-Aspirin</p> <p>relief™</p> <p>COMPARE ACTIVE INGREDIENT TO EXTRA STRENGTH Tylenol®</p> <p>Pull to Open Tire Para Abrir</p> <p>Acetaminophen 500mg/Acetaminofeno 500mg</p> <p>Pain Reliever/Fever Reducer Alivia el Dolor/Reduce la Fiebre</p> <p>125 Packets of 2 Tablets 250 Tablets Order #2222</p> <p>Distributed by Green Guard®, St Louis, MO 63045 An FDA Regulated Facility</p>	<p>Drug Facts</p> <p>Active ingredient (in each tablet) Acetaminophen 500 mg</p> <p>Purpose Pain Reliever/Fever Reducer</p> <p>Uses For the temporary relief of minor aches and pains associated with: <ul style="list-style-type: none"> headache muscular aches minor arthritis pain common cold toothache menstrual cramps </p> <p>Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: <ul style="list-style-type: none"> more than 8 tablets in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using the product </p> <p>Do not use <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. for more than 10 days for pain unless directed by a doctor for more than 3 days for fever unless directed by a doctor </p> <p>Ask a doctor before use if you have: liver disease</p> <p>Stop using and ask a doctor if: <ul style="list-style-type: none"> symptoms do not improve new symptoms occur pain or fever persists or gets worse itchiness or swelling is present </p> <p>Directions <ul style="list-style-type: none"> do not use more than directed Adults and Children (12 years and older): Take 2 tablets every 4 to 6 hours as needed. Do not take more than 8 tablets in 24 hours. Children under 12 years: Do not give this adult strength product to children under 12 years of age, this will provide more than the recommended dose (overdose) and may cause liver damage. </p> <p>Other information <ul style="list-style-type: none"> Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. may contain </p> <p>Inactive ingredients <ul style="list-style-type: none"> corn starch, hypromellose, methylcellulose, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide </p> <p>Questions or comments? 1-800-899-6870</p>	<p>MAXIMUM STRENGTH</p> <p>Non-Aspirin</p> <p>relief™</p> <p>Acetaminophen 500mg/Acetaminofeno 500mg</p> <p>Pain Reliever/Fever Reducer Alivia el Dolor/Reduce la Fiebre</p> <p>125 Packets of 2 Tablets 250 Tablets Order #2222</p> <p>Distributed by Green Guard®, St Louis, MO 63045</p>
---	--	---	---

GREEN GUARD MAXIMUM NON-ASPIRIN

acetaminophen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-009
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

POVIDONE (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white (white)	Score	no score
Shape	ROUND (ROUND)	Size	12mm
Flavor		Imprint Code	AZ;235
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-009-33	50 in 1 BOX, UNIT-DOSE	12/30/2008	
1	NDC:47682-009-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-009-48	125 in 1 BOX, UNIT-DOSE	12/30/2008	
2	NDC:47682-009-99	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	12/30/2008	

Labeler - Unifirst First Aid Corporation (832947092)