

**HEB SPF 50 MANGONADA SCENTED CONTINUOUS SUNSCREEN- avobenzone, homosalate, octisalate, octocrylene spray**

**H.E.B**

-----  
**HEB SPF 50 Mangonada Scented Continuous Sunscreen Spray**

***Active Ingredients***

Avobenzone 3.0%, Homosalate 10.0%, Octisalate 5.0%, Octocrylene 4.0%

***Purpose***

Sunscreen

***Uses***

- helps prevent sunburn
- if used as directed with other sun protection measures (see **Directions**), decreases the risk of skin cancer and early skin aging caused by the sun

***Warnings***

**For external use only**

**FLAMMABLE:**

- do not use near heat, flame or while smoking
- avoid long term storage above 104°F (40°C)

**Do not use**

- on damaged or broken skin.

**When using this product**

- keep out of eyes. Rinse with water to remove.
- do not puncture or incinerate. Contents under pressure.
- do not store at temperatures above 120°F.

**Stop use and ask a doctor if**

- rash occurs.

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

## **Directions**

- apply liberally 15 minutes before sun exposure
- hold can 4-6 inches away from body, spray evenly to ensure complete coverage
- do not spray into face. Spray into hand and apply to face.
- use in well ventilated, but not windy areas
- reapply:
  - after 80 minutes of swimming or sweating
  - immediately after towel drying
  - at least every 2 hours
  - children under 6 months of age: Ask a doctor
  - **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
    - limit time in the sun, especially from 10 a.m. - 2 p.m.
    - wear long-sleeved shirts, pants, hats, and sunglasses

## **Other Information**

- protect the product in this container from excessive heat and direct sun
- may stain or damage some fabrics, materials or surfaces

## **☐ Inactive Ingredients**

acrylates/octylacrylamide copolymer, butylene glycol, butyloctyl salicylate, caprylic/capric triglyceride, fragrance, glycerin, panthenol, saccharomyces/podophyllum peltatum ferment filtrate, SD alcohol 40-B, stearoxytrimethylsilane, tocopherol.

## **Label**

PABA, Retinyl Palmitate & Paraben Free\*  
 No Oxybenzone or Octinoxate  
 Dermatologist Tested

| Drug Facts   |                             |
|--|-----------------------------|
| <b>Active ingredients</b><br>Avobenzone 3%, Homosalate 10%,<br>Octisalate 5%, Octocrylene 4%   | <b>Purpose</b><br>Sunscreen |
| <b>Uses</b> • helps prevent sunburn • If used as directed with other sun protection measures (see <b>Directions</b> ), decreases the risk of skin cancer and early skin aging caused by the sun  |                             |
| <b>Warnings</b><br>For external use only   |                             |
| <b>Flammable:</b><br>• do not use near heat, flame or while smoking<br>• avoid long term storage above 104°F (40°C)  |                             |
| <b>Do not use</b> • on damaged or broken skin,   |                             |
| <b>When using this product</b> • keep out of eyes. Rinse with water to remove. • Keep away from face to avoid breathing it. • do not puncture or incinerate. Contents under pressure. • do not store at temperatures above 120° F.   |                             |
| <b>Stop use and ask a doctor if</b> • rash occurs.   |                             |
| <b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.  |                             |
| <b>Directions</b> • Spray liberally and spread evenly by hand 15 minutes before sun exposure • Hold can 4-6 inches away from the skin to apply • Do not spray directly into face. Spray into hands then apply to face • Do not apply in windy conditions • Use in a well-ventilated area • reapply: • after 80 minutes of swimming or sweating • immediately after towel drying • at least every 2 hours • children under 6 months of age: Ask a doctor • <b>Sun Protection Measures.</b> Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including: • limit time in the sun, especially from 10 a.m. - 2 p.m., • wear long-sleeved shirts, pants, hats, and sunglasses |                             |
| <b>Other information</b><br>• protect the product in this container from excessive heat and direct sun<br>• may stain or damage some fabrics, materials or surfaces  |                             |
| <b>Inactive ingredients</b> acrylates/octylacrylamide copolymer, butylene glycol, butyloctyl salicylate, caprylic/capric triglyceride, fragrance, glycerin, panthenol, saccharomyces/podophyllum peltatum ferment filtrate, SD alcohol 40-B, stearytrimethylsilane, tocopherol.  |                             |
| <b>Questions or comments?</b> Call toll free 1-800-527-7731  |                             |

\*NO PARABENS SEPARATELY ADDED TO PRESERVE THIS PRODUCT  
**MADE WITH PRIDE AND CARE FOR H-E-B®, SAN ANTONIO, TEXAS 78204**



**MANGONADA**  
**50**  
**MANGONADA**  
**SCENTED**

Water Resistant  
 (80 MINUTES)  
 Contains Pro-Vitamin B5  
 & Vitamin E  
 Hypoallergenic  
 UVA/UVB Protection

**BROAD SPECTRUM SPF 50**  
**SCENTED SUNSCREEN**

**NET WT.**  
**5.5 OZ (156 g)**



## HEB SPF 50 MANGONADA SCENTED CONTINUOUS SUNSCREEN

avobenzone, homosalate, octisalate, octocrylene spray

### Product Information

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:37808-172 |
| <b>Route of Administration</b> | TOPICAL        |                           |               |

### Active Ingredient/Active Moiety

| Ingredient Name   | Basis of Strength | Strength      |
|---|-------------------|---------------|
| <b>AVOBENZONE</b> (UNII: G63QQF2NOX) (AVOBENZONE - UNII:G63QQF2NOX) | AVOBENZONE        | 30 mg in 1 g  |
| <b>HOMOSALATE</b> (UNII: V06SV4M95S) (HOMOSALATE - UNII:V06SV4M95S) | HOMOSALATE        | 100 mg in 1 g |

|   |             |              |
|---|-------------|--------------|
| <b>OCTISALATE</b> (UNII: 4X49Y0596W) (OCTISALATE - UNII:4X49Y0596W)   | OCTISALATE  | 50 mg in 1 g |
| <b>OCTOCRYLENE</b> (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM) | OCTOCRYLENE | 40 mg in 1 g |

### Inactive Ingredients

| Ingredient Name  | Strength |
|--|----------|
| <b>ALCOHOL</b> (UNII: 3K9958V90M)  |          |
| <b>ACRYLATE/ISOBUTYL METHACRYLATE/N-TERT-OCTYLACRYLAMIDE COPOLYMER (75000 MW)</b> (UNII: JU3XHR8VWK) |          |
| <b>.ALPHA.-TOCOPHEROL, DL-</b> (UNII: 7QWA1RIO01)  |          |
| <b>GLYCERIN</b> (UNII: PDC6A3C0OX)   |          |
| <b>MEDIUM-CHAIN TRIGLYCERIDES</b> (UNII: C9H2L21V7U)   |          |
| <b>BUTYLENE GLYCOL</b> (UNII: 3XUS85K0RA)  |          |
| <b>BUTYLOCTYL SALICYLATE</b> (UNII: 2EH13UN8D3)  |          |
| <b>PANTHENOL</b> (UNII: WW9CM0O67Z)  |          |
| <b>PODOPHYLLUM</b> (UNII: 2S713A4VP3)  |          |
| <b>STEAROXYTRIMETHYLSILANE</b> (UNII: 9862TW94B2)  |          |

### Packaging

| # | Item Code        | Package Description                               | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:37808-172-37 | 156 g in 1 CAN; Type 0: Not a Combination Product | 01/03/2024           |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M020                                     | 01/03/2024           |                    |

**Labeler** - H.E.B (007924756)

Revised: 3/2025

H.E.B