

NEOSPORIN PLUS PAIN RELIEF- bacitracin zinc, neomycin sulfate, polymyxin b sulfate, and pramoxine hydrochloride ointment
Johnson & Johnson Consumer Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

NEOSPORIN® + PAIN RELIEF OINTMENT

Drug Facts

Active ingredients (in each gram)	Purpose
Bacitracin Zinc (500 units)	First aid antibiotic
Neomycin Sulfate (3.5 mg)	First aid antibiotic
Polymyxin B Sulfate (10,000 units)	First aid antibiotic
Pramoxine HCl (10 mg)	External analgesic

Uses

first aid to help prevent infection and for the temporary relief of pain in minor:

- cuts
- scrapes
- burns

Warnings

For external use only.

Do not use

- if you are allergic to any of the ingredients
- in the eyes
- over large areas of the body

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

Stop use and ask a doctor if

- you need to use longer than 1 week
- condition persists or gets worse
- symptoms persist for more than 1 week, or clear up and occur again within a few days
- rash or other allergic reaction develops

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Centre right away.

Directions

- adults and children 2 years of age and older:
 - clean the affected area
 - apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
 - may be covered with a sterile bandage
- children under 2 years of age: ask a doctor

Other information

- store at 20° to 25°C (68° to 77°F)

Inactive ingredient

Petrolatum

Questions?

call **800-223-0182** or **215-273-8755** (collect)

Distributed by:

**JOHNSON & JOHNSON
CONSUMER INC.**

Skillman, NJ 08558

PRINCIPAL DISPLAY PANEL - 28.3 g Tube Carton

#1

**DOCTOR
RECOMMENDED
BRAND**

NEOSPORIN®

+

PAIN RELIEF

**Maximum Strength
Pain Relief**

**24-Hour Infection
Protection**

**Soothes Cuts, Scrapes
and Burns**

NO STING

OINTMENT

FIRST AID ANTIBIOTIC/PAIN RELIEVING OINTMENT

**Bacitracin Zinc-Neomycin Sulfate-Polymyxin B Sulfate-
Pramoxine HCl**

NET WT 1 oz (28.3 g)

30047870



The makers of NEOSPORIN® do not make store brand products.

how2recycle.info



NEOSPORIN®

#1 DOCTOR RECOMMENDED BRAND



PAIN RELIEF

Maximum Strength Pain Relief

24-Hour Infection Protection

Soothes Cuts, Scrapes, and Burns



OINTMENT

FIRST AID ANTIBIOTIC/PAIN RELIEVING OINTMENT
Bacitracin Zinc - Neomycin Sulfate - Polymyxin B Sulfate - Pramoxine HCl

NET WT. 1 OZ (28.3 g)

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
 Skillman, NJ 08558
 © J&JCI 2020 www.neosporin.com

**Every Cut.
 Every Time.[®]
 Use with
 BAND-AID[®]
 Brand
 Bandages**



Drug Facts

Active ingredients (in each gram)

Bacitracin Zinc (500 units) First aid antibiotic
 Neomycin Sulfate (3.5 mg) First aid antibiotic
 Polymyxin B Sulfate (10,000 units) First aid antibiotic
 Pramoxine HCl (10 mg) External analgesic

Purpose

First aid antibiotic
 First aid antibiotic
 First aid antibiotic
 External analgesic

Uses first aid to help prevent infection and for the temporary relief of pain in minor: ■ cuts ■ scrapes ■ burns

Drug Facts (continued)

Warnings

For external use only.

Do not use

- if you are allergic to any of the ingredients ■ in the eyes
- over large areas of the body

Ask a doctor before use if you have

- deep or puncture wounds ■ animal bites ■ serious burns

Drug Facts (continued)

Stop use and ask a doctor if

- you need to use longer than 1 week
- condition persists or gets worse
- symptoms persist for more than 1 week, or clear up and occur again within a few days
- rash or other allergic reaction develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 2 years of age and older:
- clean the affected area

Drug Facts (continued)

- apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage
- children under 2 years of age: ask a doctor

Other information

- store at 20° to 25°C (68° to 77°F)

Inactive ingredient Petrolatum

Questions?

call 800-223-0182 or 215-273-8755 (collect)

NEOSPORIN[®]

+ PAIN RELIEF

ONTIMENT

EXP
LOT

CODE AREA

NEOSPORIN PLUS PAIN RELIEF

bacitracin zinc, neomycin sulfate, polymyxin b sulfate, and pramoxine hydrochloride ointment

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69968-0057
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BACITRACIN ZINC (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN	500 [USP'U] in 1 g
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN	3.5 mg in 1 g
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B	10000 [USP'U] in 1 g
PRAMOXINE HYDROCHLORIDE (UNII: 88AYB867L5) (PRAMOXINE - UNII:068X84E056)	PRAMOXINE HYDROCHLORIDE	10 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
PETROLATUM (UNII: 4T6H12BN9U)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69968-0057-2	1 in 1 CARTON	12/01/2009	
1		14.2 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:69968-0057-1	1 in 1 CARTON	12/01/2009	
2		28.3 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333B	12/01/2009	

Labeler - Johnson & Johnson Consumer Inc. (118772437)

Revised: 1/2023

Johnson & Johnson Consumer Inc.