

**DG HEALTH OMEPRAZOLE- omeprazole tablet, delayed release  
Dolgenercorp, LLC**

-----  
**Dolgenercorp, LLC Omeprazole Drug Facts**

**Active ingredient (in each tablet)**

Omeprazole 20 mg

**Purpose**

Acid reducer

**Use**

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**

**Allergy alert:** Do not use if you are allergic to omeprazole

**Do not use if you have:**

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have:**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if:**

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea

- you develop a rash or joint pain

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

**14-Day Course of Treatment**

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets

**Repeated 14-Day Courses (if needed)**

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

**Other information**

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

**Inactive ingredients**

carnauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

**Questions or comments?**

**1-888-309-9030**

**Principal Display Panel**

See current Drug Facts

Compare to Prilosec OTC<sup>®</sup>

Omeprazole

Delayed Release Tablets 20 mg

Acid Reducer

Treats Frequent Heartburn!

24 HR

THIS ITEM IS ELECTRONICALLY PROTECTED

20 mg

14 Tablets

One 14-Day Course of Treatment

Actual Tablet Size

May take 1 to 4 days for full effect



A0219

Safety Feature - Do not use if printed tablet blister unit is open or torn.



See current Drug Facts

Compare to Prilosec OTC\*

# Omeprazole

## Delayed Release Tablets 20 mg Acid Reducer

Treats Frequent Heartburn!



THIS ITEM IS ELECTRONICALLY PROTECTED



20 mg



### 14 Tablets

Actual Tablet Size One 14-Day Course of Treatment May take 1 to 4 days for full effect

91574 VT C10

### Drug Facts

Active ingredient (in each tablet)	Purpose
Omeprazole 20 mg	Acid reducer

#### Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

#### Warnings

**Allergy alert:** Do not use if you are allergic to omeprazole

#### Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
  - heartburn with light-headedness, sweating or dizziness
  - chest pain or shoulder pain with shortness of breath, sweating, pains spreading to arms, neck or shoulders; or light-headedness
  - frequent chest pain
- These may be signs of a serious condition. See your doctor.

#### Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

#### Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea

### Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

#### Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

#### 14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

#### Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

#### Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture

**Inactive ingredients** carnauba wax, ferric oxidored, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monochloramine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

100% Satisfaction Guaranteed! (888) 309-9030

MADE IN ISRAEL  
DISTRIBUTED BY DOLGENCORP, LLC  
100 MISSION RIDGE  
GOODLETTSVILLE, TN 37072 USA

\*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Questions or comments? 1-888-309-9030

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
  - Do not wear tight-fitting clothing around the stomach
  - Do not eat before bedtime
  - Raise the head of your bed
  - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
  - Eat slowly and avoid big meals
  - If overweight, lose weight
  - Quit smoking

## DG HEALTH OMEPRAZOLE

omeprazole tablet, delayed release

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:559 10-9 15
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OMEPRAZOLE (UNII: KG60484QX9) (OMEPRAZOLE - UNII:KG60484QX9)	OMEPRAZOLE	20 mg

### Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MONOETHANOLAMINE (UNII: 5KV86114PT)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SODIUM STEARATE (UNII: QU7E2XA9TG)	
SODIUM STEARYL FUMARATE (UNII: 7CV7WJK4U)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

### Product Characteristics

<b>Color</b>	BROWN	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	20
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:559 10-9 15-30	2 in 1 CARTON	06/13/20 11	
1		14 in 1 CARTON		

1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:559 10-9 15-74	14 in 1 CARTON	02/14/20 10	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:559 10-9 15-02	2 in 1 CARTON	05/20/20 12	
3		14 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:559 10-9 15-01	1 in 1 CARTON	07/06/20 13	07/06/20 13
4		14 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:559 10-9 15-03	3 in 1 CARTON	07/24/20 13	
5		14 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA022032	02/14/20 10	

**Labeler** - Dolgencorp, LLC (068331990)

Revised: 5/2019

Dolgencorp, LLC