BENZETHONIUM CHLORIDE AND DYCLONINE HYDROCHLORIDE- liquid bandage liquid
Target Corporation

Target Liquid Bandage

Active ingredients

Benzethonium chloride 0.2%
Dyclonine hydrochloride 0.75%

Purpose

First aid antiseptic
Topical analgesic

Uses

- first aid to help prevent infection in minor cuts, scrapes and burns
- first aid for the temporary relief of pain in minor cuts, scrapes, and burns

Warnings

For external use only.

Flammable:

Keep away from fire or flame.

Do not use

- in the eyes
- over large areas of the body
- longer than 1 week unless directed by a doctor

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

Stop use and ask a doctor if

- conditions persists or gets worse
- infection occurs
- symptoms persist for more than 7 days or clear up and occur again within a few days

Keep out of reach of children.

If swallowed, get medical help or call a Poison Control Center right away.

Directions

- for adults and children 2 years of age and older
- clean the affected area
- apply a small amount of this product on the area 1 to 3 times daily and let dry
- may be covered with a sterile bandage
- if bandaged, let dry first
- children under 2 years of age: do not use, consult doctor

Other information

- **cap bottle tightly**
- wipe top of bottle clean after each use
- store between 20 ^{0} to 25 ^{0}C (68 ^{0} to 77 ^{0}F)
- may temporarily sting upon application
- to remove, apply more Liquid Bandage and quickly wipe off
- do not allow to come into contact with fabrics, floors, countertops, or other surfaces - will stain

Inactive ingredients

acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD slcohol 40

Questions

Call 1-866-964-0939

Principal Display Panel

Liquid Bandage

Benzethonium chloride/First Aid Antiseptic
Dyclonine hydrochloride 0.75%/ Topical Analgesic

Eliminates 99.9% of the three most common

germs within an hour of contact

1 FL. OZ. (29.6 mL)

53-033TG-02

Liquid Bandage

Benzethonium Chloride/First Aid Antiseptic
Dyclonine Hydrochloride 0.75%/Topical Analgesic

Eliminates 99.9% of the three most common germs within an hour of contact†

Protects Small Cuts and Wounds

up&up™

1 FL. OZ. (29.6 mL)

245 07 0553 R00 C-002262-01-035-0000

0 34197 01539 4

Distributed by Target Corporation
Minneapolis, MN 55403
Made in USA with U.S. and imported parts
TM & ©2024 Target Brands, Inc.

Satisfaction guaranteed – Love it or your money back.

We welcome any questions you may have at **Target.com/comments or 1-800-910-6874.**

†Three most common germs used are K. pneumonia, E. coli, P. aeruginosa.

DO NOT USE IF NECKBAND IS BROKEN

Drug Facts

Active ingredients	Purpose
Benzethonium chloride 0.2%	First aid antiseptic
Dyclonine hydrochloride 0.75%	Topical analgesic

Uses
- first aid to help prevent infection in minor cuts, scrapes, and burns
- first aid for the temporary relief of pain in minor cuts, scrapes, and burns

Warnings

For external use only.

Flammable: Keep away from fire or flame.

Do not use
- in the eyes
- over large areas of the body
- longer than 1 week unless directed by a doctor

Ask a doctor before use if you have
- deep or puncture wounds
- animal bites
- serious burns

Stop use and ask a doctor if
- condition persists or gets worse
- infection occurs
- symptoms persist for more than 7 days or clear up and occur again within a few days

Keep out of reach of children. If swallowed, get medical help or call a Poison Control Center right away.

Directions
- for adults and children 2 years of age and older
- clean the affected area
- apply a small amount of this product on the area 1 to 3 times daily and let dry
- may be covered with a sterile bandage
- if bandaged, let dry first
- children under 2 years of age: do not use, consult a doctor

Other information
- **cap bottle tightly**
- wipe top of bottle clean after each use
- store between 20° to 25°C (68° to 77°F)
- may temporarily sting upon application
- to remove, apply more Liquid Bandage and quickly wipe off
- do not allow to come into contact with fabrics, floors, countertops or other surfaces -- will stain.

Inactive ingredients acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

Questions? Call 1-866-964-0939



BENZETHONIUM CHLORIDE AND DYCLONINE HYDROCHLORIDE

liquid bandage liquid

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:82442-019
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZETHONIUM CHLORIDE (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	2 mg in 1 mL
DYCLONINE HYDROCHLORIDE (UNII: ZEC193879Q) (DYCLONINE - UNII:078A24Q30O)	DYCLONINE HYDROCHLORIDE	7.5 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
ACETONE (UNII: 1364PS73AF)	
CASTOR OIL (UNII: D5340Y2I9G)	
ETHYL ACETATE (UNII: 76845O8NMZ)	
PYROXYLIN (UNII: KYR8BR2X6O)	
ALCOHOL (UNII: 3K9958V90M)	
AMYL ACETATE (UNII: 92Q24NH7AS)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82442-019-01	1 in 1 CARTON	04/09/2025	
1		29.6 mL in 1 BOTTLE, WITH APPLICATOR; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	04/09/2025	

Labeler - Target Corporation (006961700)

Revised: 6/2026

Target Corporation