

LOPERAMIDE HYDROCHLORIDE- loperamide hydrochloride tablet
Valu Merchandisers Company

Loperamide Hydrochloride

Drug Facts

Active ingredient (in each caplet)

Loperamide Hydrochloride USP, 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert

Do not use if you have ever had a rash or other allergic reaction to loperamide hydrochloride

Heart alert

Taking more than directed can cause serious heart problems or death

Do not use if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children.In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

| | |
|---------------------------------------|---|
| adults and children 12 years and over | 2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours |
| children 9-11 years (60-95 lbs) | 1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours |
| children 6-8 years (48-59 lbs) | 1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours |
| children 2-5 years (34 to 47 lbs) | ask a doctor |
| children under 2 years (up to 33 lbs) | do not use |

Other information

- store between 20° – 25°C (68° – 77°F)
- see side panel for lot number and expiration date
- **TAMPER EVIDENT: THIS PRODUCT PROTECTED WITH SEALED BLISTER UNITS. DO NOT USE IF ANY ARE TORN OR BROKEN.**

Inactive ingredients

anhydrous lactose, croscarmellose sodium, crospovidone, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

Questions?

Call toll-free Monday-Friday 8:30 am to 5 pm EST at **1800-406-7984**.

PROUDLY DISTRIBUTED BY:
VALU MERCHANDISERS, CO.
5000 KANSAS AVE
KANSAS CITY, KS 66106

PRINCIPAL DISPLAY PANEL - 12 Caplet Blister Pack Carton

COMPARE TO THE ACTIVE
INGREDIENT IN IMODIUM[®]A-D †

Best
Choice[®]

See New Warnings and Directions

Anti-Diarrheal
Loperamide Hydrochloride Tablets
USP, 2 mg

Anti-Diarrheal
Controls the symptoms of diarrhea

12 CAPLETS*

EACH CAPLET (*CAPSULE-SHAPED TABLET)
CONTAINS LOPERAMIDE HYDROCHLORIDE USP, 2 mg

Do not use if you have blood or black stool
 Ask a doctor before use if you have
 ■ fever ■ mucus in the stool
 ■ a history of liver disease
 ■ a history of abnormal heart rhythm
 Ask a doctor or pharmacist before use if you are
 taking a prescription drug. Lopamide may interact
 with certain prescription drugs.
 When using this product, drowsiness or
 dizziness may occur. Be careful when driving or
 operating machinery.

Stop use and ask a doctor if:

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bloating. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center. (1-800-222-1222)

Directions

- Drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- Take right dose on chart, if possible, use weight to dose; otherwise, use age.
- Adults and children 2 capsules after the first loose stool; 1 capsule after each subsequent loose stool, but no more than 4 capsules in 24 hours

| children 6-11 years (45-59 lbs) | children 12-5 years (64 to 47 lbs) | children under 2 years (up to 38 lbs) |
|---|---|--|
| 1 capsule after the first loose stool; ½ capsule after each subsequent loose stool, but no more than 3 capsules in 24 hours | 1 capsule after the first loose stool; ½ capsule after each subsequent loose stool, but no more than 2 capsules in 24 hours | ask a doctor |
| do not use | | |

Other Information

- store between 20° - 25°C (68° - 77°F)
- see side and bottom number and expiration date
- TAMPER EVIDENT: THIS PRODUCT IS PROTECTED WITH SEALED BLISTER UNITS. DO NOT USE IF ANY ARE TORN OR BROKEN.

[illegible]

Active ingredient (in each caplet) Loperamide Hydrochloride USP, 2 mg

Use controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings Allergy alert Do not use if you have ever had a rash or other allergic reaction to hypamide hydrochloride. Heart alert Taking more than directed can cause serious heart problems or death.

Lot No.:

NON VARNISH



Best Choice®

See New Warnings and Directions

Anti-Diarrheal

Controls the symptoms of diarrhea

12 CAPLETS*

EACH CAPLET (* CAPSULE-SHAPED TABLET)
CONTAINS LOPERAMIDE HYDROCHLORIDE USP, 2 mg

Anti-Diarrheal



Keep the carton. It contains important information.

R0419



PROUDLY DISTRIBUTED BY:
VALU MERCHANDISERS, CO.
5000 KANSAS AVE
KANSAS CITY, KS 66106



LOPERAMIDE HYDROCHLORIDE

loperamide hydrochloride tablet

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:63941-123 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|--------------------------|----------|
| LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II) | LOPERAMIDE HYDROCHLORIDE | 2 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK) | |
| CROSCARMELOSE SODIUM (UNII: M28OL1HH48) | |
| CROSPVIDONE (15 MPAS AT 5%) (UNII: 68401960MK) | |
| D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) | |
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | |
| HYDROGENATED COTTONSEED OIL (UNII: Z82Y2C65EA) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| POWDERED CELLULOSE (UNII: SMD1X3XO9M) | |
| STARCH, CORN (UNII: O8232NY3SJ) | |

Product Characteristics

| | | | |
|----------|---------|--------------|----------|
| Color | green | Score | 2 pieces |
| Shape | CAPSULE | Size | 9mm |
| Flavor | | Imprint Code | 123 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:63941-123-12 | 12 in 1 BLISTER PACK; Type 0: Not a Combination Product | 02/01/1993 | |

| | | | | |
|-----------------------|--|---|----------------------|--------------------|
| 2 | NDC:63941-123-24 | 24 in 1 BLISTER PACK; Type U: Not a Combination Product | 02/01/1993 | |
| | | | | |
| Marketing Information | | | | |
| Marketing Category | Application Number or Monograph Citation | | Marketing Start Date | Marketing End Date |
| ANDA | ANDA074091 | | 02/01/1993 | |
| | | | | |

Labeler - Valu Merchandisers Company (868703513)

Registrant - Ranbaxy Pharmaceuticals Inc. (937890044)

| | | | |
|-----------------------|---------|-----------|------------------------|
| Establishment | | | |
| Name | Address | ID/FEI | Business Operations |
| Ohm Laboratories Inc. | | 051565745 | manufacture(63941-123) |

Revised: 1/2024

Valu Merchandisers Company