

SALICYLIC ACID- medicated corn removers patch
Hudson Health LLC

Comfortzone Liquid Bandage

Active ingredients

Benzethonium chloride 0.2%

Dyclonine hydrochloride 0.75%

Purpose

First aid antiseptic

Topical analgesic

Uses

- first aid to prevent bacteria contamination in minor cuts and scrapes

Warnings

For external use only.

Flammable:

do not use while smoking or near heat or flame.

Do not use

- in the eyes
- over large areas of the body
- longer than 1 week unless directed by a doctor
- on infected areas or wounds that are draining
- with other first aid products such as lotions and creams
- over sutures
- on mucous membranes

Ask a doctor before use if you have

- deep puncture wounds
- deep cuts
- animal bites
- serious bleeding
- diabetes
- poor circulation
- serious burns

When using this product

- do not store at temperature above 120°F

Intentional misuse by deliberately concentrating and inhaling the contents can be harmful or fatal.

Stop use and ask a doctor if

- condition persists or gets worse
- infection occurs

Keep out of reach of children.

If swallowed, get medical help or call a Poison Control Center right away.

Directions

- Clean affected area
- apply a small amount on the area 1-3 times daily
- let dry
- a second coating may be applied for extra protection
- to remove, apply more Liquid Bandage and quickly wipe off
- finger nail polish remover may dissolve Liquid Bandage

Other information

- **cap bottle tightly**
- store at room temperature away from heat
- may temporarily sting upon application
- do not allow to come in contact with floors, countertops, or other finished surfaces - will stain

Inactive ingredients

acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

Questions

1-877-237-0194

Principal Display Panel

comfortzone

maximum strength

liquid bandage

benzethonium chloride

dyclonine hydrochloride

topical anesthetic & antiseptic

to help prevent infection

- for small cut & wounds
- antiseptic formula dries quickly to form an invisible protective cover that is flexible

NET 1.0 FL OZ (29.6 mL)

NDC 00000-000-00

DO NOT USE IF NECKBAND IS BROKEN

Active ingredients: Benzethonium Chloride 0.2% and Dyclonine Hydrochloride 0.75% **Purpose:** First aid antiseptic and topical analgesic. **Uses:** First aid to prevent bacterial contamination in minor cuts, scrapes, and burns. For the temporary relief of pain.

Warnings: For external use only. **Flammable:** Keep away from fire or flame. Do not use in eyes, over large areas of the body, and longer than 1 week unless directed by a doctor.


maximum strength
liquid bandage
*benzethonium chloride
dyclonine hydrochloride*

topical analgesic & antiseptic to help prevent infection

- for small cuts & wounds

NET 1.0 FL OZ (29.6 mL)

Ask a doctor before use if you have deep or puncture wounds, deep cuts, animal bites and serious burns.

When using this product avoid contact with eyes.

Stop use and ask a doctor if condition persists or gets worse, symptoms persist for more than 7 days or clear up and occur again within a few days, and infection occurs.

Keep out of reach of children. If swallowed, get medical help or call a Poison Control Center right away.

Directions: For adults and children 2 years of age and older. Clean affected area. Apply a small amount on the area 1 to 3 times daily and let dry. A second coating may be applied for extra protection. Children under 2 years of age: consult a doctor.

Other information: Cap bottle tightly. Store between 20° to 25°C (68° to 77°F). May temporarily sting upon application. To remove, apply more Liquid Bandage and quickly wipe off.

Inactive ingredients: acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

Questions? Call 1-877-237-0194

Distributed By:
Hudson Health LLC
Hudson, NY 12534
©2018

Questions? Call 1-877-237-0194
MADE IN USA WITH U.S. AND IMPORTED PARTS
50-0334M

NDC 00000-000-00

Compare to the active
Ingredient in New-Skin™


maximum strength
liquid bandage
*benzethonium chloride
dyclonine hydrochloride*

topical analgesic & antiseptic to help prevent infection

- for small cuts & wounds
- antiseptic formula dries quickly to form an invisible protective cover that is flexible

NET 1.0 FL OZ (29.6 mL)

USE LIQUID BANDAGE FOR:
blisters, chapped and cracked finger tips, hangnails, paper cuts, shaving nicks and help in preventing the formation of calluses

GREAT FOR:
athletes, active lifestyles, outdoor sportsmen and musicians

50-8334M


maximum strength
liquid bandage



SALICYLIC ACID

medicated corn removers patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72446-005
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZETHONIUM CHLORIDE (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	0.2 mg in 9 mL
DYCLONINE HYDROCHLORIDE (UNII: ZEC193879Q) (DYCLONINE - UNII:078A24Q300)	DYCLONINE HYDROCHLORIDE	0.75 mg in 9 mL

Inactive Ingredients

Ingredient Name	Strength
ACETONE (UNII: 1364PS73AF)	
AMYL ACETATE (UNII: 92Q24NH7AS)	

CASTOR OIL (UNII: D5340Y2I9G)	
ETHYL ACETATE (UNII: 76845O8NMZ)	
PYROXYLIN (UNII: KYR8BR2X6O)	
ALCOHOL (UNII: 3K9958V90M)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72446-005-09	9 mL in 1 BOTTLE, WTH APPLICATOR; Type 0: Not a Combination Product	09/01/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	09/01/2018	

Labeler - Hudson Health LLC (081276171)

Revised: 1/2024

Hudson Health LLC