

LAXATIVE- sennosides tablet, film coated
TOPCO ASSOCIATES LLC

TopCare 44-773

Active ingredient (in each tablet)

Sennosides USP, 25 mg

Purpose

Stimulant laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 6 to 12 hours

Warnings

Do not use

laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor.

Ask a doctor before use if you have

noticed a sudden change in bowel habits that persists over a period of 2 weeks.

Ask a doctor or pharmacist before use if you are

taking any other drug. Laxatives may affect how other drugs work. Take this product 2 or more hours before or after other drugs.

When using this product,

do not use for a period longer than 1 week.

Stop use and ask a doctor if

rectal bleeding or failure to have a bowel movement occur after use of a laxative. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- swallow tablet(s) with a glass of water
- swallow tablet(s) whole; do not crush, break, or chew

adults and children 12 years and over	2 tablets once or twice daily
children 6 to under 12 years	1 tablet once or twice daily
children under 6 years	ask a doctor

Other information

- **each tablet contains:** calcium 35 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, hypromellose, magnesium stearate, medium chain triglycerides, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, silicon dioxide, sodium lauryl sulfate, sucrose, talc, titanium dioxide

Questions or comments?

1-888-423-0139

Principal display panel

+TopCare®

health

NDC 76162-773-08

COMPARE TO EX•LAX®

MAXIMUM STRENGTH

ACTIVE INGREDIENT†

MAXIMUM STRENGTH

Laxative

SENNOSIDES USP, 25 mg • STIMULANT LAXATIVE

- **Senna Ingredient**
- **Gentle, Dependable Constipation Relief**

24 COATED TABLETS

Gentle, Dependable
Constipation Relief

actual
size

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER
UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

†This product is not manufactured or distributed
by GSK Consumer Healthcare SARL, owner of the
registered trademark ex•lax® Maximum Strength.

QUALITY GUARANTEED

50844 ORG082177308

DISTRIBUTED BY
TOPCO ASSOCIATES LLC
ELK GROVE VILLAGE, IL 60007
©TOPCO LNKA0522
QUESTIONS? 1-888-423-0139
topcare@topco.com
www.topcarebrand.com

Other information

- each tablet contains: calcium 35 mg
- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Directions

- do not take more than directed
- swallow tablet(s) with a glass of water
- swallow tablet(s) whole; do not crush, break, or chew

adults and children
2 tablets once or
twice daily

children 6 to under
12 years
1 tablet once or
twice daily

children under 6 years
ask a doctor

Drug Facts (continued)

get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Warnings

Do not use laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor.

Ask a doctor before use if you have noticed a sudden change in bowel habits that persists over a period of 2 weeks.

Ask a doctor or pharmacist before use if you are taking any other drug. Laxatives may affect how other drugs work. Take this product 2 or more hours before or after other drugs.

When using this product, do not use for a period longer than 1 week.

Stop use and ask a doctor if rectal bleeding or failure to have a bowel movement occur after use of a laxative. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose,

Drug Facts (continued)

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 6 to 12 hours

Drug Facts

Active ingredient (in each tablet)

Sennosides USP, 25 mg.....Stimulant laxative

Purpose

Stimulant laxative

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

B-1910-773-08
0FG082177308



MAXIMUM STRENGTH Laxative

SENNOSIDES USP, 25 mg • STIMULANT LAXATIVE

- Senna Ingredient
- Gentle, Dependable Constipation Relief

24 COATED TABLETS

COMPARE TO EX-LAX®
MAXIMUM STRENGTH
ACTIVE INGREDIENT†



Scan here for more information or call 1-888-423-0139



actual size

Drug Facts (continued)

Inactive ingredients corn starch, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, hypromellose, magnesium stearate, medium chain triglycerides, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, silicon dioxide, sodium lauryl sulfate, sucrose, talc, titanium dioxide

Questions or comments? 1-888-423-0139

Drug Facts (continued)

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

†This product is not manufactured or distributed by GSK Consumer Healthcare SARL, owner of the registered trademark ex-lax® Maximum Strength.

No print Area
Lot & Exp. Date

✓ QUALITY GUARANTEED
50844 0FG082177308

DISTRIBUTED BY
TOPCO ASSOCIATES, LLC
ELK GROVE VILLAGE, IL 60007
©TOPCO LNK40522
QUESTIONS? 1-888-423-0139
topcare@topco.com
www.topcarebrand.com



TopCare 44-773

sennosides tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:76162-773
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SENNOSIDES (UNII: 3FYP5M0IJX) (SENNOSIDES - UNII:3FYP5M0IJX)	SENNOSIDES	25 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I3O)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SUCROSE (UNII: C151H8M554)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	blue	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	L7
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76162-773-08	2 in 1 CARTON	06/18/2022	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
-----------	---------------------------------	-----------------	---------------

Category	Citation	Date	Date
OTC Monograph Drug	505G(a)(3)	06/18/2022	

Labeler - TOPCO ASSOCIATES LLC (006935977)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(76162-773)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(76162-773) , pack(76162-773)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(76162-773)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(76162-773)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(76162-773)

Revised: 6/2023

TOPCO ASSOCIATES LLC