

**PHENOBARBITAL, HYOSCYAMINE SULFATE, ATROPINE SULFATE,  
SCOPOLAMINE HYDROBROMIDE - phenobarbital, hyoscyamine sulfate,  
atropine sulfate, scopolamine hydrobromide elixir  
Ingenus Pharmaceuticals, LLC**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----  
**Phenobarbital, Hyoscyamine Sulfate, Atropine Sulfate, Scopolamine  
Hydrobromide Elixir  
Rx Only  
Revised: 05/21**

**DESCRIPTION**

**Phenobarbital, Hyoscyamine Sulfate, Atropine Sulfate, Scopolamine  
Hydrobromide Elixir - Grape**

Each 5 mL (teaspoonful) of elixir (alcohol not more than 23.8%) contains:  
Phenobarbital, USP..... 16.2 mg  
Hyoscyamine Sulfate, USP..... 0.1037 mg  
Atropine Sulfate, USP..... 0.0194 mg  
Scopolamine Hydrobromide, USP.... 0.0065 mg

**Inactive Ingredients**

Purified Water, Glycerin, Sorbitol, Ethyl Alcohol, Sucrose, Saccharin Sodium, Artificial and Natural Grape Flavor, FD&C Red #3, and FD&C Blue #1.

**Phenobarbital, Hyoscyamine Sulfate, Atropine Sulfate, Scopolamine  
Hydrobromide Elixir - Mint**

Each 5 mL (teaspoonful) of elixir (alcohol not more than 23.8%) contains:  
Phenobarbital, USP..... 16.2 mg  
Hyoscyamine Sulfate, USP..... 0.1037 mg  
Atropine Sulfate, USP..... 0.0194 mg  
Scopolamine Hydrobromide, USP.... 0.0065 mg

**Inactive Ingredients**

Purified Water, Glycerin, Sorbitol, Ethyl Alcohol, Sucrose, Saccharin Sodium, Natural Mint Flavor, FD&C Yellow #5, and FD&C Blue #1.

**CLINICAL PHARMACOLOGY**

This drug combination provides natural belladonna alkaloids in a specific, fixed ratio combined with phenobarbital to provide peripheral anticholinergic/antispasmodic action and mild sedation.

**INDICATIONS AND USAGE**

Based on a review of this drug by the National Academy of Sciences-National Research

Council and/or other information, FDA has classified the indications as follows: "Possibly" effective: For use as adjunctive therapy in the treatment of irritable bowel syndrome (irritable colon, spastic colon, mucous colitis) and acute enterocolitis.

May also be useful as adjunctive therapy in the treatment of duodenal ulcer.

Final classification of the less-than-effective indications requires further investigation.

IT HAS NOT BEEN SHOWN CONCLUSIVELY WHETHER ANTICHOLINERGIC/ANTISPASMODIC DRUGS AID IN THE HEALING OF A DUODENAL ULCER, DECREASE THE RATE OF RECURRENCES OR PREVENT COMPLICATIONS.

## **CONTRAINDICATIONS**

- glaucoma;
- obstructive uropathy (for example, bladder neck obstruction due to prostatic hypertrophy);
- obstructive disease of the gastrointestinal tract (as in achalasia, pyloroduodenal stenosis, etc.);
- paralytic ileus, intestinal atony of the elderly or debilitated patient;
- unstable cardiovascular status in acute hemorrhage;
- severe ulcerative colitis especially if complicated by toxic megacolon;
- myasthenia gravis;
- hiatal hernia associated with reflux esophagitis;
- in patients with known hypersensitivity to any of the ingredients.

Phenobarbital is contraindicated in acute intermittent porphyria and in those patients in whom phenobarbital produces restlessness and/or excitement.

## **WARNINGS**

Phenobarbital, Hyoscyamine Sulfate, Atropine Sulfate, Scopolamine Hydrobromide Elixir can cause fetal harm when administered to a pregnant woman. Animal reproduction studies have not been conducted with Phenobarbital, Hyoscyamine Sulfate, Atropine Sulfate, Scopolamine Hydrobromide Elixir. If this drug is used during pregnancy, or if the patient becomes pregnant while taking this drug, the patient should be apprised of the potential hazard to the fetus.

In the presence of a high environmental temperature, heat prostration can occur with belladonna alkaloids (fever and heatstroke due to decreased sweating).

Diarrhea may be an early symptom of incomplete intestinal obstruction, especially in patients with ileostomy or colostomy. In this instance, treatment with this drug would be inappropriate and possibly harmful.

Phenobarbital, Hyoscyamine Sulfate, Atropine Sulfate, Scopolamine Hydrobromide Elixir may produce drowsiness or blurred vision. The patient should be warned, should these occur, not to engage in activities requiring mental alertness, such as operating a motor vehicle or other machinery, and not to perform hazardous work.

Phenobarbital may decrease the effect of anticoagulants, and necessitate larger doses of the anticoagulant for optimal effect. When the phenobarbital is discontinued, the dose of the anticoagulant may have to be decreased.

Phenobarbital may be habit forming and should not be administered to individuals known to be addiction prone or to those with a history of physical and/or psychological dependence upon drugs.

Since barbiturates are metabolized in the liver, they should be used with caution and initial doses should be small in patients with hepatic dysfunction.

## **PRECAUTIONS**

### **General**

Use with caution in patients with:

- autonomic neuropathy
- hepatic or renal disease
- hyperthyroidism
- coronary heart disease
- congestive heart failure
- cardiac arrhythmias
- tachycardia
- hypertension

Belladonna alkaloids may produce a delay in gastric emptying (antral stasis) which would complicate the management of gastric ulcer.

Do not rely on the use of the drug in the presence of complication of biliary tract disease.

Theoretically, with overdosage, a curare-like action may occur.

Phenobarbital, Hyoscyamine Sulfate, Atropine Sulfate, Scopolamine Hydrobromide Elixir – Mint contains FD&C Yellow No. 5 (tartrazine) which may cause allergic-type reactions (including bronchial asthma) in certain susceptible persons. Although the overall incidence of FD&C Yellow No. 5 (tartrazine) sensitivity in the general population is low, it is frequently seen in patients who also have aspirin hypersensitivity.

### **Information for Patients**

Phenobarbital, Hyoscyamine Sulfate, Atropine Sulfate, Scopolamine Hydrobromide Elixir may produce drowsiness or blurred vision. The patient should be warned, should these occur, not to engage in activities requiring mental alertness, such as operating a motor vehicle or other machinery, and not to perform hazardous work.

### **Drug Interactions**

Phenobarbital may decrease the effect of anticoagulants, and necessitate larger doses of the anticoagulant for optimal effect. When the phenobarbital is discontinued, the dose of the anticoagulant may have to be decreased.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Long-term studies in animals have not been performed to evaluate carcinogenic potential.

### **Pregnancy**

Animal reproduction studies have not been conducted with Phenobarbital, Hyoscyamine Sulfate, Atropine Sulfate, Scopolamine Hydrobromide Elixir. There is positive evidence of human fetal risk based on adverse reaction data from investigational or marketing experience or studies in humans, but potential benefits may warrant use of the drug in pregnant women despite potential risks (see *WARNINGS*).

### **Nursing Mothers**

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when Phenobarbital, Hyoscyamine Sulfate, Atropine Sulfate, Scopolamine Hydrobromide Elixir is administered to a nursing woman.

### **Geriatric Use**

Elderly patients may react with symptoms of excitement, agitation, drowsiness, and other untoward manifestations to even small doses of the drug.

## **ADVERSE REACTIONS**

Adverse reactions may include xerostomia; urinary hesitancy and retention; blurred vision; tachycardia; palpitation; mydriasis; cycloplegia; increased ocular tension; loss of taste sense; headache; nervousness; drowsiness; weakness; dizziness; insomnia; nausea; vomiting; impotence; suppression of lactation; constipation; bloated feeling; musculoskeletal pain; severe allergic reaction or drug idiosyncrasies, including anaphylaxis, urticaria, and other dermal manifestations; and decreased sweating.

Acquired hypersensitivity to barbiturates consists chiefly in allergic reactions that occur especially in persons who tend to have asthma, urticaria, angioedema, and similar conditions. Hypersensitivity reactions in this category include localized swelling, particularly of the eyelids, cheeks, or lips, and erythematous dermatitis. Rarely, exfoliative dermatitis (e.g. Stevens-Johnson syndrome and toxic epidermal necrolysis) may be caused by phenobarbital and can prove fatal. The skin eruption may be associated with fever, delirium, and marked degenerative changes in the liver and other parenchymatous organs. In a few cases, megaloblastic anemia has been associated with the chronic use of phenobarbital.

Phenobarbital may produce excitement in some patients, rather than a sedative effect.

**To report SUSPECTED ADVERSE REACTIONS, contact Ingenus Pharmaceuticals, LLC at 1-877-748-1970 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

## **DRUG ABUSE AND DEPENDENCE**

## Abuse

Phenobarbital may be habit forming and should not be administered to individuals known to be addiction prone or to those with a history of physical and/or psychological dependence upon drugs (see *WARNINGS*).

## Dependence

In patients habituated to barbiturates, abrupt withdrawal may produce delirium or convulsions.

## OVERDOSAGE

The signs and symptoms of overdose are headache, nausea, vomiting, blurred vision, dilated pupils, hot and dry skin, dizziness, dryness of the mouth, difficulty in swallowing, and CNS stimulation. Treatment should consist of gastric lavage, emetics, and activated charcoal. If indicated, parenteral cholinergic agents such as physostigmine or bethanechol chloride should be used.

## DOSAGE AND ADMINISTRATION

The dosage of Phenobarbital, Hyoscyamine Sulfate, Atropine Sulfate, Scopolamine Hydrobromide Elixir should be adjusted to the needs of the individual patient to assure symptomatic control with a minimum of adverse effects.

Phenobarbital, Hyoscyamine Sulfate, Atropine Sulfate, Scopolamine Hydrobromide Elixir. Adults: One or two teaspoonfuls of elixir three or four times a day according to conditions and severity of symptoms.

Pediatric patients: may be dosed every 4 to 6 hours. Use a pediatric dosing device or oral syringe to measure the dose.

### Starting Dosage

Body weight	Every 4 hours	Every 6 hours
10 lb. (4.5 kg)	0.5 mL	0.75 mL
20 lb. (9.1 kg)	1 mL	1.5 mL
30 lb. (13.6 kg)	1.5 mL	2 mL
50 lb. (22.7 kg)	2.5 mL	3.75 mL
75 lb. (34 kg)	3.75 mL	5 mL
100 lb. (45.4 kg)	5 mL	7.5 mL

## HOW SUPPLIED

**Phenobarbital, Hyoscyamine Sulfate, Atropine Sulfate, Scopolamine Hydrobromide Elixir - Grape** is a purple colored, grape flavored liquid.

- 4 fl oz (118 mL) bottles- NDC 50742-666-04.
- 1 Pint (473 mL) bottles- NDC 50742-666-16.

**Phenobarbital, Hyoscyamine Sulfate, Atropine Sulfate, Scopolamine Hydrobromide Elixir - Mint** is a green colored, mint flavored liquid.

- 4 fl oz (118 mL) bottles- NDC 50742-665-04.

- 1 Pint (473 mL) bottles- NDC 50742-665-16.

#### Avoid Freezing

Store Phenobarbital, Hyoscyamine Sulfate, Atropine Sulfate, Scopolamine Hydrobromide Elixir at 20° - 25°C (68° - 77°F) [see USP Controlled Room Temperature]. Protect from light and moisture.

Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.

DEA EXEMPT PRODUCT

#### **Manufactured for:**

Ingenus Pharmaceuticals, LLC  
Orlando, FL 32839-6408

Revised: 05/2021  
LF-122298-01  
MPL-100



#### **PACKAGE LABEL.PRINCIPAL DISPLAY PANEL - 4 OZ GRAPE**

NDC 50742-666-04

Phenobarbital, Hyoscyamine Sulfate, Atropine Sulfate, Scopolamine Hydrobromide Elixir Grape Flavored

Each 5 mL (1 teaspoonful) contains:

Phenobarbital, USP..... 16.2 mg  
Hyoscyamine Sulfate, USP..... 0.1037 mg  
Atropine Sulfate, USP..... 0.0194 mg  
Scopolamine Hydrobromide, USP....0.0065 mg

Alcohol not more than 23.8%

**DO NOT USE IF TAMPER-EVIDENT SEAL UNDER CAP IS BROKEN OR MISSING**

Rx Only

4 FL OZ

Ingenus

**Manufactured for:**  
 Ingenus Pharmaceuticals, LLC  
 Orlando, FL 32839-6408

Made in United States  
 LB-122291-01  
 MPL-096

NDC 50742-666-04  
**Phenobarbital,  
 Hyoscyamine Sulfate,  
 Atropine Sulfate,  
 Scopolamine Hydrobromide**  
**Elixir Grape Flavored**

Each 5 mL (1 teaspoonful) contains:  
 Phenobarbital, USP.....16.2 mg  
 Hyoscyamine Sulfate, USP.....0.1037 mg  
 Atropine Sulfate, USP.....0.0194 mg  
 Scopolamine Hydrobromide, USP.....0.0065 mg  
 Alcohol not more than 23.8%

**DO NOT USE IF TAMPER-EVIDENT SEAL  
 UNDER CAP IS BROKEN OR MISSING.**

**Rx Only**  
**4 FL OZ**  


**Warnings: Keep this and all drugs out of reach of children. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.**  
 Store at 20°-25°C (68°-77°F) [see USP Controlled Room Temperature].  
 Avoid freezing.  
 Protect from light and moisture.  
 Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.

**Dosage and Administration:** See accompanying prescribing information.  
 DEA EXEMPT PRODUCT

3  
 50742  
 666 04  
 1

666-04 v1/05-2021

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL - 1 PINT GRAPE**

NDC 50742-666-16

Phenobarbital, Hyoscyamine Sulfate, Atropine Sulfate, Scopolamine Hydrobromide  
 Elixir Grape Flavored

Each 5 mL (1 teaspoonful) contains:

- Phenobarbital, USP..... 16.2 mg
- Hyoscyamine Sulfate, USP..... 0.1037 mg
- Atropine Sulfate, USP..... 0.0194 mg
- Scopolamine Hydrobromide, USP....0.0065 mg
- Alcohol not more than 23.8%

**DO NOT USE IF TAMPER-EVIDENT SEAL UNDER CAP IS BROKEN OR MISSING**

**Bulk Container-Not For Household Use**

Rx Only

1 Pint

Ingenus

Manufactured for:  
Ingenus Pharmaceuticals, LLC  
Orlando, FL 32839-6408

Made in United States

LB-122289-01  
MPL-097

NDC 50742-666-16

# Phenobarbital, Hyoscyamine Sulfate, Atropine Sulfate, Scopolamine Hydrobromide

**Elixir Grape Flavored**

Each 5 mL (1 teaspoonful) contains:  
Phenobarbital, USP.....16.2 mg  
Hyoscyamine Sulfate, USP.....0.1037 mg  
Atropine Sulfate, USP.....0.0194 mg  
Scopolamine Hydrobromide, USP.....0.0065 mg


Alcohol not more than 23.8%

**DO NOT USE IF TAMPER-EVIDENT SEAL  
UNDER CAP IS BROKEN OR MISSING.**

**Bulk Container- Not For Household Use**

**Rx Only**

**1 Pint**



**Warnings: Keep this and all drugs out of reach of children. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.**

Store at 20°-25°C (68°-77°F) [see USP Controlled Room Temperature].

Avoid freezing.

Protect from light and moisture.

Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.

**Dosage and Administration:** See accompanying prescribing information.

DEA EXEMPT PRODUCT

666-16 v1/05-2021



## PACKAGE LABEL.PRINCIPAL DISPLAY PANEL - 4 OZ MINT

NDC 50742-665-04

Phenobarbital, Hyoscyamine Sulfate, Atropine Sulfate, Scopolamine Hydrobromide  
Elixir Mint Flavored

Each 5 mL (1 teaspoonful) contains:

Phenobarbital, USP..... 16.2 mg  
Hyoscyamine Sulfate, USP..... 0.1037 mg  
Atropine Sulfate, USP..... 0.0194 mg  
Scopolamine Hydrobromide, USP....0.0065 mg

Alcohol not more than 23.8%

**DO NOT USE IF TAMPER-EVIDENT SEAL UNDER CAP IS BROKEN OR MISSING**

Rx Only

4 FL OZ

Ingenus




**Manufactured for:**  
 Ingenus Pharmaceuticals, LLC  
 Orlando, FL 32839-6408  
 Made in United States  
 LB-122296-01  
 MPL-098

NDC 50742-665-04  
**Phenobarbital,  
 Hyoscyamine Sulfate,  
 Atropine Sulfate,  
 Scopolamine Hydrobromide**  
**Elixir Mint Flavored**

Each 5 mL (1 teaspoonful) contains:  
 Phenobarbital, USP.....16.2 mg  
 Hyoscyamine Sulfate, USP.....0.1037 mg  
 Atropine Sulfate, USP.....0.0194 mg  
 Scopolamine Hydrobromide, USP.....0.0065 mg  
 Alcohol not more than 23.8%

**DO NOT USE IF TAMPER-EVIDENT SEAL  
 UNDER CAP IS BROKEN OR MISSING.**

**Rx Only**  
**4 FL OZ**  


**Warnings: Keep this and all drugs out of reach of children. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.**  
 Store at 20°-25°C (68°-77°F) [see USP Controlled Room Temperature].  
 Avoid freezing.  
 Protect from light and moisture.  
 Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.  
 Contains color additives including FD&C Yellow No. 5 (tartrazine).  
**Dosage and Administration:** See accompanying prescribing information.  
 DEA EXEMPT PRODUCT

3  
 50742  
 665  
 04  
 4

665-04 v1/05-2021

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL - 1 PINT MINT**

NDC 50742-665-16

Phenobarbital, Hyoscyamine Sulfate, Atropine Sulfate, Scopolamine Hydrobromide  
 Elixir Mint Flavored

Each 5 mL (1 teaspoonful) contains:  
 Phenobarbital, USP..... 16.2 mg  
 Hyoscyamine Sulfate, USP..... 0.1037 mg  
 Atropine Sulfate, USP..... 0.0194 mg  
 Scopolamine Hydrobromide, USP.....0.0065 mg

Alcohol not more than 23.8%

contains color additives including FD&C Yellow No. 5 (tartrazine)

**DO NOT USE IF TAMPER-EVIDENT SEAL UNDER CAP IS BROKEN OR MISSING**

**Bulk Container - Not For Household Use**

Rx Only

1 Pint

Ingenus

Manufactured for:  
Ingenus Pharmaceuticals, LLC  
Orlando, FL 32839-6408

Made in United States

LB-122294-01  
MPL-099

NDC 50742-665-16  
**Phenobarbital,  
Hyoscyamine Sulfate,  
Atropine Sulfate,  
Scopolamine Hydrobromide**

**Elixir Mint Flavored**

Each 5 mL (1 teaspoonful) contains:

Phenobarbital, USP.....16.2 mg  
Hyoscyamine Sulfate, USP.....0.1037 mg  
Atropine Sulfate, USP.....0.0194 mg  
Scopolamine Hydrobromide, USP.....0.0065 mg

Alcohol not more than 23.8%  
contains color additives including  
FD&C Yellow No. 5 (tartrazine)

**DO NOT USE IF TAMPER-EVIDENT SEAL  
UNDER CAP IS BROKEN OR MISSING.**

**Bulk Container- Not For Household Use**

**Rx Only**

**1 Pint**

ingenus

**Warnings: Keep this and all drugs  
out of reach of children. In case of  
accidental overdose, seek  
professional assistance or contact a  
Poison Control Center immediately.**

Store at 20°-25°C (68°-77°F) [see  
USP Controlled Room Temperature].

Avoid freezing.


Protect from light and moisture.

Dispense in a tight, light-resistant  
container as defined in the USP  
using a child-resistant closure.

**Dosage and Administration:** See  
accompanying prescribing  
information.

DEA EXEMPT PRODUCT

665-16 v1/05-2021



## PHENOBARBITAL, HYOSCYAMINE SULFATE, ATROPINE SULFATE, SCOPOLAMINE HYDROBROMIDE

phenobarbital, hyoscyamine sulfate, atropine sulfate, scopolamine hydrobromide elixir

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:50742-666
<b>Route of Administration</b>	ORAL	<b>DEA Schedule</b>	CIV

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>PHENOBARBITAL</b> (UNII: YQE403BP4D) (PHENOBARBITAL - UNII:YQE403BP4D)	PHENOBARBITAL	16.2 mg in 5 mL
<b>HYOSCYAMINE SULFATE</b> (UNII: F2R8V82B84) (HYOSCYAMINE - UNII:PX44XO846X)	HYOSCYAMINE SULFATE	0.1037 mg in 5 mL
<b>ATROPINE SULFATE</b> (UNII: 03J5ZE7KA5) (ATROPINE - UNII:7C0697DR9I)	ATROPINE SULFATE	0.0194 mg in 5 mL
<b>SCOPOLAMINE HYDROBROMIDE</b> (UNII: 4511FR0GXB) (SCOPOLAMINE - UNII:DL48G20X8X)	SCOPOLAMINE HYDROBROMIDE	0.0065 mg in 5 mL

Inactive Ingredients	
Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>FD&amp;C RED NO. 3</b> (UNII: PN2ZH5LOQY)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	

Product Characteristics			
<b>Color</b>	PURPLE	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	GRAPE (Artificial and Natural Grape)	<b>Imprint Code</b>	
<b>Contains</b>			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50742-666-04	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/01/2021	
2	NDC:50742-666-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/01/2021	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
UNAPPROVED DRUG OTHER		08/01/2021	

## PHENOBARBITAL, HYOSCYAMINE SULFATE, ATROPINE SULFATE, SCOPOLAMINE HYDROBROMIDE

phenobarbital, hyoscyamine sulfate, atropine sulfate, scopolamine hydrobromide elixir

Product Information			
<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:50742-665
<b>Route of Administration</b>	ORAL	<b>DEA Schedule</b>	CIV

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
<b>PHENOBARBITAL</b> (UNII: YQE403BP4D) (PHENOBARBITAL - UNII:YQE403BP4D)	PHENOBARBITAL	16.2 mg in 5 mL

<b>HYOSCYAMINE SULFATE</b> (UNII: F2R8V82B84) (HYOSCYAMINE - UNII:PX44XO846X)	HYOSCYAMINE SULFATE	0.1037 mg in 5 mL
<b>ATROPINE SULFATE</b> (UNII: 03J5ZE7KA5) (ATROPINE - UNII:7C0697DR9I)	ATROPINE SULFATE	0.0194 mg in 5 mL
<b>SCOPOLAMINE HYDROBROMIDE</b> (UNII: 451IFR0GXB) (SCOPOLAMINE - UNII:DL48G20X8X)	SCOPOLAMINE HYDROBROMIDE	0.0065 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>FD&amp;C YELLOW NO. 5</b> (UNII: I753WB2F1M)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	

### Product Characteristics

<b>Color</b>	GREEN	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	MINT (Natural Mint)	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50742-665-04	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/01/2021	
2	NDC:50742-665-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/01/2021	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
UNAPPROVED DRUG OTHER		08/01/2021	

**Labeler** - Ingenus Pharmaceuticals, LLC (833250017)