

OMEPRAZOLE- omeprazole tablet, delayed release
Twin Med LLC

Omeprazole

Drug Facts

Active ingredient (in each tablet)

Omeprazole USP 20 mg

Purpose

Acid reducer

Use

- treats frequent heartburn (occurs ***2 or more*** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert:

- do not use if you are allergic to omeprazole.
- omeprazole may cause severe skin reactions.
Symptoms may include:
 - skin reddening
 - blisters
 - rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20 to 25° C (68 to 77° F) and protect from moisture

Inactive ingredients

anhydrous lactose, hypromellose, hypromellose acetate succinate, iron oxide red, iron oxide yellow, lactose monohydrate, methyl cellulose, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, triethyl citrate and titanium dioxide.

The imprinting ink contains ammonium hydroxide, black iron oxide, n-butyl alcohol, propylene glycol and shellac.

Questions or Comments?

Call toll free 1-800-818-4555 weekdays.

Distributed by:

ProCure Products

Santa Fe Springs, CA 90670

PRINCIPAL DISPLAY PANEL - 20 mg Tablet Bottle Carton

SWALLOW - DO NOT CHEW

NDC 55681-306-42

PROCURE™

HEALTHCARE PRODUCTS

Compare to active ingredient of Prilose OTC®*

OMEPRAZOLE

20 mg / ACID REDUCER

DELAYED-RELEASE TABLETS

actual size

PROCURE PRODUCTS

SYMBOL OF EXCELLENCE

24

HR

Treats frequent heartburn!

See new warning information

42 TABLETS

THREE 14-DAY COURSES OF TREATMENT

MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT

PROCURE™
HEALTHCARE PRODUCTS

OMEPRAZOLE

20 MG / ACID REDUCER
DELAYED-RELEASE TABLETS

42 TABLETS
THREE 14-DAY COURSE OF TREATMENT
MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT

SWALLOW - DO NOT CHEW

ISS: 12/2022

PROCURE™
HEALTHCARE PRODUCTS

OMEPRAZOLE

20 MG / ACID REDUCER
DELAYED-RELEASE TABLETS
3 BOTTLES INSIDE

Safety Feature - Do not use if printed seal under cap is broken or missing.




Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Distributed by:
ProCure Products
Santa Fe Springs, CA 90670
Made in India

SWALLOW - DO NOT CHEW

NDC 55881-306-42



Compare to active ingredient of Prilosec OTC®

PROCURE™
HEALTHCARE PRODUCTS

OMEPRAZOLE

20 mg / ACID REDUCER
DELAYED-RELEASE TABLETS



actual size



Treats frequent heartburn!

See new warning information

42 TABLETS

THREE 14-DAY COURSES OF TREATMENT

MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT



8 16516 01383 6

DNH/DRUGS/NH/138

5 2 3 6 5 1 6

**Unvarnish Area:
44x33 mm**

*All trademarks are the property of their respective owner. This product is not affiliated with the maker/owner of Prilosec OTC®.

Glue-NO COATING

Drug Facts

Active ingredient (in each tablet)
Omeprazole USP 20 mg Acid reducer

Use
■ treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert:
■ do not use if you are allergic to omeprazole.
■ omeprazole may cause severe skin reactions.
Symptoms may include:
■ skin redness ■ hives ■ rash
If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
■ heartburn with lightheadedness, sweating or dizziness
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
■ frequent chest pain
These may be signs of a serious condition. See your doctor.

Drug Facts (continued)

Ask a doctor before use if you have:
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting
■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and seek a doctor if:
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea
■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
■ for adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours
14-Day Course of Treatment
■ swallow 1 tablet with a glass of water before eating in the morning

Drug Facts (continued)

■ take every day for 14 days
■ do not take more than 1 tablet a day
■ do not use for more than 14 days unless directed by your doctor
■ swallow whole. Do not chew or crush tablets.
Repeated 14-Day Courses (if needed)
■ you may repeat a 14-day course every 4 months
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor
■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other Information
■ read the directions and warnings before use
■ keep the carton. It contains important information.
■ store at 20 to 25° C (68 to 77° F) and protect from moisture

Inactive ingredients anhydrous lactose, hydroxypropylcellulose, hydroxypropylcellulose acetate succinate, iron oxide red, iron oxide yellow, lactose monohydrate, methylcellulose, monobasic aluminum phosphate, polyethylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, toluidine blue, triethyl citrate and titanium dioxide.
The imprinting ink contains ammonium hydroxide, black iron oxide, n-butyl alcohol, propylene glycol and shellac.

Questions or Comments?
Call toll free 1-800-918-4555 weekdays.

Glue- NO COATING

OMEPRAZOLE

omeprazole tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55681-306
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OMEPRAZOLE (UNII: KG60484QX9) (OMEPRAZOLE - UNII:KG60484QX9)	OMEPRAZOLE	20 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
HYPROMELLOSE ACETATE SUCCINATE 12070923 (3 MM2/S) (UNII: 36BGF0E889)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MONOETHANOLAMINE (UNII: 5KV86114PT)	
METHYLCELLULOSE (1500 MPA.S) (UNII: PONTE48364)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
SODIUM STEARATE (UNII: QU7E2XA9TG)	
SODIUM STEARYL FUMARATE (UNII: 7CV7WJK4UI)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
AMMONIA (UNII: 5138Q19F1X)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
BUTYL ALCOHOL (UNII: 8PJ61P6TS3)	
SHELLAC (UNII: 46N107B71O)	

Product Characteristics

Color	BROWN (brownish pink)	Score	no score
-------	-----------------------	-------	----------

Shape	OVAL (biconvex)		Size	12mm
Flavor			Imprint Code	20
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55681-306-14	1 in 1 CARTON	04/14/2022	
1		14 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:55681-306-42	3 in 1 CARTON	04/14/2022	
2		14 in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA207891		04/14/2022	

Labeler - Twin Med LLC (009579330)

Establishment			
Name	Address	ID/FEI	Business Operations
Sun Pharmaceutical Industries Limited		650445203	MANUFACTURE(55681-306)

Revised: 1/2023

Twin Med LLC