

BLISTER BLASTER- benzalkonium chloride liquid
Quadex Pharmaceuticals, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

blisterblaster®

Drug Facts

Active ingredients

Benzalkonium Chloride 0.13%

Purposes

Cold Sore/Fever Blister Treatment Topical Antiseptic

Uses

- To treat cold sores/fever blisters.
- To help guard against infection.

Warning

- For external use only.
- Flammable; keep away from fire or flame.

Do not use:

- in the eyes.
- over large areas of the body.
- longer than one week unless directed by a doctor.
- more than 3 times per day.
- if you are allergic to any ingredient in this product.

Stop use and ask a doctor if:

- condition persists or worsens.
- symptoms persist for more than 7 days.

Ask a doctor if:

- used to treat deep or puncture wounds, animal bites, or serious burns.
- you are pregnant or nursing a baby.

When using this product you may feel a brief stinging sensation when you apply it. The sting should go away in a short time.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions - Adults and children over 2 years of age

- Using only warm water and a washcloth, clean the lip area of any lip preparations, lotions, ointments, residual beverages, or cosmetics, including lipstick.
- Remove the cardboard cap from the applicator end and place the cardboard cap onto the glass/plastic

vial end, opposite the brush end of the product.

- Squeeze the cardboard cap to break open the inner glass vial.
- Saturate the applicator end with solution by holding the brush end down and squeezing the container until you can see liquid on the brush applicator.
- For best results, massage the solution into the cold sore by rubbing. Rub firmly, but take care not to damage the tissue. The purpose of the rubbing is to deliver the drug to the infection site. Hold the vial so the solution flows to the sore.
- To treat most cold sores, usually one treatment is enough. If your symptoms go away and then return later, apply another dose for the second cold sore.
- Do not use more than 3 times per day.
- Children under 2 years of age — ask a doctor.
- Discard after use.

Other information

Store at room temperature. The ingredients in toothpaste, soft drinks, and some fruit juices can deactivate the active ingredient in Blister Blaster. For best results, avoid brushing your teeth with toothpaste or drinking soft drinks or fruit juices for at least one hour after applying the drug.

Inactive ingredients

Isopropyl alcohol (70% v/v), water.

Questions or to report an adverse event: 1-877-825-7153

PRINCIPAL DISPLAY PANEL - 0.6 ml Vial Blister Pack

NDC 66774-1001-2

blisterblaster®

**One Day,
One Dose,
Healing Begins.**

*Finally, a cold sore treatment that actually works—so you can clear away that **THING** on your lip right now!*

Relieves Pain and Burning

Not available through retail stores; avoid imitations.

Contents: 2 Vials (0.6 ml)

NDC 66774-1001-2

blisterblaster® 



**One Day,
One Dose,
Healing Begins.**

*Finally, a cold sore treatment that actually works—so you can clear away that **THING** on your lip right now!*

Relieves Pain and Burning

Not available through retail stores; avoid imitations.

Contents: 2 Vials (0.6 ml)

**Cold Sore/Fever Blister
Treatment & Topical Antiseptic**

Drug Facts

Active ingredients

Benzalkonium Chloride 0.13%

Purposes

Cold Sore/Fever Blister
Treatment Topical Antiseptic

Uses

- To treat cold sores/fever blisters.
- To help guard against infection.

Warning

- For external use only.
- Flammable; keep away from fire or flame.

Do not use:

- in the eyes.
- longer than one week unless directed by a doctor.
- if you are allergic to any ingredient in this product.
- over large areas of the body.
- more than 3 times per day.

Stop use and ask a doctor if:

- condition persists or worsens.
- symptoms persist for more than 7 days.

Ask a doctor if:

- used to treat deep or puncture wounds, animal bites, or serious burns.
- you are pregnant or nursing a baby.

When using this product you may feel a brief stinging sensation when you apply it. The sting should go away in a short time.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions - Adults and children over 2 years of age.

- Using only warm water and a washcloth, clean the lip area of any lip preparations, lotions, ointments, residual beverages, or cosmetics, including lipstick.
- Remove the cardboard cap from the applicator end and place the cardboard cap onto the glass/plastic vial end, opposite the brush end of the product.
- Squeeze the cardboard cap to break open the inner glass vial.
- Saturate the applicator end with solution by holding the brush end down and squeezing the container until you can see liquid on the brush applicator.
- For best results, massage the solution into the cold sore by rubbing. Rub firmly, but take care not to damage the tissue. The purpose of the rubbing is to deliver the drug to the infection site. Hold the vial so the solution flows to the sore.
- To treat most cold sores, usually one treatment is enough. If your symptoms go away and then return later, apply another dose for the second cold sore.
- Do not use more than 3 times per day.
- Children under 2 years of age – ask a doctor.
- Discard after use.

Other information

Store at room temperature. The ingredients in toothpaste, soft drinks, and some fruit juices can deactivate the active ingredient in Blister Blaster. For best results, avoid brushing your teeth with toothpaste or drinking soft drinks or fruit juices for at least one hour after applying the drug.

Inactive ingredients:

Isopropyl alcohol (70% v/v), water.

**Questions or to report an
adverse event: 1-877-825-7153**



Quadex Pharmaceuticals, LLC
7070 Union Park Center #380
Salt Lake City, Utah 84047

Made in the U.S.A.

www.BlisterBlaster.com

Cold Sore/Fever Blister Treatment & Topical Antiseptic

Did You Know:

A cold sore is caused by the herpes virus (Typically HSV-1). "Cold sore" and "fever blister" are just different popular names for the same lesion, which is caused by the herpes simplex virus.

Herpes is highly contagious.

- A kiss can transfer the virus to another individual.
- Children are highly at risk.

After touching your cold sore:

- Wash your hands with warm water and soap.
- Do not touch your eyes.
- Do not touch your genitals or other parts of the body.
It is possible to transfer the virus to other parts of the body.

Additional Treatments:

- Occasionally a lesion will start healing and then appear to recur. This is typically caused by a new viral release, and will require an additional treatment for the second lesion.
- Deactivation of the active ingredient, caused by toothpaste/mouthwash, soft drinks, soaps/detergents, and some fruit juices, may require an additional treatment.

How It Works:

Blister Blaster[®] is a patented approach to treating cold sores/fever blisters. Rubbing the cold sore with the applicator tip delivers the germicidal medicine to the site of the infection while eliminating the need to touch the sore with your fingers.

blister blaster 

Quadex Pharmaceuticals, LLC
7070 Union Park Center #380
Salt Lake City, Utah 84047

US Patents: 6,758,434 6,423,750 6,211,243 6,410,599 6,414,032 6,420,431
Other US and foreign patents issued and pending.

Quadex Pharmaceuticals, LLC
7070 Union Park Center #380
Salt Lake City, Utah 84047

Made in the U.S.A.

www.BlisterBlaster.com

blisterblaster®



Instructions:

Apply Blister Blaster® as soon as you feel the tingly, itchy sensation of a cold sore beginning.

Clean it

Prior to treatment, clean the area to be treated. Do not use soap or other cleansers. A dry wipe maybe sufficient; you can use water or alcohol, if necessary.



Break it

Remove the paper cap and place it on the other end, over the clear plastic tube. Pinch the vial in the center of the blue paper cap until the inner glass vial breaks.

Hold with the white applicator tip down and allow medication to saturate the swab.

If necessary: pinch the vial gently until a drop of medication just appears.

When finished, recap the vial and dispose immediately. Do not disassemble. Keep out of the reach of children.



Always carry Blister Blaster® with you. The sooner you apply, the sooner healing begins. Minutes can make a difference.

Rub it

The key is to **vigorously rub** on the tingly, itching site of the cold sore and the surrounding area. Rub firmly, but take care not to damage the tissue.

On Open Sores:

Vigorously rub the surrounding area.
Then, firmly press the applicator tip to the cold sore and rub, moving the lip against the teeth.
Avoid scrubbing; this will further open the lesion.

Keep the applicator saturated at all times. If necessary, pause and hold the vial so as to allow the medication to flow into the applicator.

Use all the medication in the vial.

Additional Treatments?

Yes, if the burning and pain returns:

- Because of a new viral release, or
- After deactivation of the active ingredient by contact with soap, toothpaste, or fruit juice.



Always carry Blister Blaster[®] with you.
The sooner you apply, the sooner healing begins. Minutes can make a difference.

CLEAN IT
wipe your lips

BREAK IT
break the vial

RUB IT
massage lips



Quadex Pharmaceuticals, LLC
7070 Union Park Center #380
Salt Lake City, Utah 84047
www.BlisterBlaster.com
1-800-398-9924

COUPON



Go to BlisterBlaster.com to order your next supply of Blister Blaster.
Enter this Offer Code P0005 and get the same price as your last order.

That would be two vials for just \$19.95 plus shipping and handling.

Quadex Pharmaceuticals, LLC • 7070 Union Park Center #380 • Salt Lake City, Utah 84047 • www.BlisterBlaster.com • 1-800-398-9924

CLEAN IT
wipe your lips

BREAK IT
break the vial

RUB IT
massage lips



Quadex Pharmaceuticals, LLC
7070 Union Park Center #380
Salt Lake City, Utah 84047
www.BlisterBlaster.com
1-800-398-9924

BLISTER BLASTER

benzalkonium chloride liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66774-1001
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Benzalkonium chloride (UNII: F5UM2KM3W7) (Benzalkonium - UNII:7N6JUD5X6Y)	Benzalkonium chloride	0.12 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
Isopropyl Alcohol (UNII: ND2M416302)	70 mL in 100 mL
Water (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66774-1001-2	2 in 1 BLISTER PACK		
1		0.6 mL in 1 VIAL		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part333A	04/01/2010	

Labeler - Quadex Pharmaceuticals, LLC (090438909)

Establishment

Name	Address	ID/FEI	Business Operations
James Alexander Corporation		040756421	MANUFACTURE

Revised: 7/2011

Quadex Pharmaceuticals, LLC