

ASPIRIN- aspirin tablet, film coated
L.N.K. International, Inc.

Quality Plus 44-157

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves:

- minor pain of arthritis
- headache
- pain and fever of colds
- toothache
- menstrual pain
- muscle pain

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- hives
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take more or for a longer time than directed
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you are taking a diuretic
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

corn starch, hypromellose, polyethylene glycol, propylene glycol

Questions or comments?

1-800-426-9391

Principal display panel

**QUALITY
+PLUS**

NDC 50844-157-11

†Compare to the
active ingredient in
Genuine Bayer ® Aspirin

ASPIRIN
PAIN RELIEVER/
FEVER REDUCER
(NSAID)

325 mg

**60
Tablets**

Talk to your doctor
or other healthcare
provider before
using this product
for your heart.

ACTUAL SIZE

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed by Bayer AG, owner of the
registered trademark Genuine Bayer® Aspirin. 50844 ORG012215711

Distributed by
LNK INTERNATIONAL, INC.
60 Arkay Drive
Hauppauge, NY 11788
USA



NDC 50844-157-11

†Compare to the active ingredient in Genuine Bayer® Aspirin

ASPIRIN

PAIN RELIEVER/ FEVER REDUCER (NSAID)

325 mg



ACTUAL SIZE

60 Tablets

Talk to your doctor or other healthcare provider before using this product for your heart.

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves:
 - minor pain of arthritis
 - headache
 - pain and fever of colds
 - toothache
 - menstrual pain
 - muscle pain

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
 - hives
 - shock
 - asthma (wheezing)
- Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
- are age 60 or older
 - have had stomach ulcers or bleeding problems
 - take a blood thinning (anticoagulant) or steroid drug
 - take more or for a longer time than directed
 - take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

No print/No varnish
Lot & Exp date

This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Genuine Bayer® Aspirin.

Distributed by

LINK INTERNATIONAL, INC.

60 A-ckey Drive
Hauppauge, NY 11788
USA



0 50844 15711 8

PEEL HERE FOR MORE DRUG FACTS

Adhesive Only

STOP PEELING

Drug Facts (continued)

■ have 3 or more alcoholic drinks every day while using this product

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer

■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ stomach bleeding warning

applies to you ■ you are taking a diuretic

■ you have a history of stomach problems, such as heartburn

■ you have high blood pressure, heart disease, liver cirrhosis,

or kidney disease ■ you have asthma

Ask a doctor or pharmacist before use if you are taking a

prescription drug for ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if

■ an allergic reaction occurs. Seek medical help right away.

■ you experience any of the following signs of stomach bleeding:

■ feel faint ■ vomit blood ■ have bloody or black stools

■ have stomach pain that does not get better

■ ringing in the ears or a loss of hearing occurs

■ pain gets worse or lasts more than 10 days

■ fever gets worse or lasts more than 3 days

■ new symptoms occur ■ redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or

later in pregnancy unless definitely directed to do so by a doctor

because it may cause problems in the unborn child or

complications during delivery.

Keep out of reach of children. In case of overdose, get medical

help or contact a Poison Control Center right away.

Directions ■ do not take more than directed

■ drink a full glass of water with each dose

■ adults and children 12 years and over: take 1 or 2 tablets every

4 hours or 3 tablets every 6 hours not to exceed 12 tablets in

24 hours unless directed by a doctor

■ children under 12 years: do not use unless directed by a doctor

Other information

■ store at 25°C (77°F); excursions permitted between 15°-30°C

(59°-86°F) ■ use by expiration date on package

(59°-86°F) ■ use by expiration date on package

Inactive ingredients corn starch, hypromellose,

polyethylene glycol, propylene glycol

Questions or comments? 1-800-426-9391

Quality Plus 44-157

ASPIRIN

aspirin tablet, film coated

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:50844-157

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	ASPIRIN;44;157
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-157-11	60 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/10/1996	
2	NDC:50844-157-12	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/10/1996	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/10/1996	

Labeler - L.N.K. International, Inc. (038154464)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(50844-157)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(50844-157)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(50844-157)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(50844-157)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(50844-157)

Revised: 1/2025

L.N.K. International, Inc.