

# MINOCYCLINE HYDROCHLORIDE- minocycline hydrochloride tablet Torrent Pharmaceuticals Limited

## MINOCYCLINE HYDROCHLORIDE TABLETS, USP


### Rx only

To reduce the development of drug-resistant bacteria and maintain the effectiveness of minocycline hydrochloride tablets, USP and other antibacterial drugs, minocycline hydrochloride tablets, USP should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.

### DESCRIPTION

Minocycline hydrochloride, USP is a semisynthetic derivative of tetracycline, 4,7-Bis(dimethylamino)-1,4,4a,5,5a,6,11,12a-octahydro-3,10,12,12a-tetrahydroxy-1,11-dioxo-2-naphthacencarboxamide monohydrochloride.

Its structural formula is:



Minocycline hydrochloride tablets, USP for oral administration, contains minocycline hydrochloride, USP equivalent to 50 mg, 75 mg or 100 mg of minocycline. In addition, each film-coated tablet contains the following inactive ingredients: colloidal silicon dioxide, crospovidone, hypromellose, iron oxide yellow, lactose monohydrate, magnesium stearate, polyethylene glycol 6000, silicified microcrystalline cellulose, stearic acid, and titanium dioxide.

### CLINICAL PHARMACOLOGY

Following a single dose of two 100 mg tablet of minocycline hydrochloride administered to 28 normal fasting adult volunteers, maximum serum concentrations were attained in 1 to 3 hours (average 1.71 hours) and ranged from 491.71 to 1,292.70 ng/mL (average 758.29 ng/mL). The serum half-life in the normal volunteers ranged from 11.38 to 24.31 hours (average 17.03 hours).

When minocycline hydrochloride tablets were given concomitantly with a meal, which included dairy products, the extent of absorption of minocycline hydrochloride tablets was slightly decreased (6%). The peak plasma concentrations were slightly decreased

(12%) and delayed by 1.09 hours when administered with food, compared to dosing under fasting conditions. Minocycline hydrochloride tablets may be administered with or without food.

In previous studies with other minocycline dosage forms, the minocycline serum half-life ranged from 11 to 16 hours in 7 patients with hepatic dysfunction, and from 18 to 69 hours in 5 patients with renal dysfunction. The urinary and fecal recovery of minocycline when administered to 12 normal volunteers was one-half to one-third that of other tetracyclines.

## **Microbiology**

### Mechanism of Action

The tetracyclines are primarily bacteriostatic and are thought to exert their antimicrobial effect by the inhibition of protein synthesis. The tetracyclines, including minocycline, have a similar antimicrobial spectrum of activity against a wide range of gram-positive and gram-negative organisms. Cross-resistance of these organisms to tetracycline is common.

### Antimicrobial Activity

Minocycline has been shown to be active against most strains of the following microorganisms, both in vitro and in clinical infections as described in the **INDICATIONS AND USAGE** section:

#### **Gram-positive Bacteria**

*Bacillus anthracis*  
*Listeria monocytogenes*  
*Staphylococcus aureus*  
*Streptococcus pneumoniae*

#### **Gram-negative Bacteria**

*Bartonella bacilliformis*  
*Brucella* species  
*Klebsiella granulomatis*  
*Campylobacter fetus*  
*Francisella tularensis*  
*Haemophilus ducreyi*  
*Vibrio cholerae*  
*Yersinia pestis*  
*Acinetobacter* species  
*Enterobacter aerogenes*  
*Escherichia coli*  
*Haemophilus influenzae*  
*Klebsiella* species  
*Neisseria gonorrhoeae*<sup>1</sup>  
*Neisseria meningitidis*<sup>1</sup>  
*Shigella* species

#### **Other Microorganisms**

*Actinomyces* species

*Borrelia recurrentis*  
*Chlamydophila psittaci*  
*Chlamydia trachomatis*  
*Clostridium* species  
*Entamoeba* species  
*Fusobacterium nucleatum subspecies fusiforme*  
*Mycobacterium marinum*  
*Mycoplasma pneumoniae*  
*Propionibacterium acnes*  
Rickettsiae  
*Treponema pallidum subspecies pallidum*  
*Treponema pallidum subspecies pertenue*  
*Ureaplasma urealyticum*

## Susceptibility Test Methods

When available, the clinical microbiology laboratory should provide cumulative report of *in vitro* susceptibility test results for antimicrobial drugs used in local hospitals and practice areas to the physician as periodic reports that describe the susceptibility profile of nosocomial and community-acquired pathogens. These reports should aid the physician in selecting an antibacterial drug for treatment.

### Dilution techniques:

Quantitative methods are used to determine antimicrobial minimal inhibitory concentrations (MICs). These MICs provide estimates of the susceptibility of bacteria to antimicrobial compounds. The MICs should be determined using a standardized test method (broth or agar) or equivalent with standardized inoculum concentrations and standardized concentrations of tetracycline (class) or minocycline powder<sup>1,2</sup>. The MIC values should be interpreted according to the criteria provided in Table 1.

### Diffusion techniques:

Quantitative methods that require measurement of zone diameters also provide reproducible estimates of the susceptibility of bacteria to antimicrobial compounds. The zone size provides an estimate of the susceptibility of bacteria to antimicrobial compounds. The zone size should be determined using a standardized method.<sup>2,3</sup> This procedure uses paper disks impregnated with 30 µg tetracycline (class disk) or 30 µg minocycline to test the susceptibility of microorganisms to minocycline. The disk diffusion interpretive criteria are provided in Table 1.

**Table 1: Susceptibility Test Interpretive Criteria for Minocycline and Tetracycline**

Species	Minimal Inhibitory Concentration (mcg/mL)			Zone Diameter (mm)			Agar Dilution (mcg/mL)		
	S	I	R	S	I	R	S	I	R
<i>Enterobacteriaceae</i> <sup>a</sup>									
Minocycline	≤ 4	8	≥16	≥16	13 to 15	≤ 12			
Tetracycline	≤ 4	8	≥16	≥15	12 to 14	≤ 11			
<i>Acinetobacter</i> <sup>a</sup>									
Minocycline	≤ 4	8	≥16	≥16	13 to 15	≤ 12			

Tetracycline	≤ 4	8	≥16	≥15	12 to 14	≤ 11			
<i>Haemophilus influenzae</i>									
Tetracycline	≤ 2	4	≥8	≥29	26 to 28	≤ 25			
<i>Streptococcus pneumoniae</i>									
Tetracycline	≤ 1	2	≥4	≥28	25 to 27	≤ 24			
<i>Staphylococcus aureus</i> <sup>a</sup>									
Minocycline	≤ 4	8	≥16	≥19	15 to 18	≤ 14			
Tetracycline	≤ 4	8	≥16	≥19	15 to 18	≤ 14			
<i>Vibrio cholerae</i> <sup>a</sup>									
Minocycline	≤ 4	8	≥16	≥16	13 to 15	≤ 12			
Tetracycline	≤ 4	8	≥16	≥19	15 to 18	≤ 14			
<i>Neisseria meningitidis</i> <sup>b</sup>									
Minocycline	--	--	--	≥26	--	--	≤2	--	--
<i>Bacillus anthracis</i> <sup>b</sup>									
Tetracycline	≤ 1	--	--						
<i>Francisella tularensis</i> <sup>b</sup>									
Tetracycline	≤ 4	--	--						
<i>Yersinia pestis</i>									
Tetracycline	≤ 4	8	≥16						

<sup>a</sup> Organisms that are susceptible to tetracycline are also considered susceptible to minocycline. However, some organisms that are intermediate or resistant to tetracycline may be susceptible to minocycline.

<sup>b</sup> The current absence of resistance isolates precludes defining any result other than “susceptible”. If isolates yielding MIC results other than susceptible, they should be submitted to a reference laboratory for further testing.

A report of *Susceptible* (S) indicates that the antimicrobial drug is likely to inhibit growth of the microorganism if the antimicrobial drug reaches the concentration usually achievable at the site of infection. A report of *Intermediate* (I) indicates that the result should be considered equivocal, and, if the microorganism is not fully susceptible to alternative, clinically feasible drugs, the test should be repeated. This category implies possible clinical applicability in body sites where the drug is physiologically concentrated or in situations where high dosage of drug can be used. This category also provides a buffer zone which prevents small uncontrolled technical factors from causing major discrepancies in interpretation. A report of *Resistant* (R) indicates that the antimicrobial drug is not likely to inhibit growth of the microorganism, if the antimicrobial drug-reaches the concentrations usually achievable at the infection site; other therapy should be selected.

### Quality Control

Standardized susceptibility test procedures require the use of laboratory controls to monitor and ensure the accuracy and precision of supplies and reagents used in the assay, and the techniques of the individuals performing the test.<sup>1,2,3</sup> Standard tetracycline (class compound) or minocycline powder should provide the following range of MIC values noted in Table 2. For the disc diffusion technique, using the 30 mcg tetracycline or 30 mcg minocycline disk the criteria in Table 2 should be achieved.

**Table 2: Acceptable Quality Control Ranges for Minocycline and Tetracycline**

<b>Species</b>	<b>Minimal Inhibitory Concentration (mcg/mL)</b>	<b>Zone Diameter (mm)</b>	<b>Agar Dilution (mcg/mL)</b>
<i>Enterococcus faecalis</i> ATCC 29212 Minocycline Tetracycline	1 to 4 8 to 32	-- --	-- --
<i>Escherichia coli</i> ATCC 25922 Minocycline Tetracycline	0.25 to 1 0.5 to 2	19 to 25 18 to 25	-- --
<i>Haemophilus influenzae</i> ATCC 49247 Tetracycline	4 to 32	14 to 22	--
<i>Neisseria gonorrhoeae</i> ATCC 49226 Tetracycline	--	30 to 42	0.25 to 1
<i>Staphylococcus aureus</i> ATCC 25923 Minocycline Tetracycline		25 to 30 24 to 30	-- --
<i>Staphylococcus aureus</i> ATCC 29213 Minocycline Tetracycline	0.06 to 0.5 0.12 to 1		-- --
<i>Streptococcus pneumoniae</i> ATCC 49619 Tetracycline	0.06 to 0.5	27 to 31	--

## **INDICATIONS AND USAGE**

Minocycline hydrochloride tablets, USP are indicated in the treatment of the following infections due to susceptible strains of the designated microorganisms:

Rocky Mountain spotted fever, typhus fever and the typhus group, Q fever, rickettsialpox and tick fevers caused by rickettsiae.

Respiratory tract infections caused by *Mycoplasma pneumoniae*.

Lymphogranuloma venereum caused by *Chlamydia trachomatis*.

Psittacosis (Ornithosis) due to *Chlamydophila psittaci*.

Trachoma caused by *Chlamydia trachomatis*, although the infectious agent is not always eliminated, as judged by immunofluorescence.

Inclusion conjunctivitis caused by *Chlamydia trachomatis*.

Nongonococcal urethritis, endocervical, or rectal infections in adults caused by *Ureaplasma urealyticum* or *Chlamydia trachomatis*.

Relapsing fever due to *Borrelia recurrentis*.

Relapsing fever due to *Borrelia recurrentis*.

Plague due to *Yersinia pestis*.

Tularemia due to *Francisella tularensis*.

Cholera caused by *Vibrio cholerae*.

Campylobacter fetus infections caused by *Campylobacter fetus*.

Brucellosis due to *Brucella* species (in conjunction with streptomycin).

Bartonellosis due to *Bartonella bacilliformis*.

Granuloma inguinale caused by *Klebsiella granulomatis*.

Minocycline is indicated for the treatment of infections caused by the following gram-negative microorganisms when bacteriologic testing indicates appropriate susceptibility to the drug:

*Escherichia coli*.

*Enterobacter aerogenes*.

*Shigella* species.

*Acinetobacter* species.

Respiratory tract infections caused by *Haemophilus influenzae*.

Respiratory tract and urinary tract infections caused by *Klebsiella* species.

Minocycline hydrochloride tablets, USP are indicated for the treatment of infections caused by the following gram-positive microorganisms when bacteriologic testing indicates appropriate susceptibility to the drug:

Upper respiratory tract infections caused by *Streptococcus pneumoniae*.

Skin and skin structure infections caused by *Staphylococcus aureus* (Note: Minocycline is not the drug of choice in the treatment of any type of staphylococcal infection).

When penicillin is contraindicated, minocycline is an alternative drug in the treatment of the following infections:

Uncomplicated urethritis in men due to *Neisseria gonorrhoeae* and for the treatment of other gonococcal infections.

Infections in women caused by *Neisseria gonorrhoeae*.

Syphilis caused by *Treponema pallidum* subspecies *pallidum*.

Yaws caused by *Treponema pallidum* subspecies *pertenue*.

Listeriosis due to *Listeria monocytogenes*.

Anthrax due to *Bacillus anthracis*.

Vincent's infection caused by *Fusobacterium fusiforme*.

Actinomycosis caused by *Actinomyces israelii*.

Infections caused by *Clostridium* species.

In *acute intestinal amebiasis*, minocycline may be a useful adjunct to amebicides.

In severe *acne*, minocycline may be useful adjunctive therapy.

Oral minocycline is indicated in the treatment of asymptomatic carriers of *Neisseria meningitidis* to eliminate meningococci from the nasopharynx. In order to preserve the usefulness of minocycline in the treatment of asymptomatic meningococcal carriers,

diagnostic laboratory procedures, including serotyping and susceptibility testing, should be performed to establish the carrier state and the correct treatment. It is recommended that the prophylactic use of minocycline be reserved for situations in which the risk of meningococcal meningitis is high.

*Oral minocycline is not indicated for the treatment of meningococcal infection.*

Although no controlled clinical efficacy studies have been conducted, limited clinical data show that oral minocycline hydrochloride has been used successfully in the treatment of infections caused by *Mycobacterium marinum*.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of minocycline hydrochloride tablets, USP and other antibacterial drugs, minocycline hydrochloride tablets, USP should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

## **CONTRAINDICATIONS**

This drug is contraindicated in persons who have shown hypersensitivity to any of the tetracyclines or to any of the components of the product formulation.

## **WARNINGS**

### **Tooth Development**

Minocycline hydrochloride, like other tetracycline-class antibiotics, can cause fetal harm when administered to a pregnant woman. If any tetracycline is used during pregnancy or if the patient becomes pregnant while taking these drugs, the patient should be apprised of the potential hazard to the fetus. The use of drugs of the tetracycline class during tooth development (last half of pregnancy, infancy, and childhood to the age of 8 years) may cause permanent discoloration of the teeth (yellow-gray-brown).

This adverse reaction is more common during long-term use of the drug but has been observed following repeated short-term courses. Enamel hypoplasia has also been reported. Tetracycline drugs, therefore, should not be used during tooth development unless other drugs are not likely to be effective or are contraindicated.

### **Skeletal Development**

All tetracyclines form a stable calcium complex in any bone-forming tissue. A decrease in the fibula growth rate has been observed in premature human infants given oral tetracycline in doses of 25 mg/kg every six hours. This reaction was shown to be reversible when the drug was discontinued.

### **Use in Pregnancy**

Results of animal studies indicate that tetracyclines cross the placenta, are found in fetal tissues, and can have toxic effects on the developing fetus (often related to retardation of skeletal development). Evidence of embryotoxicity has been noted in animals treated

early in pregnancy.

## **Dermatologic Reaction**

Drug Rash with Eosinophilia and Systemic Symptoms (DRESS) including fatal cases have been reported with minocycline use. If this syndrome is recognized, the drug should be discontinued immediately.

## **Antianabolic Action**

The antianabolic action of the tetracyclines may cause an increase in BUN. While this is not a problem in those with normal renal function, in patients with significantly impaired function, higher serum levels of tetracycline may lead to azotemia, hyperphosphatemia, and acidosis. Under such conditions, monitoring of creatinine and BUN is recommended, and the total daily dosage should not exceed 200 mg in 24 hours (see **DOSAGE AND ADMINISTRATION**). If renal impairment exists, even usual oral or parenteral doses may lead to systemic accumulation of the drug and possible liver toxicity.

## **Photosensitivity**

Photosensitivity manifested by an exaggerated sunburn reaction has been observed in some individuals taking tetracyclines. This has been reported with minocycline.

## **Central Nervous System**

Central nervous system side effects including light-headedness, dizziness, or vertigo have been reported with minocycline therapy. Patients who experience these symptoms should be cautioned about driving vehicles or using hazardous machinery while on minocycline therapy. These symptoms may disappear during therapy and usually disappear rapidly when the drug is discontinued.

## ***Clostridium difficile* Associated Diarrhea**

*Clostridium difficile* associated diarrhea (CDAD) has been reported with use of nearly all antibacterial agents, including minocycline hydrochloride, and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

*C. difficile* produces toxins A and B which contribute to the development of CDAD. Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibiotic use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing antibiotic use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibiotic treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated.

## **Intracranial Hypertension**

Intracranial hypertension (IH, pseudotumor cerebri) has been associated with the use of tetracyclines including minocycline hydrochloride. Clinical manifestations of IH include



headache, blurred vision, diplopia, and vision loss; papilledema can be found on fundoscopy. Women of childbearing age who are overweight or have a history of IH are at greater risk for developing tetracycline associated IH. Concomitant use of isotretinoin and minocycline hydrochloride should be avoided because isotretinoin is also known to cause pseudotumor cerebri.

Although IH typically resolves after discontinuation of treatment, the possibility for permanent visual loss exists. If visual disturbance occurs during treatment, prompt ophthalmologic evaluation is warranted. Since intracranial pressure can remain elevated for weeks after drug cessation patients should be monitored until they stabilize.

## **PRECAUTIONS**

### **General**

As with other antibiotic preparations, use of this drug may result in overgrowth of non-susceptible organisms, including fungi. If superinfection occurs, the antibiotic should be discontinued and appropriate therapy instituted.

Hepatotoxicity has been reported with minocycline; therefore, minocycline should be used with caution in patients with hepatic dysfunction and in conjunction with other hepatotoxic drugs.

Incision and drainage or other surgical procedures should be performed in conjunction with antibiotic therapy when indicated.

Prescribing minocycline hydrochloride tablets in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

### **Information For Patients**

Diarrhea is a common problem caused by antibiotics which usually ends when the antibiotic is discontinued. Sometimes after starting treatment with antibiotics, patients can develop watery and bloody stools (with or without stomach cramps and fever) even as late as two or more months after having taken the last dose of the antibiotic. If this occurs, patients should contact their physician as soon as possible.

Photosensitivity manifested by an exaggerated sunburn reaction has been observed in some individuals taking tetracyclines. Patients apt to be exposed to direct sunlight or ultraviolet light should be advised that this reaction can occur with tetracycline drugs, and treatment should be discontinued at the first evidence of skin erythema. This reaction has been reported with use of minocycline.

Patients who experience central nervous system symptoms should be cautioned about driving vehicles or using hazardous machinery while on minocycline therapy (See **WARNINGS**).

Concurrent use of tetracycline with oral contraceptives may render oral contraceptives less effective (See **Drug Interactions**).

Patients should be counseled that antibacterial drugs including minocycline hydrochloride tablets should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When minocycline hydrochloride tablets are

prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by minocycline hydrochloride tablets or other antibacterial drugs in the future.

Unused supplies of tetracycline antibiotics should be discarded by the expiration date.

### **Laboratory Tests**

In venereal disease when coexistent syphilis is suspected, a dark-field examination should be done before treatment is started and the blood serology repeated monthly for at least four months.

Periodic laboratory evaluations of organ systems, including hematopoietic, renal, and hepatic, should be performed.

### **Drug Interactions**

Because tetracyclines have been shown to depress plasma prothrombin activity, patients who are on anticoagulant therapy may require downward adjustment of their anticoagulant dosage.

Since bacteriostatic drugs may interfere with the bactericidal action of penicillin, it is advisable to avoid giving tetracycline-class drugs in conjunction with penicillin.

Absorption of tetracyclines is impaired by antacids containing aluminum, calcium, or magnesium, and iron-containing preparations.

The concurrent use of tetracycline and methoxyflurane has been reported to result in fatal renal toxicity.

Concurrent use of tetracyclines with oral contraceptives may render oral contraceptives less effective.

Administration of isotretinoin should be avoided shortly before, during, and shortly after minocycline therapy. Each drug alone has been associated with pseudotumor cerebri (See **PRECAUTIONS**).

Increased risk of ergotism when ergot alkaloids or their derivatives are given with tetracyclines.

### **Drug/Laboratory Test Interactions**

False elevations of urinary catecholamine levels may occur due to interference with the fluorescence test.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Dietary administration of minocycline in long term tumorigenicity studies in rats resulted in evidence of thyroid tumor production. Minocycline has also been found to produce thyroid hyperplasia in rats and dogs. In addition, there has been evidence of oncogenic activity in rats in studies with a related antibiotic, oxytetracycline (i.e., adrenal and pituitary tumors). Likewise, although mutagenicity studies of minocycline have not been conducted, positive results in *in vitro* mammalian cell assays (i.e., mouse lymphoma and

Chinese hamster lung cells) have been reported for related antibiotics (tetracycline hydrochloride and oxytetracycline). Segment I (fertility and general reproduction) studies have provided evidence that minocycline impairs fertility in male rats.

## **Pregnancy**

### **Risk Summary**

All pregnancies have a background risk of birth defects, loss, or other adverse outcome regardless of drug exposure. There are no adequate and well-controlled studies on the use of minocycline in pregnant women. Minocycline, like other tetracycline-class antibiotics, crosses the placenta and may cause fetal harm when administered to a pregnant woman. Rare spontaneous reports of congenital anomalies including limb reduction have been reported in post-marketing experience. Only limited information is available regarding these reports; therefore, no conclusion on causal association can be established. If minocycline is used during pregnancy or if the patient becomes pregnant while taking this drug, the patient should be apprised of the potential hazard to the fetus.

*Nonteratogenic Effects:* (See **WARNINGS**).

## **Labor and Delivery**

The effect of tetracyclines on labor and delivery is unknown.

## **Nursing Mothers**

Tetracyclines are excreted in human milk. Because of the potential for serious adverse reactions in nursing infants from the tetracyclines, a decision should be made whether to discontinue nursing or discontinue the drug, taking into account the importance of the drug to the mother (See **WARNINGS**).

## **Pediatric Use**

Minocycline is not recommended for the use in children below 8 years of age unless the expected benefits of therapy outweigh the risks (See **WARNINGS**).

## **Geriatric Use**

Clinical studies of oral minocycline did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy (See **WARNINGS, DOSAGE AND ADMINISTRATION**).

## **ADVERSE REACTIONS**

Due to oral minocycline's virtually complete absorption, side effects to the lower bowel, particularly diarrhea, have been infrequent. The following adverse reactions have been observed in patients receiving tetracyclines.

*Body as a whole:* Fever, and discoloration of secretions.

*Gastrointestinal:* Anorexia, nausea, vomiting, diarrhea, dyspepsia, stomatitis, glossitis, dysphagia, enamel hypoplasia, enterocolitis, pseudomembranous colitis, pancreatitis, inflammatory lesions (with monilial overgrowth) in the oral and anogenital regions. Instances of esophagitis and esophageal ulcerations have been reported in patients taking the tetracycline-class antibiotics in capsule and tablet form. Most of these patients took the medication immediately before going to bed (See **DOSAGE AND ADMINISTRATION**).

*Genitourinary:* Vulvovaginitis.

*Hepatic toxicity:* Hyperbilirubinemia, hepatic cholestasis, increases in liver enzymes, fatal hepatic failure, and jaundice. Hepatitis, including autoimmune hepatitis, and liver failure have been reported (See **PRECAUTIONS**).

*Skin:* Alopecia, erythema nodosum, hyperpigmentation of nails, pruritus, toxic epidermal necrolysis, vasculitis, maculopapular rash and erythematous rash. Exfoliative dermatitis has been reported. Fixed drug eruptions have been reported. Lesions occurring on the glans penis have caused balanitis. Erythema multiforme and Stevens-Johnson syndrome have been reported. Photosensitivity is discussed above (See **WARNINGS**). Pigmentation of the skin and mucous membranes has been reported.

*Respiratory:* Cough, dyspnea, bronchospasm, exacerbation of asthma, and pneumonitis.

*Renal toxicity:* Interstitial nephritis. Elevations in BUN have been reported and are apparently dose related (See **WARNINGS**). Reversible acute renal failure has been reported.

*Musculoskeletal:* Arthralgia, arthritis, bone discoloration, myalgia, joint stiffness, and joint swelling.

*Hypersensitivity reactions:* Urticaria, angioneurotic edema, polyarthralgia, anaphylaxis/anaphylactoid reaction (including shock and fatalities), anaphylactoid purpura, myocarditis, pericarditis, exacerbation of systemic lupus erythematosus and pulmonary infiltrates with eosinophilia have been reported. A transient lupus-like syndrome and serum sickness-like reactions also have been reported.

*Blood:* Agranulocytosis, hemolytic anemia, thrombocytopenia, leukopenia, neutropenia, pancytopenia, and eosinophilia have been reported.

*Central Nervous System:* Convulsions, dizziness, hypesthesia, paresthesia, sedation, and vertigo. Bulging fontanel in infants and benign intracranial hypertension (pseudotumor cerebri) in adults have been reported (See **PRECAUTIONS - General**). Headache has also been reported.

*Other:* Thyroid cancer has been reported in the post-marketing setting in association with minocycline products. When minocycline therapy is given over prolonged periods, monitoring for signs of thyroid cancer should be considered. When given over prolonged periods, tetracyclines have been reported to produce brown-black microscopic discoloration of the thyroid gland. Cases of abnormal thyroid function have been reported.

Tooth discoloration in children less than 8 years of age (see **WARNINGS**) and also, in adults has been reported.

Oral cavity discoloration (including tongue, lip, and gum) has been reported.

Tinnitus and decreased hearing have been reported in patients on minocycline hydrochloride.

The following syndromes have been reported. In some cases involving these syndromes, death has been reported. As with other serious adverse reactions, if any of these syndromes are recognized, the drug should be discontinued immediately:

Hypersensitivity syndrome consisting of cutaneous reaction (such as rash or exfoliative dermatitis), eosinophilia, and one or more of the following: hepatitis, pneumonitis, nephritis, myocarditis, and pericarditis. Fever and lymphadenopathy may be present.

Lupus-like syndrome consisting of positive antinuclear antibody; arthralgia, arthritis, joint stiffness, or joint swelling; and one or more of the following: fever, myalgia, hepatitis, rash, and vasculitis.

Serum sickness-like syndrome consisting of fever; urticaria or rash; and arthralgia, arthritis, joint stiffness, or joint swelling and lymphadenopathy. Eosinophilia may be present.

**To report SUSPECTED ADVERSE REACTIONS, contact Torrent Pharmaceuticals Inc. at 1-800-912-9561 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

## **OVERDOSAGE**

The adverse events more commonly seen in overdose are dizziness, nausea, and vomiting.

No specific antidote for minocycline is known.

In case of overdosage, discontinue medication, treat symptomatically, and institute supportive measures. Minocycline is not removed in significant quantities by hemodialysis or peritoneal dialysis.

## **DOSAGE AND ADMINISTRATION**

**THE USUAL DOSAGE AND FREQUENCY OF ADMINISTRATION OF MINOCYCLINE DIFFERS FROM THAT OF THE OTHER TETRACYCLINES. EXCEEDING THE RECOMMENDED DOSAGE MAY RESULT IN AN INCREASED INCIDENCE OF SIDE EFFECTS.**

Minocycline hydrochloride tablets may be taken with or without food (See **CLINICAL PHARMACOLOGY**).

Ingestion of adequate amounts of fluids along with capsule and tablet forms of drugs in the tetracycline-class is recommended to reduce the risk of esophageal irritation and ulceration. The tablets should be swallowed whole.

### **For Pediatric Patients Above 8 Years Of Age**

Usual pediatric dose: 4 mg/kg initially followed by 2 mg/kg every 12 hours, not to exceed the usual adult dose.

## Adults

The usual dosage of minocycline hydrochloride tablets is 200 mg initially followed by 100 mg every 12 hours. Alternatively, if more frequent doses are preferred, two or four 50 mg tablets may be given initially followed by one 50 mg tablet 4 times daily.

Uncomplicated gonococcal infections other than urethritis and anorectal infections in men: 200 mg initially, followed by 100 mg every 12 hours for a minimum of 4 days, with post-therapy cultures within 2 to 3 days.

In the treatment of uncomplicated gonococcal urethritis in men, 100 mg every 12 hours for 5 days is recommended.

For the treatment of syphilis, the usual dosage of minocycline hydrochloride should be administered over a period of 10 to 15 days. Close follow-up, including laboratory tests, is recommended.

In the treatment of meningococcal carrier state, the recommended dosage is 100 mg every 12 hours for 5 days.

*Mycobacterium marinum* infections: Although optimal doses have not been established, 100 mg every 12 hours for 6 to 8 weeks have been used successfully in a limited number of cases.

Uncomplicated urethral, endocervical, or rectal infection in adults caused by *Chlamydia trachomatis* or *Ureaplasma urealyticum*: 100 mg orally, every 12 hours for at least 7 days.

Ingestion of adequate amounts of fluids along with capsule and tablet forms of drugs in the tetracycline-class is recommended to reduce the risk of esophageal irritation and ulceration.

The pharmacokinetics of minocycline in patients with renal impairment ( $CL_{CR} < 80$  mL/min) have not been fully characterized. Current data are insufficient to determine if a dosage adjustment is warranted. The total daily dosage should not exceed 200 mg in 24 hours. However, due to the anti-anabolic effect of tetracyclines, BUN and creatinine should be monitored (See **WARNINGS**).

## HOW SUPPLIED

Minocycline hydrochloride tablets, USP equivalent to 50 mg minocycline are yellow colored, oval-shaped, film-coated tablets, debossed with “**R189**” on one side and plain on the other side. They are supplied as follows:

NDC 13668-485-01

Bottles of 100

Minocycline hydrochloride tablets, USP equivalent to 75 mg minocycline are yellow colored, oval-shaped, film-coated tablets, debossed with “**R190**” on one side and plain on the other side. They are supplied as follows:

NDC 13668-486-01

Bottles of 100

Minocycline hydrochloride tablets, USP equivalent to 100 mg minocycline are yellow colored, oval-shaped, film-coated tablets, debossed with “**R191**” on one side and plain on the other side. They are supplied as follows:

NDC 13668-487-50

Bottles of 50

**Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].**

**Protect from light, moisture and excessive heat.**

**Dispense in a tight, light-resistant container with child-resistant closure.**

## **ANIMAL PHARMACOLOGY AND TOXICOLOGY**

Minocycline hydrochloride has been observed to cause a dark discoloration of the thyroid in experimental animals (rats, minipigs, dogs, and monkeys). In the rat, chronic treatment with minocycline hydrochloride has resulted in goiter accompanied by elevated radioactive iodine uptake and evidence of thyroid tumor production. Minocycline hydrochloride has also been found to produce thyroid hyperplasia in rats and dogs.

## **REFERENCES**

1. Clinical and Laboratory Standards Institute (CLSI). Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria that Grow Aerobically; Approved Standard-Tenth Edition; CLSI Document M07-A10, Vol. 32, No. 2, January, 2015. Clinical and Laboratory Standards, 940 West Valley Rd., Suite 2500, Wayne, PA 19087-1898.
2. Clinical and Laboratory Standards Institute (CLSI). Performance Standards for Antimicrobial Disk Susceptibility Tests; Approved Standard-Twelfth Edition; CLSI Document M02-A12, Vol. 32, No. 1, January, 2015. Clinical and Laboratory Standards, 940 West Valley Rd., Suite 2500, Wayne, PA 19087-1898.
3. Clinical and Laboratory Standards Institute (CLSI). Performance Standards for Antimicrobial Susceptibility Testing; Twenty-sixth Informational Supplement. Document M100-S26, Vol. 32, No. 3, January, 2016. Clinical and Laboratory Standards, 940 West Valley Rd., Suite 2500, Wayne, PA 19087-1898.

## **PATIENT INFORMATION**

### **PATIENT INFORMATION**

### **MINOCYCLINE HYDROCHLORIDE**

#### **(Mi-no-syk-lin hi-droh-clor-ride) TABLETS, USP**

**50 mg, 75 mg and 100 mg**

#### **Rx only**

Read the Patient Information that comes with minocycline hydrochloride tablets, USP before you or a family member starts taking it and each time you get a refill. There may be new information. This leaflet does not take the place of talking to your doctor about your medical condition or treatment.

#### **What are minocycline hydrochloride tablets, USP?**

Minocycline hydrochloride tablets, USP are a tetracycline-class antibiotic medicine.

Minocycline hydrochloride tablets, USP are used to treat certain infections caused by bacteria. These include infections of the skin, respiratory tract, urinary tract, some sexually transmitted diseases, and others. Minocycline hydrochloride tablets, USP may be used along with other treatments for severe acne.

Sometimes, other germs, called viruses cause infections. The common cold is a virus. Minocycline hydrochloride tablets, USP, like other antibiotics, does not treat viruses.

### **Who should not use minocycline hydrochloride tablets, USP?**

**Do not take minocycline hydrochloride tablets, USP if you are allergic to minocycline or other tetracycline antibiotics.**

Ask your doctor or pharmacist for a list of these medications if you are not sure. See the end of this leaflet for a complete list of ingredients in minocycline hydrochloride tablets, USP. **Minocycline hydrochloride tablets, USP are not recommended for pregnant women or children up to 8 years old because:**

1. **Minocycline hydrochloride tablets, USP may harm an unborn baby**
2. **Minocycline hydrochloride tablets, USP may permanently turn a baby's or child's teeth yellow-gray-brown during tooth development.** Tooth development happens in the last half of pregnancy and birth to age 8 years.

### **What should I tell my doctor before starting minocycline hydrochloride tablets, USP?**

Tell your doctor about all of your medical conditions, including if you:

- **have liver or kidney problems**
- **are pregnant or planning to become pregnant.** Minocycline hydrochloride tablets, USP may harm your unborn baby. **Stop taking minocycline hydrochloride tablets, USP and call your doctor if you become pregnant while taking it.**
- **are breast feeding.** Minocycline passes into your milk and may harm your baby. You should decide whether to use minocycline hydrochloride tablets, USP or breastfeed, but not both.

Tell your doctor about all the medicines you are taking including prescription and non-prescription medications, vitamins, and herbal supplements. Minocycline hydrochloride tablets, USP and other medicines may interact. Especially tell your doctor if you take:

- **birth control pills.** Minocycline hydrochloride tablets, USP may make your birth control pills less effective.
- **a blood thinner medicine.** The dose of your blood thinner may have to be lowered.
- **a penicillin antibiotic medicine.** Minocycline hydrochloride tablets, USP and penicillins should not be used together.
- **Migraine medicines called ergot alkaloids.**
- **An acne medicine called isotretinoin (Accutane, Amnesteem, Claravis, Sotret).**
- **Antacids that contain aluminum, calcium, or magnesium, or iron-containing products.**

Know the medicines you take, keep a list of them to show your doctor and pharmacist each time you get a new medicine.

### **How should I take minocycline hydrochloride tablets, USP?**



- **Take minocycline hydrochloride tablets, USP exactly as your doctor tells you to take them.** Skipping doses or not taking all your minocycline hydrochloride tablets, USP may:
  - o Decrease the effectiveness of the treatment.
  - o Increase the chance that bacteria will develop resistance to minocycline hydrochloride tablets, USP.
- **Take minocycline hydrochloride tablets, USP with a full glass of liquid.** Taking minocycline hydrochloride tablets, USP with enough liquid may lower your chance of getting irritation or ulcers in your esophagus. Your esophagus is the tube that connects your mouth to your stomach.
- **Minocycline hydrochloride tablets, USP** may be taken with or without food. If you forget to take minocycline hydrochloride tablets, USP, take it as soon as you remember.
- If you take too much minocycline hydrochloride tablets, USP, call your doctor or poison control center right away.

### **What are the possible side effects of minocycline hydrochloride tablets, USP?**

**Minocycline hydrochloride tablets, USP may cause serious side effects. Stop minocycline hydrochloride tablets, USP and call your doctor if you have:**

- watery diarrhea
- bloody stools
- stomach cramps
- unusual headaches
- blurred vision
- fever
- rash
- joint pain
- feeling very tired
- swollen lymph nodes

### **Minocycline hydrochloride tablets, USP may also cause:**

- **central nervous system effects.** Symptoms include light-headedness, dizziness, and a spinning feeling (vertigo). You should not drive or operate machines if you have these symptoms.
- **sun sensitivity (photosensitivity).** You may get a worse sunburn with minocycline hydrochloride tablets, USP. Avoid sun exposure and the use of sunlamps or tanning beds. Protect your skin while out in the sunlight. Stop minocycline hydrochloride tablets, USP and call your doctor if your skin turns red.

These are not all the side effects with minocycline hydrochloride tablets, USP. Ask your doctor or pharmacist for more information.

**CALL YOUR DOCTOR FOR MEDICAL ADVICE ABOUT SIDE EFFECTS. YOU MAY REPORT SIDE EFFECTS TO THE FDA AT 1-800-FDA-1088.**

### **How should I store minocycline hydrochloride tablets, USP?**

- Store minocycline hydrochloride tablets, USP at room temperature and away from excess heat and moisture.
- Throw away any minocycline hydrochloride tablets, USP that is outdated or no longer

needed.

- **Keep minocycline hydrochloride tablets, USP and all medicines out of the reach of children.**

### **General advice about minocycline hydrochloride tablets, USP**

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use minocycline hydrochloride tablets, USP for a condition for which it was not prescribed. Do not give minocycline hydrochloride tablets, USP to other people, even if they have the same symptoms you have. It may harm them.

This Patient Information leaflet summarizes the most important information about minocycline hydrochloride tablets, USP.

If you would like more information, talk with your doctor. Your doctor or pharmacist can give you information about minocycline hydrochloride tablets, USP that is written for health care professionals. For more information, you can also call Torrent Pharmaceuticals Inc. at 1-800-912-9561.

### **What are the ingredients in minocycline hydrochloride tablets, USP?**

**Active ingredient:** minocycline hydrochloride, USP, 50 mg, 75 mg and 100 mg

**Inactive ingredients:** colloidal silicon dioxide, crospovidone, hypromellose, iron oxide yellow, lactose monohydrate, magnesium stearate, polyethylene glycol 6000, silicified microcrystalline cellulose, stearic acid, and titanium dioxide.

This Patient information has been approved by the U.S. Food and Drug Administration.

Trademarks are the property of their respective owners.



### **Manufactured by:**

TORRENT PHARMACEUTICALS LTD., INDIA

### **Manufactured for:**

TORRENT PHARMA INC., Levittown, PA 19057.

8083734

August 2021

### **PACKAGE LABEL.PRINCIPAL DISPLAY PANEL - 50 mg**

Minocycline Hydrochloride Tablets, USP

50 mg

### **Rx only**

**PHARMACIST:** Dispense the accompanying Patient Information Sheet to each patient.

**Manufactured by:** TORRENT PHARMACEUTICALS LTD.



## PACKAGE LABEL.PRINCIPAL DISPLAY PANEL - 75 mg

Minocycline Hydrochloride Tablets, USP

75 mg

**Rx only**

**PHARMACIST:** Dispense the accompanying Patient Information Sheet to each patient.

**Manufactured by:** TORRENT PHARMACEUTICALS LTD.



## PACKAGE LABEL.PRINCIPAL DISPLAY PANEL - 100 mg

Minocycline Hydrochloride Tablets, USP

100 mg

**Rx only**

**PHARMACIST:** Dispense the accompanying Patient Information Sheet to each patient.

**Manufactured by:** TORRENT PHARMACEUTICALS LTD.

\*Each film-coated tablet contains:  
 Minocycline Hydrochloride, USP equivalent to  
 100 mg minocycline.  
**Keep this and all drugs out of the reach of children.**  
 Store at 20° to 25°C (68° to 77°F); excursions  
 permitted to 15° to 30°C (59° to 86°F) [see USP  
 Controlled Room Temperature].  
 Protect from light, moisture and excessive heat.  
 Dispense in a tight, light-resistant container with  
 child-resistant closure.  
**Usual dosage:** See accompanying prescribing  
 Information.  
 This package not for household use.

Mfg. Lic. No.: G/25/2010  
 Manufactured by: TORRENT PHARMACEUTICALS LTD.  
 Bharuch-392130, INDIA.  
 Manufactured for: TORRENT PHARMA INC.  
 Levittown, PA 19057.

50 Tablets NDC 13668-487-50


**Minocycline  
 Hydrochloride Tablets,  
 USP**

**100 mg\***

**PHARMACIST:** Dispense the  
 accompanying Patient Information Sheet  
 to each patient.



**Rx only**



## MINOCYCLINE HYDROCHLORIDE

minocycline hydrochloride tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:13668-485
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>MINOCYCLINE HYDROCHLORIDE</b> (UNII: 0020414E5U) (MINOCYCLINE - UNII:FYY3R43WGO)	MINOCYCLINE	50 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>CROSPVIDONE (15 MPA.S AT 5%)</b> (UNII: 68401960MK)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>POLYETHYLENE GLYCOL 6000</b> (UNII: 30IQX730WE)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	RI89
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:13668-485-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	04/02/2015	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA065156	04/02/2015	

## MINOCYCLINE HYDROCHLORIDE

minocycline hydrochloride tablet

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:13668-486
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MINOCYCLINE HYDROCHLORIDE (UNII: 0020414E5U) (MINOCYCLINE - UNII:FYY3R43WGO)	MINOCYCLINE	75 mg

### Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
CROSPVIDONE (15 MPA.S AT 5%) (UNII: 68401960MK)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOL 6000 (UNII: 30IQX730WE)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

### Product Characteristics

Color	yellow	Score	no score
Shape	OVAL	Size	14mm
Flavor		Imprint Code	RI90

**Contains****Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:13668-486-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	04/02/2015	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA065156	04/02/2015	

**MINOCYCLINE HYDROCHLORIDE**

minocycline hydrochloride tablet

**Product Information**

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:13668-487
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>MINOCYCLINE HYDROCHLORIDE</b> (UNII: 0020414E5U) (MINOCYCLINE - UNII:FYY3R43WGO)	MINOCYCLINE	100 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>CROSPROVIDONE (15 MPA.S AT 5%)</b> (UNII: 68401960MK)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX43802MRT)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>POLYETHYLENE GLYCOL 6000</b> (UNII: 30IQX730WE)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

**Product Characteristics**

<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	15mm

<b>Flavor</b>		<b>Imprint Code</b>	RI91	
<b>Contains</b>				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:13668-487-50	50 in 1 BOTTLE; Type 0: Not a Combination Product	04/02/2015	
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA065156	04/02/2015		

**Labeler** - Torrent Pharmaceuticals Limited (916488547)

**Registrant** - Torrent Pharma, Inc. (790033935)

<b>Establishment</b>			
Name	Address	ID/FEI	Business Operations
Torrent Pharmaceuticals Limited		864147745	manufacture(13668-485, 13668-485, 13668-487)

<b>Establishment</b>			
Name	Address	ID/FEI	Business Operations
Torrent Pharmaceuticals Limited		916488547	manufacture(13668-485, 13668-486, 13668-487) , label(13668-485, 13668-486, 13668-487) , pack(13668-485, 13668-486, 13668-487)