

**ANTACID MAXIMUM STRENGTH- aluminum hydroxide, magnesium hydroxide,
dimethicone liquid
STRATEGIC SOURCING SERVICES LLC**

SUNMARK ANTACID MAX

Active ingredients (in each 5 mL teaspoonful)

Aluminum hydroxide 400mg (equivalent to dried gel, USP)
Magnesium hydroxide 400 mg
Simethicone 40mg

Purposes

Antacid

Antigas

Uses

relieves

- heartburn
- sour stomach
- acid indigestion
- upset stomach due to these symptoms
- symptoms of gas

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are

presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use.

Keep this and all drugs out of the reach of children.

in case of accidental overdose, seek professional assistance or contact a poison control center (1-800-222-1222) immediately.

Directions

- **shake well before use**

- do not exceed 12 teaspoons (60 mL) in a 24 hour period or use the maximum dosage for more than 2 weeks
- adults and children 12 years and older: take 2 to 4 teaspoons (10-20 mL) between meals, at bedtime, or as directed by a doctor
- children under 12 years: ask a doctor

Other information

- **each 5 mL teaspoonful contains:** magnesium 165 mg, sodium 5 mg
- do not freeze
- store at room temperature tightly closed

Inactive ingredients

benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor (contains alcohol), hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

package label

Sunmark

compare to MYLANTA Maximum Strength active ingredients

NDC 49348-303-39

antacid

Maximum Strength

antacid/antigas

Aluminum Hydroxide 400 mg

Magnesium Hydroxide 400 mg

Simethicone 40 mg

Fast acting

fast soothing relief of

heartburn,

acid indigestion,

sour stomach,

pressure & bloating

ORIGINAL

Alcohol: 0,5%

12 FL OZ (355 mL)

sunmark[®] COMPARE TO MYLANTA[®] MAXIMUM STRENGTH ACTIVE INGREDIENTS* NDC 49348-303-39

antacid

Maximum Strength Antacid/Anti-Gas
 Aluminum Hydroxide 400 mg
 Magnesium Hydroxide 400 mg
 Simethicone 40 mg

Fast acting
 Fast soothing relief of heartburn, acid indigestion, sour stomach, pressure & bloating

ORIGINAL
 Alcohol: 0.5%
 12 FL OZ (355 mL)

273-06122-1
 REV GC-0317

Drug Facts TAMPER EVIDENT: DO NOT USE IF THE BREAKAWAY BAND ON CAP IS BROKEN OR MISSING.

Active ingredients (in each 5 mL teaspoonful)	Purpose
Aluminum hydroxide (equivalent to Dried Gel, USP) 400 mg	Antacid
Magnesium hydroxide 400 mg	Antacid
Simethicone 40 mg	Antigas

Uses relieves • heartburn • sour stomach • acid indigestion • upset stomach due to these symptoms • symptoms of gas

Warnings
 Ask a doctor before use if you have
 • kidney disease • a magnesium restricted diet
 Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.
 Stop use and ask a doctor if symptoms last for more than 2 weeks
 If pregnant or breast-feeding, ask a health professional before use.
 Keep this and all drugs out of the reach of children.
 In case of overdose, seek professional assistance or contact a Poison Control Center (1-800-222-1222) immediately.

Directions • shake well before use
 • do not exceed 12 teaspoonfuls (60 mL) in a 24 hour period or use the maximum dosage for more than 2 weeks
 • adults and children 12 years of age and older: take 2 to 4 teaspoonfuls (10-20 mL) between meals, at bedtime or as directed by a doctor
 • children under 12 years of age: ask a doctor

Other information • each 5 mL teaspoonful contains: magnesium 165 mg, sodium 5 mg
 • do not freeze
 • store at room temperature tightly closed

Inactive ingredients benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor (contains alcohol), hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

*This product is not manufactured or distributed by Johnson & Johnson Merck, distributor of Mylanta[®]

McKesson Another Quality Product
 Distributed By McKesson
 One Post Street, San Francisco, CA 94104
 Money Back Guarantee
 Please visit us at www.sunmarkbrand.com

Questions or Comments?
 1-866-534-4631

973-06122-2
 REV GC-0317

0 10939 57844 0

ANTACID MAXIMUM STRENGTH

aluminum hydroxide, magnesium hydroxide, dimethicone liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49348-303
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	400 mg in 5 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	400 mg in 5 mL
DIMETHICONE (UNII: 92RU3N3Y10) (DIMETHICONE - UNII:92RU3N3Y10)	DIMETHICONE	40 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)	
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0K00R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	LEMON (Lemon)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49348-303-39	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2012	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	06/01/2012	03/01/2025

Labeler - STRATEGIC SOURCING SERVICES LLC (116956644)

Registrant - GCP Laboratories (965480861)

Establishment

Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(49348-303)

Revised: 11/2023

STRATEGIC SOURCING SERVICES LLC