

FEVERALL JR. STRENGTH- acetaminophen suppository
Taro Pharmaceuticals U.S.A. Inc.

FeverAll®
Jr. Strength

Drug Facts

Active ingredient (in each rectal suppository)

Acetaminophen, USP 325 mg

Purposes

Pain reliever/fever reducer

Uses

temporarily

- reduces fever
- relieves minor aches, pains, and headache

Warnings

Liver warning

This product contains acetaminophen.

Severe liver damage may occur if

- a child 6 to 12 years takes more than 5 doses in 24 hours
- an adult or child 12 years and older takes more than 6 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- an adult takes 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

For rectal use only.

Do not use

- in children under 6 years
- if you are allergic to acetaminophen.
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease.
- you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- fever lasts more than 3 days (72 hours), or recurs.
- pain gets worse or lasts more than 10 days.
- new symptoms occur.
- redness or swelling is present in the painful area.

These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical in case of overdose for adults and for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- remove wrapper
- carefully insert suppository well up into the rectum

Dosing Chart

| Age | Dose |
|--|--|
| under 6 years | do not use |
| 6 to 12 years | Use 1 suppository every 4 to 6 hours. (maximum of 5 doses in 24 hours) |
| adults and children 12 years and older | Use 2 suppositories every 4 to 6 hours. (maximum of 6 doses) |

Other information

- store at 2°-27°C (35°-80°F)
- do not use if imprinted suppository wrapper is opened or damaged

Inactive ingredients

glycerol monostearate, hydrogenated vegetable oil, polyoxyethylene stearate, polysorbate 80

Questions?

call 1-866-923-4914

Dist. by: **Taro Pharmaceuticals U.S.A., Inc.**
Hawthorne, NY 10532

PRINCIPAL DISPLAY PANEL - 325 mg Suppository Blister Pack Carton

NDC 51672-2116-2

RECTAL
SUPPOSITORY*

*actual size

Doctor Recommended

Pain Reliever/Fever Reducer

- No Parabens or Any Other Preservatives
- No Artificial Colors

JR. STRENGTH

ages 6-12 years

FeverAll®

ACETAMINOPHEN SUPPOSITORIES

Pain Reliever/Fever Reducer

6

Rectal

Suppositories

325

mg

each

NDC 51672-2116-2



*actual size

Doctor Recommended Pain Reliever/Fever Reducer

- ✓ No Parabens or Any Other Preservatives
- ✓ No Artificial Colors

JR. STRENGTH
ages 6-12 years

FeverAll®

ACETAMINOPHEN SUPPOSITORIES

Pain Reliever/Fever Reducer

JR. STRENGTH
ages 6-12 years

FeverAll®

ACETAMINOPHEN SUPPOSITORIES

Pain Reliever/Fever Reducer

JR. STRENGTH
ages 6-12 years

FeverAll®

ACETAMINOPHEN SUPPOSITORIES

Pain Reliever/Fever Reducer

6 Rectal Suppositories

325 mg each



Dist. by: Taro Pharmaceuticals U.S.A., Inc.
Hawthorne, NY 10532 Made in USA
GW 710012 TBD

Tamper-evident: Suppositories are individually wrapped. Do not use if imprinted wrapper is opened or damaged.



Other information

- store at 20°-25°C (35°-80°F)
- do not use if imprinted suppository wrapper is opened or damaged

Inactive ingredients

oil, polyoxyethylene stearate, poly sorbate 80, glycerol monostearate, hydrogenated vegetable

Questions? call 1-866-923-4914

As a doctor before use if

- you have liver disease.
- you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- fever lasts more than 3 days (72 hours), or recurs.
- pain gets worse or lasts more than 10 days.
- new symptoms occur.
- redness or swelling is present in the painful area.


These may be signs of a serious condition.

Ask a doctor before use if

- you are unsure whether a drug contains acetaminophen, ask a doctor or pharmacist.

(press upon or nonprescription, if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.)

6 doses) (years and older)



FEVERALL JR. STRENGTH

acetaminophen suppository

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:51672-2116 |
| Route of Administration | RECTAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|----------|
| Acetaminophen (UNII: 362O9ITL9D) (Acetaminophen - UNII:362O9ITL9D) | Acetaminophen | 325 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| glyceryl monostearate (UNII: 230OU9XXE4) | |
| hydrogenated palm kernel oil (UNII: FM8D1RE2VP) | |
| PEG-100 stearate (UNII: YD01N1999R) | |
| polysorbate 80 (UNII: 6OZP39ZG8H) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:51672-2116-2 | 6 in 1 CARTON | 12/12/2013 | |
| 1 | NDC:51672-2116-0 | 1 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |
| 2 | NDC:51672-2116-4 | 50 in 1 CARTON | 12/12/2013 | |

| | | | | |
|------------------------------|---|--|---------------------------|--|
| 2 | NDC:51672-2116-0 | 1 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |
| Marketing Information | | | | |
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | |
| NDA | NDA018337 | 12/12/2013 | | |

Labeler - Taro Pharmaceuticals U.S.A. Inc. (145186370)

| | | | |
|---------------------------|----------------|---------------|----------------------------|
| Establishment | | | |
| Name | Address | ID/FEI | Business Operations |
| G&W NC Laboratories, Inc. | | 079419931 | MANUFACTURE(51672-2116) |

Revised: 7/2019

Taro Pharmaceuticals U.S.A. Inc.