

## **MILK OF MAGNESIA- magnesium hydroxide suspension**

### **Major Pharmaceuticals**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **Major Pharmaceuticals Milk Of Magnesia Drug Facts**

#### **Active ingredient (in each 15 mL)**

Magnesium hydroxide 1200 mg

#### **Purpose**

Saline laxative

#### **Uses**

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

#### **Warnings**

##### **Ask a doctor before use if you have**

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

##### **Ask a doctor or pharmacist before use if you are**

taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

##### **Stop use and ask a doctor if**

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

##### **If pregnant or breast-feeding,**

ask a health professional before use.

##### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

#### **Directions**

- do not exceed the maximum recommended daily dose in a 24 hour period

- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- mL = milliliter

adults and children 12 years and older	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

**Other information**

- **each 15 mL contains:** calcium 20 mg and magnesium 595 mg
- does not meet USP requirements for preservative effectiveness
- store at 20°-25° C (68°-77° F)
- close cap tightly after use
- avoid freezing

**Inactive ingredient**

purified water

**Questions or comments?**

**1-800-616-2471**

**Principal Display Panel**

Milk of Magnesia

Saline Laxative

ORIGINAL FLAVOR

Effective overnight relief

Cramp free

Stimulant free

Sugar free

Compare to the Active Ingredient of Phillips'® Milk of Magnesia

12 FL OZ (355 mL)

Magnesium Hydroxide, 1200 mg per 15 mL



NDC 0904-6756-14

# Milk of Magnesia

Saline Laxative  
**ORIGINAL FLAVOR**

- Effective overnight relief
- Cramp free
- Stimulant free
- Sugar free

Compare to the Active Ingredient  
of Phillips'® Milk of Magnesia\*

12 FL OZ (355 mL)

Magnesium Hydroxide,  
1200 mg per 15 mL

: 39640 5C F1

## Drug Facts

**DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING.**

### Active ingredient (in each 15 mL)

Magnesium hydroxide 1200 mg.....Saline laxative

### Purpose

### Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

### Warnings

Ask a **doctor** before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a **doctor** or **pharmacist** before use if you are taking any other drug.

Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

**Stop use and ask a doctor if** ■ you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.

- you need to use a laxative for more than 1 week

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### Directions

- do not exceed the maximum recommended daily dose in a 24 hour period
- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided ■ mL = milliliter

adults and children 12 years and older	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

### Other information

- each 15 mL contains: calcium 20 mg and magnesium 595 mg
- does not meet USP requirements for preservative effectiveness
- store at 20°-25°C (68°-77°F) ■ close cap tightly after use ■ avoid freezing

**Inactive ingredient** purified water

**Questions or comments?** 1-800-616-2471

**Gluten Free**

Distributed by  
**MAJOR®**  
**PHARMACEUTICALS**  
17177 N Laurel Park Drive  
Suite 233  
Livonia, MI 48152  
M-05 REV. 04/18  
Re-Order No. 700957

\*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Phillips'® Milk of Magnesia.



: 39640 5C B1

## MILK OF MAGNESIA

magnesium hydroxide suspension

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0904-6756
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>MAGNESIUM HYDRO XIDE</b> (UNII: NBZ3Q Y004S) (MAGNESIUM CATION - UNII:T6 V3LHY8 38, HYDRO XIDE ION - UNII:9 159 UV38 1P)	MAGNESIUM HYDROXIDE	1200 mg in 15 mL

**Inactive Ingredients**

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

**Product Characteristics**

Color	WHITE (opaque)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-6756-14	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/12/2019	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	02/12/2019	

**Labeler** - Major Pharmaceuticals (191427277)

Revised: 2/2019

Major Pharmaceuticals