

**MIGRAINE RELIEF- acetaminophen, aspirin, caffeine tablet, film coated  
Walgreen Company**

-----  
**Walgreen Co. Migraine Relief Drug Facts**

**Active ingredients (in each tablet)**

Acetaminophen 250 mg

Aspirin 250 mg (NSAID\*)

Caffeine 65 mg

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever

Pain reliever aid

**Use**

- treats migraine

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 2 gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Medication overuse headache warning:** Headaches may worsen if this product is used for 10 or more days per month.

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause severe nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

### **Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

### **Ask a doctor before use if**

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have vomiting with migraine headache

## **Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for:
- diabetes
- gout
- arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

## **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

## **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

## **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## **Directions**

- **do not use more than directed**
- adults: take 2 gels with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 gels in 24 hours, unless directed by a doctor
- under 18 years of age: as a doctor

## **Other information**

- store at 20°-25°C (68°-77°F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

## **Inactive ingredients**

black iron oxide, carnauba wax, crospovidone, D&C yellow #10 aluminum lake, FD&C blue #1 aluminum lake, gelatin, glycerin, hypromellose, maltitol solution, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polyvinyl acetate phthalate, povidone, pregelatinized starch, propylene glycol, stearic acid, titanium dioxide

## **Questions or comments?**

**1-800-719-9260**

## **Package/Label Principal Display Panel**

Walgreens

WALGREENS PHARMACIST RECOMMENDED

Compare to the active ingredients in Excedrin<sup>®</sup> Migraine

Migraine Relief

ACETAMINOPHEN / ASPIRIN (NSAID) / CAFFIENE TABLETS

PAIN RELIEVER / PAIN RELIEVER AID

20 GELTABS\*\*

(\*\*gelatin-coated tablets)

ACTUAL SIZE



22560 94 C 1

ITEM 492352 W10106-0722-L  
 4922 Walgreens  
 WALGREENS COMPANY, 240 W. MILWAUKEE, DEERFIELD, IL 60015

©2012 Walgreens  
 This product is not manufactured or distributed by GSK Consumer Healthcare, owner of the registered trademark Excedrin®.  
 We invite you to compare to national brands.  
 Our pharmacists recommend the Walgreens brand.

**Drug Facts**

**Active ingredients (in each geltab)**      **Purposes**

Acetaminophen 250 mg      Pain reliever

Aspirin 250 mg (NSAID)<sup>1</sup>      Pain reliever

Caffeine 65 mg      Pain reliever aid

<sup>1</sup>nonsteroidal anti-inflammatory drug

**Use**      Treats migraine

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:      skin redness      blisters      rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:      hives      facial swelling      asthma (wheezing)      shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take      more than 2 geltabs in 24 hours, which is the maximum daily amount      with other drugs containing acetaminophen      3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you      are age 60 or older      have had stomach ulcers or bleeding problems      take a blood thinning (anticoagulant) or steroid drug      take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)      have 3 or more alcoholic drinks every day while using this product      take more or for longer time than directed

Medicines were used with the warning: Headaches may worsen if this product is used for 10 or more days per month.

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally rapid heartbeat.

**Drug Facts (continued)**

**Do not use**      if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer      with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if**      you have never had migraines diagnosed by a health professional      you have a headache that is different from your usual migraines      you have the worst headache of your life      you have fever and stiff neck      you have headaches beginning after or caused by head injury, exertion, coughing or bending      you experienced your first headache after the age of 50      you have daily headaches      you have a migraine as often as to require bed rest      you have liver disease      stomach bleeding warning applies to you      you have a history of stomach problems, such as heartburn      you have high blood pressure, heart disease, liver cirrhosis, or kidney disease      you are taking a diuretic      you have asthma      you have problems or serious side effects from taking pain relievers or fever reducers      you have vomiting with your migraine headache

**Ask a doctor or pharmacist before use if you are**      taking a prescription drug (for:      diabetes      gout      arthritis      under a doctor's care for any serious condition      taking any other drug      taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer


**Stop use and ask a doctor if**      an allergic reaction occurs. Seek medical help right away      you experience any of the following signs of stomach bleeding:      feel faint      vomit blood      have bloody or black stools      have stomach pain that does not get better      your migraine is not relieved or worsens after first dose      new or unexpected symptoms occur      ringing in the ears or loss of hearing occurs

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions**      do not use more than directed      adults take



## MIGRAINE RELIEF

acetaminophen, aspirin, caffeine tablet, film coated

### Product Information

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:0363-1235 |
| <b>Route of Administration</b> | ORAL           |                           |               |

### Active Ingredient/Active Moiety

| Ingredient Name   | Basis of Strength | Strength |
|---|-------------------|----------|
| <b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN     | 250 mg   |
| <b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)             | ASPIRIN           | 250 mg   |
| <b>CAFFEINE</b> (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)           | CAFFEINE          | 65 mg    |

### Inactive Ingredients

| Ingredient Name                                       | Strength |
|---|----------|
| <b>FERROSO FERRIC OXIDE</b> (UNII: XM0M87F357)        |          |
| <b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)                |          |
| <b>CROSPVIDONE</b> (UNII: 2S7830E561)                 |          |
| <b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)       |          |
| <b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)         |          |
| <b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)        |          |
| <b>GLYCERIN</b> (UNII: PDC6A3C0OX)                    |          |
| <b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)   |          |
| <b>MALTITOL</b> (UNII: D65DG142WK)                    |          |
| <b>MEDIUM-CHAIN TRIGLYCERIDES</b> (UNII: C9H2L21V7U)  |          |
| <b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)  |          |
| <b>POLYVINYL ACETATE PHTHALATE</b> (UNII: 58QVG85GW3) |          |
| <b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)       |          |
| <b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)            |          |
| <b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)                |          |
| <b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)            |          |

### Product Characteristics

|              |              |              |          |
|--------------|--------------|--------------|----------|
| <b>Color</b> | GREEN, WHITE | <b>Score</b> | no score |
| <b>Shape</b> | ROUND        | <b>Size</b>  | 13mm     |

| <b>Flavor</b>                |  | <b>Imprint Code</b>                               | M                    |                    |
|------------------------------|--|---|----------------------|--------------------|
| <b>Contains</b>              |  |   |                      |                    |
| <b>Packaging</b>             |  |   |                      |                    |
| #                            | Item Code                                | Package Description                               | Marketing Start Date | Marketing End Date |
| 1                            | NDC:0363-1235-60                         | 1 in 1 CARTON                                     | 06/16/2022           |                    |
| 1                            |  | 20 in 1 BOTTLE; Type 0: Not a Combination Product |                      |                    |
| <b>Marketing Information</b> |  |   |                      |                    |
| Marketing Category           | Application Number or Monograph Citation | Marketing Start Date                              | Marketing End Date   |                    |
| ANDA                         | ANDA075794                               | 06/16/2022  |                      |                    |

**Labeler** - Walgreen Company (008965063)

Revised: 6/2022

Walgreen Company