

**COLD, FLU AND SORE THROAT- acetaminophen, dextromethorphan hbr,
guaifenesin, phenylephrine hcl solution
Walgreen Company**

Walgreens 44-005

Active ingredients (in each 20 mL)

Acetaminophen 650 mg
Dextromethorphan HBr 20 mg
Guaifenesin 400 mg
Phenylephrine HCl 10 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Expectorant
Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - minor aches and pains
 - sinus congestion and pressure
 - stuffy nose
 - nasal congestion
 - sore throat
 - cough
 - headache
- temporarily promotes nasal and/or sinus drainage
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- rash
- blisters

- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- liver disease
- heart disease
- diabetes
- difficulty in urination due to enlargement of the prostate gland
- high blood pressure
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- thyroid disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- new symptoms occur
- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- do not take more than 6 doses in any 24-hour period
- mL = milliliter
- only use the dose cup provided
- dose as follows or as directed by a doctor
- adults and children 12 years and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years: do not use

Other information

- **each 20 mL contains:** sodium 10 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

anhydrous citric acid, FD&C blue #1, FD&C red #40, flavors, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sorbitol, sucralose

Questions or comments?

1-800-426-9391

Principal Display Panel

NDC 0363-8005-45

Walgreens

WALGREENS PHARMACIST RECOMMENDED[†]

Compare to the active ingredients
in Maximum Strength Mucinex[®]
FAST-MAX[®] Cold, Flu & Sore Throat^{††}

Cold, Flu

& Sore Throat

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER

DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT

GUAIFENESIN / EXPECTORANT

PHENYLEPHRINE HCl / NASAL DECONGESTANT

Maximum Strength

- Relieves body pain, headache,

fever & sore throat

- Controls cough
- Relieves nasal & chest congestion
- Thins & loosens mucus
- Ages 12 years & over

6 FL OZ (177 mL)

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

50844 ORG052300545

DISTRIBUTED BY:

WALGREEN CO.

DEERFIELD, IL 60015

100% SATISFACTION

GUARANTEED

walgreens.com

©2023 Walgreen Co.

†Our pharmacists recommend the Walgreens brand.

We invite you to compare to national brands.

††This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademark Maximum Strength Mucinex® FAST-MAX® Cold, Flu & Sore Throat.



PARENTS:
Learn how to best protect your child.
www.DrugFactsOnline.com

50844 ORG052300545
 B-005-45 ORG

DISTRIBUTED BY:
 WALGREEN CO.
 DEERFIELD, IL 60015
 100% SATISFACTION
 GUARANTEED
 walgreens.com
 ©2023 Walgreen Co.

ITEM 769447 W00000-0000-0

3 11917 03411 9

W3ORG0923-F

Our pharmacists recommend the Walgreens brand.
 We invite you to compare to national brands.
 ††This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademark Maximum Strength Mucinex® FAST-MAX® Cold, Flu & Sore Throat.

Drug Facts TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Active ingredients (in each 20 mL) **Purpose**
 Acetaminophen 650 mg.....Pain reliever/fever reducer
 Dextromethorphan HBr 20 mg.....Cough suppressant
 Guaifenesin 400 mg.....Expectorant
 Phenylephrine HCl 10 mg.....Nasal decongestant

Uses ■ temporarily relieves these common cold and flu symptoms:
 ■ minor aches and pains
 ■ sinus congestion and pressure
 ■ stuffy nose ■ nasal congestion
 ■ sore throat ■ cough ■ headache
 ■ temporarily promotes nasal and/or sinus drainage
 ■ temporarily reduces fever
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**No Print / No Varnish Area
 Lot # and Exp. Info**

Drug Facts (continued)

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ rash ■ blisters ■ skin reddening. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
 ■ cough that occurs with too much phlegm (mucus)
 ■ liver disease ■ heart disease ■ diabetes
 ■ difficulty in urination due to enlargement of the prostate gland ■ high blood pressure ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ thyroid disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Drug Facts (continued)

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if
 ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ new symptoms occur
 ■ nervousness, dizziness, or sleeplessness occur
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed
 ■ do not take more than 6 doses in any 24-hour period
 ■ mL = milliliter ■ only use the dose cup provided
 ■ dose as follows or as directed by a doctor
 ■ adults and children 12 years and over: 20 mL in dosing cup provided every 4 hours
 ■ children under 12 years: do not use

Other information ■ each 20 mL contains: sodium 10 mg ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package

Inactive ingredients anhydrous citric acid, FD&C blue #1, FD&C red #40, flavors, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sorbitol, sucralose

(Questions or comments?) 1-800-426-9391

Walgreens 44-005

COLD, FLU AND SORE THROAT

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-8005
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1VS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	blue	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-8005-45	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/04/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/04/2023	

Labeler - Walgreen Company (008965063)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	manufacture(0363-8005) , pack(0363-8005)

Revised: 10/2023

Walgreen Company