

COLD MULTI-SYMPTOM DAYTIME AND NIGHTTIME- acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, and phenylephrine hydrochloride
GoodSense

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

GDS - 1150 - 2022-1118

COLD MAX DAY

Drug Facts

<i>Active ingredients (in each caplet)</i>	<i>Purpose</i>
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these common cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - cough
 - sinus congestion and pressure
- helps clear nasal passages
- promotes nasal and sinus drainage
- temporarily reduces fever

COLD MAX NIGHT

Drug Facts

<i>Active ingredients (in each caplet)</i>	<i>Purpose</i>
Acetaminophen 325 mg	Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg	Antihistamine
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these common cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - runny nose and sneezing
 - cough
 - sinus congestion and pressure
- helps clear nasal passages
- relieves cough to help you sleep
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dosage**

In addition, when using Cold Max Night:

- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
 - do not take Day and Night caplets at the same time
 - do not take more than a total of 10 caplets in 24 hours
-

adults and children 12 years and over	<ul style="list-style-type: none">▪ take 2 caplets every 4 hours▪ swallow whole – do not crush, chew, or dissolve
children under 12 years	<ul style="list-style-type: none">▪ ask a doctor

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information and warnings

Inactive ingredients

Cold Max Day

acesulfame potassium, colloidal silicon dioxide, croscarmellose sodium, crospovidone, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

Cold Max Night

acesulfame potassium, colloidal silicon dioxide, croscarmellose sodium, FD&C blue #1 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

PRINCIPAL DISPLAY PANEL

GoodSense®

NDC 50804-551-02

For Adults

Cold Max Multi-Symptom

Cool Taste

Instant Cooling Sensation

Acetaminophen, Phenylephrine HCl, Dextromethorphan HBr, Chlorpheniramine Maleate*

Pain Reliever/Fever Reducer, Nasal Decongestant, Cough Suppressant, Antihistamine*

Day

- Head & Body Aches
- Fever & Sore Throat
- Cough
- Nasal Congestion

12 CAPLETS

Night

- Head & Body Aches
- Fever & Sore Throat
- Cough
- Nasal Congestion
- Runny Nose*

*Antihistamine in Nighttime Only

8 CAPLETS

LIFT PANEL FOR MORE DRUG FACTS INFORMATION

<p>Do not use</p> <ul style="list-style-type: none"> ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> ■ skin reddening ■ blisters ■ rash <p>■ 3 or more alcoholic drinks every day while using this product</p> <p>■ more than 4,000 mg of acetaminophen in 24 hours</p> <p>■ with other drugs containing acetaminophen</p> <p>Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take</p>	
<p>Warnings</p> <p>temporarily relieves these common cold/flu symptoms:</p> <ul style="list-style-type: none"> ■ minor aches and pains ■ headache ■ sore throat ■ nasal congestion ■ cough ■ sinus congestion and pressure ■ helps clear nasal passages ■ promotes nasal and sinus drainage ■ temporarily reduces fever 	
<p>Uses</p> <p>temporarily relieves these common cold/flu symptoms:</p> <ul style="list-style-type: none"> ■ minor aches and pains ■ headache ■ sore throat ■ nasal congestion ■ runny nose and sneezing ■ cough ■ sinus congestion and pressure ■ helps clear nasal passages ■ relieves cough to help you sleep ■ temporarily reduces fever 	
<p>Drug Facts</p> <p>Active ingredients</p> <p>(in each Day caplet)</p> <p>Acetaminophen 325 mg...Pain reliever/fever reducer</p> <p>Chlorpheniramine maleate 2 mg.....Antihistamine</p> <p>Dextromethorphan HBr 10 mg.....Cough suppressant</p> <p>Phenylephrine HCl 5 mg.....Nasal decongestant</p>	<p>Drug Facts</p> <p>Active ingredients</p> <p>(in each Night caplet)</p> <p>Acetaminophen 325 mg...Pain reliever/fever reducer</p> <p>Chlorpheniramine maleate 2 mg.....Antihistamine</p> <p>Dextromethorphan HBr 10 mg.....Cough suppressant</p> <p>Phenylephrine HCl 5 mg.....Nasal decongestant</p>
<p>Purpose</p> <p>temporarily relieves these common cold/flu symptoms:</p> <ul style="list-style-type: none"> ■ minor aches and pains ■ headache ■ sore throat ■ nasal congestion ■ runny nose and sneezing ■ cough ■ sinus congestion and pressure ■ helps clear nasal passages ■ relieves cough to help you sleep ■ temporarily reduces fever 	

DO NOT TAKE THE DAY AND NIGHT CAPLETS AT THE SAME TIME.
Do not take more than a total of 10 caplets in a 24-hour period. Take only as directed.

COLD MAX DAY **COLD MAX NIGHT**

GOODSENSE.
Cold Max Day & Night
Multi-Symptom For Adults

GOODSENSE.

NDC 50804-551-02

For Adults

Cold Max Cool Taste Instant Cooling Sensation

Multi-Symptom

**Acetaminophen, Phenylephrine HCl,
Dextromethorphan HBr, Chlorpheniramine Maleate***

**Pain Reliever/Fever Reducer, Nasal Decongestant,
Cough Suppressant, Antihistamine***

Day

- Head & Body Aches
- Fever & Sore Throat
- Cough
- Nasal Congestion



Actual Size

Night

- Head & Body Aches
- Fever & Sore Throat
- Cough
- Nasal Congestion
- Runny Nose*



Actual Size

* Antihistamine in Night Only

12 CAPLETS

8 CAPLETS

NC

NC

Drug Facts (continued)

Inactive ingredients
Cold Max Day acetaminophen potassium, colloidal silicon dioxide, corn starch, croscarmellose sodium, croscopolone, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

Cold Max Night acetaminophen potassium, colloidal silicon dioxide, corn starch, croscarmellose sodium, croscopolone, FD&C blue #1 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

Other information ■ store between 20-25°C (68-77°F) in a dry place ■ retain carton for complete product information and warnings

Directions ■ do not take more than directed (see overdose warning) ■ do not take Day and Night caplets at the same time ■ do not take more than a total of 10 caplets in 24 hours

adults and children ■ take 2 caplets every 4 hours ■ swallow whole; do not crush, chew, or dissolve

children under 12 years ■ ask a doctor

Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occur ■ path, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

When using this product ■ do not exceed recommended dosage ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers

Ask a doctor or pharmacist before use if you are ■ a breathing problem such as emphysema or chronic bronchitis ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ trouble urinating due to an enlarged prostate gland ■ cough that occurs with too much phlegm (mucus)

In addition, when using Cold Max Night ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ avoid alcoholic drinks


Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occur ■ path, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

DO NOT USE IF BUSTER UNITS ARE TORN OR BROKEN

Distributed by:
Perrigo Direct, Inc.
Peachtree City, GA 30269
www.PerrigoDirect.com
1-844-705-4384
GoodSense® is a registered trademark of L. Perrigo Company





COLD MULTI-SYMPATOM DAYTIME AND NIGHTTIME

acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, and phenylephrine hydrochloride kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50804-551
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50804-551-02	1 in 1 CARTON	12/31/2011	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BLISTER PACK	12
Part 2	1 BLISTER PACK	8

Part 1 of 2

ACETAMINOPHEN, DEXTROMETHORPHAN HYDROBROMIDE, PHENYLEPHRINE HYDROCHLORIDE

acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride tablet, coated

Product Information

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH)	DEXTROMETHORPHAN	10 mg

(DEXTROMETHORPHAN - UNII:7355X3ROTS)	HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL (capsule-shaped)	Size	17mm
Flavor	MINT	Imprint Code	AAA;1138
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341		

Part 2 of 2

ACETAMINOPHEN, CHLORPHENIRAMINE MALEATE, DEXTROMETHORPHAN HYDROBROMIDE, PHENYLEPHRINE HYDROCHLORIDE

acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, and phenylephrine

hydrochloride tablet, coated

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
MAGNESIUM STEARATE (UNII: 70097M6130)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	blue	Score	no score
Shape	OVAL (capsule-shaped)	Size	17mm
Flavor	MINT	Imprint Code	AAA;1139
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	12/31/2011	

Labeler - GoodSense (076059836)

Revised: 11/2022

GoodSense