


CHLOROTHIAZIDE SODIUM- chlorothiazide sodium injection, powder, lyophilized, for solution
Sagent Pharmaceuticals

Chlorothiazide Sodium for Injection, USP
(For Intravenous Use)

SAGENT®
Rx only


DESCRIPTION

Chlorothiazide Sodium for Injection, USP is a diuretic and antihypertensive. It is 6-chloro-2*H*-1,2,4-benzothiadiazine-7-sulfonamide 1,1-dioxide monosodium salt and its molecular weight is 317.71. Its empirical formula is $C_7H_5ClN_3NaO_4S_2$ and its structural formula is:



Chlorothiazide Sodium for Injection, USP is a sterile lyophilized white powder and is supplied in a vial containing: Chlorothiazide sodium equivalent to chlorothiazide 500 mg, and the inactive ingredient mannitol 250 mg with sodium hydroxide to adjust pH.

Chlorothiazide is a diuretic and antihypertensive. It is 6-chloro-2*H*-1,2,4-benzothiadiazine-7-sulfonamide 1,1-dioxide. Its empirical formula is $C_7H_6ClN_3O_4S_2$ and its structural formula is:



It is a white, or practically white, crystalline powder with a molecular weight of 295.72, which is very slightly soluble in water, but readily soluble in dilute aqueous sodium hydroxide. It is soluble in urine to the extent of about 150 mg per 100 mL at pH 7.

CLINICAL PHARMACOLOGY

The mechanism of the antihypertensive effect of thiazides is unknown. Chlorothiazide does not usually affect normal blood pressure.

Chlorothiazide affects the distal renal tubular mechanism of electrolyte reabsorption. At maximal therapeutic dosage all thiazides are approximately equal in their diuretic efficacy.

Chlorothiazide increases excretion of sodium and chloride in approximately equivalent amounts. Natriuresis may be accompanied by some loss of potassium and bicarbonate.

After oral use diuresis begins within 2 hours, peaks in about 4 hours and lasts about 6 to 12 hours. Following intravenous use of chlorothiazide sodium, onset of the diuretic action occurs in 15 minutes and the maximal action in 30 minutes.

Pharmacokinetics and Metabolism

Chlorothiazide is not metabolized but is eliminated rapidly by the kidney; 96 percent of an intravenous dose is excreted unchanged in the urine within 23 hours. The plasma half-life of chlorothiazide is 45 to 120 minutes. Chlorothiazide crosses the placental but not the blood-brain barrier and is excreted in breast milk.

INDICATIONS AND USAGE

Chlorothiazide Sodium for Injection, USP is indicated as adjunctive therapy in edema associated with congestive heart failure, hepatic cirrhosis, and corticosteroid and estrogen therapy.

Chlorothiazide Sodium for Injection, USP has also been found useful in edema due to various forms of renal dysfunction such as nephrotic syndrome, acute glomerulonephritis, and chronic renal failure.

Use in Pregnancy

Routine use of diuretics during normal pregnancy is inappropriate and exposes mother and fetus to unnecessary hazard. Diuretics do not prevent development of toxemia of pregnancy and there is no satisfactory evidence that they are useful in the treatment of toxemia.

Edema during pregnancy may arise from pathologic causes or from the physiologic and mechanical consequences of pregnancy. Thiazides are indicated in pregnancy when edema is due to pathologic causes, just as they are in the absence of pregnancy (see **PRECAUTIONS, Pregnancy**). Dependent edema in pregnancy, resulting from restriction of venous return by the gravid uterus, is properly treated through elevation of the lower extremities and use of support stockings. Use of diuretics to lower intravascular volume in this instance is illogical and unnecessary. During normal pregnancy there is hypervolemia which is not harmful to the fetus or the mother in the absence of cardiovascular disease. However, it may be associated with edema, rarely generalized edema. If such edema causes discomfort, increased recumbency will often provide relief. Rarely this edema may cause extreme discomfort which is not relieved by rest. In these instances, a short course of diuretic therapy may provide relief and be appropriate.

CONTRAINDICATIONS

Anuria.

Hypersensitivity to any component of this product or to other sulfonamide-derived drugs.

WARNINGS

Intravenous use in infants and children has been limited and is not generally recommended.

Use with caution in severe renal disease. In patients with renal disease, thiazides may precipitate azotemia. Cumulative effects of the drug may develop in patients with impaired renal function.

Thiazides should be used with caution in patients with impaired hepatic function or progressive liver disease, since minor alterations of fluid and electrolyte balance may precipitate hepatic coma.

Thiazides may add to or potentiate the action of other antihypertensive drugs.

Sensitivity reactions may occur in patients with or without a history of allergy or bronchial asthma.

The possibility of exacerbation or activation of systemic lupus erythematosus has been reported.

Lithium generally should not be given with diuretics (see **PRECAUTIONS, Drug Interactions**).

PRECAUTIONS

General

All patients receiving diuretic therapy should be observed for evidence of fluid or electrolyte imbalance: namely, hyponatremia, hypochloremic alkalosis, and hypokalemia. Serum and urine electrolyte determinations are particularly important when the patient is vomiting excessively or receiving parenteral fluids. Warning signs or symptoms of fluid and electrolyte imbalance, irrespective of cause, include dryness of mouth, thirst, weakness, lethargy, drowsiness, restlessness, confusion, seizures, muscle pains or cramps, muscular fatigue, hypotension, oliguria, tachycardia, and gastrointestinal disturbances such as nausea and vomiting.

Hypokalemia may develop especially with brisk diuresis, when severe cirrhosis is present or after prolonged therapy.

Interference with adequate oral electrolyte intake will also contribute to hypokalemia. Hypokalemia may cause cardiac arrhythmias and may also sensitize or exaggerate the response of the heart to the toxic effects of digitalis (e.g., increased ventricular irritability). Hypokalemia may be avoided or treated by use of potassium-sparing diuretics or potassium supplements such as foods with a high potassium content.

Although any chloride deficit is generally mild and usually does not require specific treatment except under extraordinary circumstances (as in liver disease or renal disease), chloride replacement may be required in the treatment of metabolic alkalosis.

Dilutional hyponatremia may occur in edematous patients in hot weather; appropriate

therapy is water restriction, rather than administration of salt, except in rare instances when the hyponatremia is life-threatening. In actual salt depletion, appropriate replacement is the therapy of choice.

Hyperuricemia may occur or acute gout may be precipitated in certain patients receiving thiazides.

In diabetic patients dosage adjustments of insulin or oral hypoglycemic agents may be required. Hyperglycemia may occur with thiazide diuretics. Thus latent diabetes mellitus may become manifest during thiazide therapy.

The antihypertensive effects of the drug may be enhanced in the postsympathectomy patient.

If progressive renal impairment becomes evident, consider withholding or discontinuing diuretic therapy.

Thiazides have been shown to increase the urinary excretion of magnesium; this may result in hypomagnesemia.

Thiazides may decrease urinary calcium excretion. Thiazides may cause intermittent and slight elevation of serum calcium in the absence of known disorders of calcium metabolism. Marked hypercalcemia may be evidence of hidden hyperparathyroidism. Thiazides should be discontinued before carrying out tests for parathyroid function.

Increases in cholesterol and triglyceride levels may be associated with thiazide diuretic therapy.

Laboratory Tests

Periodic determination of serum electrolytes to detect possible electrolyte imbalance should be done at appropriate intervals.

Drug Interactions

When given concurrently the following drugs may interact with thiazide diuretics.

Alcohol, barbiturates, or narcotics - potentiation of orthostatic hypotension may occur.

Antidiabetic drugs - (oral agents and insulin) - dosage adjustment of the antidiabetic drug may be required.

Other antihypertensive drugs - additive effect or potentiation.

Corticosteroids, ACTH - intensified electrolyte depletion, particularly hypokalemia.

Pressor amines (e.g., norepinephrine) - possible decreased response to pressor amines but not sufficient to preclude their use.

Skeletal muscle relaxants, nondepolarizing (e.g., tubocurarine) - possible increased responsiveness to the muscle relaxant.

Lithium - generally should not be given with diuretics. Diuretic agents reduce the renal clearance of lithium and add a high risk of lithium toxicity. Refer to the package insert for lithium preparations before use of such preparations with chlorothiazide sodium.

Non-steroidal Anti-inflammatory Drugs - In some patients, the administration of a non-steroidal anti-inflammatory agent can reduce the diuretic, natriuretic, and

antihypertensive effects of loop, potassium-sparing and thiazide diuretics. Therefore, when chlorothiazide sodium and non-steroidal anti-inflammatory agents are used concomitantly, the patient should be observed closely to determine if the desired effect of the diuretic is obtained.

Drug/Laboratory Test Interactions

Thiazides should be discontinued before carrying out tests for parathyroid function (see **PRECAUTIONS, General**).

Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenicity studies have not been conducted with chlorothiazide.

Chlorothiazide was not mutagenic *in vitro* in the Ames microbial mutagen test (using a maximum concentration of 5 mg/plate and *Salmonella typhimurium* strains TA98 and TA100) and was not mutagenic and did not induce mitotic nondisjunction in diploid-strains of *Aspergillus nidulans*.

Chlorothiazide had no adverse effects on fertility in female rats at doses up to 60 mg/kg/day and no adverse effects on fertility in male rats at doses up to 40 mg/kg/day. These doses are 1.5 and 1.0 times** the recommended maximum human dose, respectively, when compared on a body weight basis.

Pregnancy

Teratogenic Effects - Pregnancy Category C: Although reproduction studies performed with chlorothiazide doses of 50 mg/kg/day in rabbits, 60 mg/kg/day in rats and 500 mg/kg/day in mice revealed no external abnormalities of the fetus or impairment of growth and survival of the fetus due to chlorothiazide, such studies did not include complete examinations for visceral and skeletal abnormalities. It is not known whether chlorothiazide can cause fetal harm when administered to a pregnant woman; however, thiazides cross the placental barrier and appear in cord blood. Chlorothiazide should be used during pregnancy only if clearly needed (see **INDICATIONS AND USAGE**).

Nonteratogenic Effects: Chlorothiazide may cause fetal or neonatal jaundice, thrombocytopenia, and possibly other adverse reactions which have occurred in the adult.

Nursing Mothers

Because of the potential for serious adverse reactions in nursing infants from chlorothiazide sodium for injection, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use

Safety and effectiveness of chlorothiazide sodium for injection in pediatric patients have not been established.

Geriatric Use

Clinical studies of chlorothiazide sodium for injection did not include sufficient numbers

of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function (see **WARNINGS**).

ADVERSE REACTIONS

The following adverse reactions have been reported and, within each category, are listed in order of decreasing severity.

Body as a Whole: Weakness.

Cardiovascular: Hypotension including orthostatic hypotension (may be aggravated by alcohol, barbiturates, narcotics or antihypertensive drugs).

Digestive: Pancreatitis, jaundice (intrahepatic cholestatic jaundice), diarrhea, vomiting, sialadenitis, cramping, constipation, gastric irritation, nausea, anorexia.

Hematologic: Aplastic anemia, agranulocytosis, leukopenia, hemolytic anemia, thrombocytopenia.

Hypersensitivity: Anaphylactic reactions, necrotizing angitis (vasculitis and cutaneous vasculitis), respiratory distress including pneumonitis and pulmonary edema, photosensitivity, fever, urticaria, rash, purpura.

Metabolic: Electrolyte imbalance (see **PRECAUTIONS**), hyperglycemia, glycosuria, hyperuricemia.

Musculoskeletal: Muscle spasm.

Nervous System/Psychiatric: Vertigo, paresthesias, dizziness, headache, restlessness.

Skin: Erythema multiforme including Stevens-Johnson syndrome, exfoliative dermatitis including toxic epidermal necrolysis, alopecia.

Special Senses: Transient blurred vision, xanthopsia.

Renal: Renal failure, renal dysfunction, interstitial nephritis, (see **WARNINGS**); hematuria (following intravenous use).

Urogenital: Impotence.

Whenever adverse reactions are moderate or severe, thiazide dosage should be reduced or therapy withdrawn.

To report SUSPECTED ADVERSE REACTIONS, contact Sagent Pharmaceuticals, Inc. at 1-866-625-1618 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

OVERDOSAGE

The most common signs and symptoms observed are those caused by electrolyte depletion (hypokalemia, hypochloremia, hyponatremia) and dehydration resulting from excessive diuresis. If digitalis has also been administered, hypokalemia may accentuate cardiac arrhythmias.

In the event of overdosage, symptomatic and supportive measures should be employed. Correct dehydration, electrolyte imbalance, hepatic coma and hypotension by established procedures. If required, give oxygen or artificial respiration for respiratory impairment.

The degree to which chlorothiazide sodium is removed by hemodialysis has not been established.

The intravenous LD₅₀ of chlorothiazide in the mouse is 1.1 g/kg.

DOSAGE AND ADMINISTRATION

Chlorothiazide Sodium for Injection, USP should be reserved for patients unable to take oral medication or for emergency situations.

Therapy should be individualized according to patient response. Use the smallest dosage necessary to achieve the required response.

Intravenous use in infants and children has been limited and is not generally recommended.

When medication can be taken orally, therapy with chlorothiazide tablets or oral suspension may be substituted for intravenous therapy, using the same dosage schedule as for the parenteral route.

Chlorothiazide Sodium for Injection, USP may be given slowly by direct intravenous injection or by intravenous infusion.

Extravasation must be rigidly avoided. Do not give subcutaneously or intramuscularly.

The usual adult dosage is 500 mg to 1 g once or twice a day. Many patients with edema respond to intermittent therapy, i.e., administration on alternate days or on three to five days each week. With an intermittent schedule, excessive response and the resulting undesirable electrolyte imbalance are less likely to occur.

Directions for Reconstitution

Use aseptic technique. Because Chlorothiazide Sodium for Injection, USP contains no preservative, a fresh solution should be prepared immediately prior to each administration, and the unused portion should be discarded.

Add 18 mL of Sterile Water for Injection to the vial to form an isotonic solution for intravenous injection. Never add less than 18 mL. When reconstituted with 18 mL of Sterile Water, the final concentration of Chlorothiazide Sodium for Injection, USP is 28 mg per mL. The reconstituted solution is clear and essentially free from visible particles. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to use whenever solution and container permit. The solution is

compatible with dextrose or sodium chloride solutions for intravenous infusion. Avoid simultaneous administration of solutions of chlorothiazide with whole blood or its derivatives.

HOW SUPPLIED

Chlorothiazide Sodium for Injection, USP is supplied as follows:

NDC	Chlorothiazide Sodium for Injection, USP	Package Factor
25021-305-20	500 mg Single-Dose Vial	1 vial per carton

Chlorothiazide Sodium for Injection, USP is a dry, sterile lyophilized white powder usually in cake form, supplied in vials containing chlorothiazide sodium equivalent to 500 mg of chlorothiazide.

Storage Conditions

Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]

For single dose only. Use solution immediately after reconstitution (see **DOSAGE AND ADMINISTRATION, Directions for Reconstitution**). Discard unused portion of the reconstituted solution.

Sterile, Nonpyrogenic, Preservative-free.

The container closure is not made with natural rubber latex.

** Calculations based on a human body weight of 50 kg.

SAGENT®

Mfd. by SAGENT Pharmaceuticals
Schaumburg, IL 60195
Made in USA

©2020 Sagent Pharmaceuticals, Inc.

May 2020

SAGENT Pharmaceuticals®

PACKAGE LABEL – PRINCIPAL DISPLAY PANEL – Vial Label

NDC 25021-305-20

Rx only

CHLOROTHIAZIDE SODIUM for Injection, USP

500 mg per vial

FOR THE PREPARATION OF INTRAVENOUS SOLUTIONS

For Intravenous Use

Single-Dose Vial

NDC 25021-305-20

CHLOROTHIAZIDE SODIUM
for Injection, USP

500 mg*
per vial

FOR THE PREPARATION
OF INTRAVENOUS SOLUTIONS

**For Intravenous Use
Single-Dose Vial**

R_xonly

Sterile, Nonpyrogenic, Preservative-free.
This container closure is not made with natural rubber latex.

***Each vial contains:** chlorothiazide sodium equivalent to 500 mg chlorothiazide, 250 mg mannitol, sodium hydroxide to adjust pH.

Usual Dosage: See package insert for dosage information.

Store at 20° to 25°C (68° to 77°F).
[See USP Controlled Room Temperature.]
Reconstitute with a minimum of 18 mL of Sterile Water for Injection. (An isotonic solution is obtained if 18 mL is added.)
For single-dose only. Use reconstituted solution **immediately**. Discard unused portion.

SAGENT®
Mfd. by SAGENT Pharmaceuticals
Schaumburg, IL 60195
Made in USA
©2021 Sagent Pharmaceuticals, Inc.

626142



(01)00325021305201

Lot: _____
Exp.: _____

CHLOROTHIAZIDE SODIUM

chlorothiazide sodium injection, powder, lyophilized, for solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:25021-305
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
chlorothiazide sodium (UNII: SN86FG7N2K) (chlorothiazide - UNII: 77W477J15H)	chlorothiazide	500 mg in 18 mL

Inactive Ingredients

Ingredient Name	Strength
mannitol (UNII: 3OWL53L36A)	
sodium hydroxide (UNII: 55X04QC32I)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:25021-305-20	1 in 1 CARTON	10/12/2015	
1		18 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202462	10/12/2015	

Labeler - Sagent Pharmaceuticals (796852890)

Revised: 5/2021

Sagent Pharmaceuticals