FLEET- glycerin liquid C.B. Fleet Company, Inc.

-----

Fleet Glycerin Liquid 0132-018

## **Drug Facts**

## Active Ingredient (in each 5.4 mL average delivered dose)

Glycerin 5.4 g

## **Purpose**

Hyperosmotic Laxative

#### Use

for relief of occasional constipation.

This product usually produces a bowel movement in 15 minutes to 1 hour.

## **Warnings**

May cause rectal discomfort or a burning sensation.

For rectal use only.

Do not use when abdominal pain, nausea, or vomiting are present unless directed by a doctor.

# Ask a doctor before using any laxative if you

- have a sudden change in bowel habits lasting more than 2 weeks
- already used a laxative for more than 1 week

# Stop using this product and consult a doctor if you have

- rectal bleeding
- no bowel movement within 1 hour of using this product

These symptoms may indicate a serious condition.

# Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

### **Directions**

Single Daily Dosage (per 24 hours)

Adults and children 6 years and over 1 suppository or as directed by a doctor

Children 2 to under 6 years use Fleet® Pedia-Lax® Liquid

Children under 2 years consult a doctor

**Left-side position:** Lie on left side with right knee bent and arms resting comfortably.

**Knee-chest position:** Kneel, then lower head and chest forward until left side of face is resting on surface with left arm folded comfortably

### Administering the laxative:

CAUTION: REMOVE ORANGE PROTECTIVE SHIELD BEFORE INSERTING.

Hold unit upright, grasping bulb of unit with fingers. Grasp orange protective shield with other hand, pull gently to remove. With steady pressure, insert tip into rectum with a slight side-to-side movement, with tip pointing toward navel. Do not force the applicator tip into the rectum, as this can cause injury. Insertion may be easier if person receiving suppository bears down as if having a bowel movement. This helps relax the muscles around the anus. Squeeze the bulb until nearly all liquid is expelled. While continuing to squeeze the bulb, remove tip from rectum and discard unit. It is not necessary to empty unit completely. The unit contains more than the amount of liquid needed for effective use. A small amount of liquid will remain in the unit after use.

For a more detailed step-by-step instructional video on how to use Fleet<sup>®</sup> products, visit our website **www.fleetlabs.com** to watch the "how to use" video.

### **Inactive Ingredients**

edetate disodium, purified water

#### Questions?

1-866-255-6960 or www.fleetlabs.com

#### PRINCIPAL DISPLAY PANEL

Fleet Laxative
Liquid Glycerin Suppositories
4Adult Suppositories
7.5 ml each



## **FLEET**

glycerin liquid

Product	Inform	ation
Product	Intorm	ation

Product Type HUMAN OTC DRUG Item Code (Source) NDC:0132-0185

**Route of Administration** RECTAL

### **Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
GLYCERIN (UNII: PDC6A3C0OX) (GLYCERIN - UNII:PDC6A3C0OX)	GLYCERIN	5.4 a

Inactive Ingredients			
Ingredient Name	Strength		
EDETATE DISODIUM (UNII: 7FLD91C86K)			
WATER (UNII: 059QF0KO0R)			

l	Packaging					
	# Item Code	Package Description	Marketing Start Date	Marketing End Date		
	NDC:0132-0185-	4 in 1 CARTON; Type 0: Not a Combination Product	06/03/2002			
	NDC:0132-0185- 01	1 in 1 PACKAGE; Type 0: Not a Combination Product	06/03/2002	08/31/2022		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M007	06/03/2002		

Labeler - C.B. Fleet Company, Inc. (003119054)

Revised: 5/2024 C.B. Fleet Company, Inc.