

**TOPCO TUSSIN CF MAX- acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl solution**

**Topco Associates LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Topco Associates LLC. Tussin CF Max Drug Facts**

**Active ingredients (in each 20 mL)**

Acetaminophen, USP 650 mg

Dextromethorphan HBr, USP 20 mg

Guaifenesin, USP 400 mg

Phenylephrine HCl, USP 10 mg

**Purposes**

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

**Uses**

- temporarily relieves these symptoms occurring with a cold or flu:
- cough due to minor throat and bronchial irritation
- nasal congestion
- sinus congestion and pressure
- minor aches and pains
- sore throat
- headache
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen

- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

### **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

### **When using this product**

**do not use more than directed**

### **Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days

- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than 5 doses in any 24-hour period
- do not exceed recommended dosage. Taking more than the recommended dose (overdose) may cause serious liver damage.
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 mL every 4 hours
children under 12 years	do not use

**Other information**

- **each 20 mL contains:** sodium 15 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients**

anhydrous citric acid, benzyl alcohol, edetate disodium, FD&C red #40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, triacetin, xanthan gum

**Questions or comments?**

**1-888-423-0139**

**Package/Label Principal Display Panel**

TopCare® health

COMPARE TO ROBITUSSIN® MAXIMUM STRENGTH SEVERE MULTI-SYMPTOM  
COUGH COLD + FLU ACTIVE INGREDIENTS

NON-DROWSY

Tussin CF Max

Multi-Symptom Cough, Cold & Flu

PAIN RELIEVER/FEVER REDUCER-ACETAMINOPHEN

COUGH SUPPRESSANT-DEXTROMETHORPHAN HBr

EXPECTORANT-GUAIFENESIN

NASAL DECONGESTANT-PHENYLEPHRINE HCl

MAXIMUM STRENGTH

SEVERE

RELIEVES:

- Cough, Sore Throat
- Body Aches, Fever
- Nasal Congestion
- Chest Congestion

Adult

For Ages 12 & Over

4 FL OZ (118 mL)

NDC 36800-707-26



COMPARE TO ROBITUSSIN®  
MAXIMUM STRENGTH SEVERE MULTI-SYMPTOM  
COUGH COLD + FLU ACTIVE INGREDIENTS\*

**NON-DROWSY**

# Tussin CF Max

Multi-Symptom  
Cough, Cold & Flu

PAIN RELIEVER/FEVER REDUCER-**ACETAMINOPHEN**  
COUGH SUPPRESSANT-**DEXTROMETHORPHAN HBr**  
EXPECTORANT-**GUAIFENESIN**  
NASAL DECONGESTANT-**PHENYLEPHRINE HCl**

**MAXIMUM STRENGTH**

**SEVERE**


**RELIEVES:**

- Cough,
- Sore Throat
- Body Aches, Fever
- Nasal Congestion
- Chest Congestion

**Adult**  
For Ages 12 & Over



4 FL OZ (118 mL)



### Drug Facts

#### Active ingredients (in each 20 mL) Purposes

Acetaminophen, USP 650 mg.....Pain reliever/fever reducer  
 Dextromethorphan HBr, USP 20 mg.....Cough suppressant  
 Guaifenesin, USP 400 mg.....Expectorant  
 Phenylephrine HCl, USP 10 mg.....Nasal decongestant

#### Uses

- temporarily relieves these symptoms occurring with a cold or flu:
  - cough due to minor throat and bronchial irritation
  - nasal congestion
  - sinus congestion and pressure
  - minor aches and pains
  - headache
  - temporarily reduces fever
  - helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive
- sore throat
- diabetes

#### Warnings

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take
- more than 4,000 mg of acetaminophen in 24 hours
  - with other drugs containing acetaminophen
  - 3 or more alcoholic drinks every day while using this product
- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
  - blisters
  - rash
- If a skin reaction occurs, stop use and seek medical help right away.
- Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
- Do not use**
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.



### Drug Facts (continued)

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

#### Ask a doctor before use if you have

- liver disease
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

#### Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

#### When using this product do not use more than directed

- Stop use and ask a doctor if**
- you get nervous, dizzy or sleepless
  - pain, cough, or nasal congestion gets worse or lasts more than 7 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

#### If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.**  
**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

- do not take more than 5 doses in any 24-hour period
- do not exceed recommended dosage. Taking more than the recommended dose (overdose) may cause serious liver damage.

### Drug Facts (continued)

- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 mL every 4 hours
children under 12 years	do not use

#### Other information

- each 20 mL contains: sodium 15 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

#### Inactive ingredients

anhydrous citric acid, benzyl alcohol, edetate disodium, FD&C red #40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, triacetin, xanthan gum

#### Questions or comments?

1-888-423-0139  
 DISTRIBUTED BY TOPCO ASSOCIATES LLC  
 ELK GROVE VILLAGE, IL 60007  
 ©TOPCO PERA0521 QUESTIONS? 1-888-423-0139  
 topcare@topco.com www.topcarebrand.com



Scan here for more information or call 1-888-423-0139



This TopCare® product is laboratory tested to guarantee its highest quality. Your total satisfaction is guaranteed.



\*This product is not manufactured or distributed by Pfizer, distributor of Robitussin® Maximum Strength Severe Multi-Symptom Cough Cold + Flu.

**DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING**

**PARENTS:**  
 Learn about teen medicine abuse  
 www.StopMedicineAbuse.org



7C126 88 C3

Code Above

## TOPCO TUSSIN CF MAX

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:36800-707
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>SORBITOL</b> (UNII: 506T60A25R)	

**Product Characteristics**

<b>Color</b>	RED	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BERRY	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:36800-707-26	1 in 1 CARTON	09/11/2017	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:36800-707-34	1 in 1 CARTON	09/14/2017	
2		237 mL in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph final	part341	09/11/2017	

**Labeler** - Topco Associates LLC (006935977)

Revised: 8/2022

Topco Associates LLC