

TRIPROLIDINE HYDROCHLORIDE- triprolidine hydrochloride liquid
Rugby Laboratories

TRIPROLIDINE HYDROCHLORIDE- triprolidine hydrochloride

Rugby Laboratories Triprolidine HCl 0.938mg/1mL Bubblegum Liquid

NDC: 0536-1257-75

Drug Facts

Active ingredient

Purpose

(in each 1 mL dropperful)

Tripolidine HCl 0.938 mg

Purpose

Antihistamine

Uses

temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies:

■ runny nose ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes

Warnings

Do not exceed recommended dosage.

Ask a doctor before use if you have

- glaucoma
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland

Ask a doctor before use if you are

taking sedatives or tranquilizers

When using this product

- may cause drowsiness

- excitability may occur, especially in children
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- new symptoms occur

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

do not exceed recommended dosage.

- use only with enclosed dropper
- do not use enclosed dropper for any other drug product

Age	Dose
adult and children 12 year of age or older:	2.67 mL (2.5 milligrams) every 4 to 6 hours, not to exceed 10.67 mL (10 milligrams) in 24 hours, or as directed by a doctor.
children 6 to under 12 years of age:	1.33 mL (1.25 milligrams) every 4 to 6 hours, not to exceed 5.33 mL (5 milligrams) in 24 hours, or as directed by a doctor.
children under 6 years of age:	consult a doctor

Other information

- this packaging is child-resistant.
- store at room temperature 15°-30°C (59°-86°F)
- Tamper evident by foil seal under cap. Do not use if foil seal is missing or broken

Inactive ingredients

bubble gum flavor, citric acid, glycerin, methylparaben, monoammonium glycyrrhizinate, potassium citrate, potassium sorbate, propylene glycol, propylparaben, purified water, sucralose

Questions or comments?

Call 1-800-645-2158.

PRINCIPAL DISPLAY PANEL

NDC 0536-1257-75

Rugby
Triprolidine HCl
Antihistamine
0.938mg/1mL
Bubblegum Liquid

<p>Drug Facts</p> <p>Active ingredient (in each 1 mL dropperful) Triprolidine HCl 0.938 mg Antihistamine</p> <p>Purpose</p> <p>Uses temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies:</p> <ul style="list-style-type: none"> runny nose sneezing itching of the nose or throat itchy, watery eyes <p>Warnings Do not exceed recommended dosage.</p>	<p>Drug Facts (continued)</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> glaucoma a breathing problem such as emphysema or chronic bronchitis trouble urinating due to an enlarged prostate gland <p>Ask a doctor before use if you are taking sedatives or tranquilizers</p> <p>When using this product</p> <ul style="list-style-type: none"> may cause drowsiness excitability may occur, especially in children alcohol, sedatives, and tranquilizers may increase the drowsiness effect avoid alcoholic beverages use caution when driving a motor vehicle or operating machinery 	<p>Drug Facts (continued)</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> new symptoms occur <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)</p> <p>Directions do not exceed recommended dosage. use only with enclosed dropper do not use enclosed dropper for any other drug product</p>								
<p>Drug Facts (continued)</p> <table border="1"> <thead> <tr> <th>Age</th> <th>Dose</th> </tr> </thead> <tbody> <tr> <td>adults and children 12 years of age or older:</td> <td>2.67 mL (2.5 milligrams) every 4 to 6 hours, not to exceed 10.67 mL (10 milligrams) in 24 hours, or as directed by a doctor.</td> </tr> <tr> <td>children 6 to under 12 years of age:</td> <td>1.33 mL (1.25 milligrams) every 4 to 6 hours, not to exceed 5.33 mL (5 milligrams) in 24 hours, or as directed by a doctor.</td> </tr> <tr> <td>children under 6 years of age:</td> <td>consult a doctor</td> </tr> </tbody> </table>	Age	Dose	adults and children 12 years of age or older:	2.67 mL (2.5 milligrams) every 4 to 6 hours, not to exceed 10.67 mL (10 milligrams) in 24 hours, or as directed by a doctor.	children 6 to under 12 years of age:	1.33 mL (1.25 milligrams) every 4 to 6 hours, not to exceed 5.33 mL (5 milligrams) in 24 hours, or as directed by a doctor.	children under 6 years of age:	consult a doctor	<p>Drug Facts (continued)</p> <p>Other information</p> <ul style="list-style-type: none"> this packaging is child-resistant. store at room temperature 15°-30°C (59°-86°F) Tamper evident by foil seal under cap. Do not use if foil seal is missing or broken <p>Inactive ingredients bubble gum flavor, citric acid, glycerin, methylparaben, monoammonium glycyrrhizinate, potassium citrate, potassium sorbate, propylene glycol, propylparaben, purified water, sucralose</p> <p>Questions or comments? Call 1-800-645-2158.</p>	<p>Do not use if foil seal under cap is missing or broken.</p> <p>Usual Dosage: See attached labeling for complete product information. Store at 15°-30°C (59°-86°F) KEEP OUT OF REACH OF CHILDREN</p> <p>Dist. by: RUGBY® LABORATORIES 17177 N Laurel Park Dr, Suite 233 Livonia, MI 48152 www.rugbylaboratories.com</p> <p>Rugby® NDC 0536-1257-75</p> <p>Triprolidine HCl</p> <p>Each dropperful (1 mL) contains: Triprolidine HCl 0.938 mg</p> <p>Antihistamine</p> <p>0.938 mg/1 mL</p> <p>Sugar Free, Dye Free Alcohol Free Bubble Gum Flavored Liquid</p> <p>1 FL. OZ (30 mL)</p> <p>Rev. 10/15 IP-166 No-order No. 310605</p>
Age	Dose									
adults and children 12 years of age or older:	2.67 mL (2.5 milligrams) every 4 to 6 hours, not to exceed 10.67 mL (10 milligrams) in 24 hours, or as directed by a doctor.									
children 6 to under 12 years of age:	1.33 mL (1.25 milligrams) every 4 to 6 hours, not to exceed 5.33 mL (5 milligrams) in 24 hours, or as directed by a doctor.									
children under 6 years of age:	consult a doctor									

TRIPROLIDINE HYDROCHLORIDE

triprolidine hydrochloride liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0536-1257
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TRIPROLIDINE HYDROCHLORIDE (UNII: YAN7R5L890) (TRIPROLIDINE - UNII:2L8T9S52QM)	TRIPROLIDINE HYDROCHLORIDE	0.938 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
GLYCERIN (UNII: PDC6A3C0OX)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
AMMONIUM GLYCYRRHIZATE (UNII: 3VRD35U26C)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	BUBBLE GUM	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0536-1257-75	30 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product	11/09/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/09/2019	10/31/2024

Labeler - Rugby Laboratories (079246066)

Establishment

Name	Address	ID/FEI	Business Operations
MONARCH PCM, LLC		080000294	manufacture(0536-1257)

Revised: 1/2024

Rugby Laboratories