

NICOTINE- nicotine polacrilex lozenge
AmerisourceBergen (Good Neighbor Pharmacy) 46122

Drug Facts

Active ingredient (in each lozenge)

Nicotine polacrilex USP, 4 mg

Purpose

Stop smoking aid

Use

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding,

only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth problems occur
- persistent indigestion or severe sore throat occurs
- irregular heartbeat or palpitations occur

- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away

Directions

- **if you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.**
- before using this product, read the enclosed User’s Guide for complete directions and other important information
- begin using the lozenge on your quit day
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine lozenge
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours

- **nicotine lozenge is a medicine and must be used a certain way to get the best results**
- place the lozenge in your mouth and allow the lozenge to slowly dissolve (about 20 to 30 minutes). Minimize swallowing. **Do not chew or swallow lozenge.**
- you may feel a warm or tingling sensation
- occasionally move the lozenge from one side of your mouth to the other until completely dissolved (about 20 to 30 minutes)
- do not eat or drink 15 minutes before using or while the lozenge is in your mouth
- to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
- do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
- **do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.**
- It is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

Other information

- **each lozenge contains:** sodium 14 mg
- store at 20° to 25°C (68° to 77°F)
- keep bottle tightly closed and protect from light

Inactive ingredients

acesulfame potassium, colloidal silicon dioxide, flavor, hydroxypropyl cellulose, magnesium stearate, mannitol, potassium bicarbonate, povidone, sodium alginate, sodium carbonate, sucralose, xanthan gum

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to Nicorette® Lozenge active ingredient*

Nicotine Lozenges

nicotine polacrilex lozenge, 4 mg (nicotine)

Stop Smoking Aid

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.

If you smoke your first cigarette **MORE THAN 30 MINUTES** after waking up, use 2 mg Nicotine Polacrilex Lozenge.

LOZENGES

TO INCREASE YOUR SUCCESS IN QUITTING:

- 1.** You must be motivated to quit.
- 2.** Use Enough - Use **at least 9** Nicotine Polacrilex lozenges per day during the first six weeks.
- 3.** Use Long Enough - Use Nicotine Polacrilex lozenges for the full 12 weeks.
- 4. Use With a Support Program** as directed in the enclosed User's Guide.
 - **not for sale to those under 18 years of age**
 - **proof of age required**
 - **not for sale in vending machines or from any source where proof of age cannot be verified**

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Lozenge

TAMPER EVIDENT FEATURE: DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING.

RETAIN OUTER CARTON FOR FULL PRODUCT USES, DIRECTIONS AND WARNINGS.

Distributed by:

AmerisourceBergen

1 West First Avenue,
 Conshohocken, PA 19428
 www.mygnp.com

Package Label

<p>Drug Facts</p> <p>Active ingredient (in each lozenge) Nicotine polacrilex USP, 4 mg. Purpose Stop smoking aid.</p> <p>Use</p> <ul style="list-style-type: none"> ● reduce withdrawal symptoms, including nicotine craving, associated with quitting smoking <p>Warnings</p> <p>If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ● a sodium-restricted diet ● heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate. ● high blood pressure not controlled with medication. Nicotine can increase your blood pressure. ● stomach ulcer or diabetes. ● history of seizures <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> ● using a non-nicotine stop smoking drug ● taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted. <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ● mouth problems occur ● persistent hoarseness or severe sore throat occurs ● irregular heartbeat or palpitations occur ● you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat ● you have symptoms of an allergic reaction (such as difficulty breathing or rash) <p>Keep out of reach of children and pets. Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wash it off paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.</p> <p>Directions</p> <ul style="list-style-type: none"> ● If you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you. ● before using this product, read the enclosed User's Guide for complete directions and other important information ● begin using the lozenge on your quit day ● If you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine lozenge ● If you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine lozenge according to the following 12 week schedule: <table border="1"> <thead> <tr> <th>Weeks 1 to 6</th> <th>Weeks 7 to 9</th> <th>Weeks 10 to 12</th> </tr> </thead> <tbody> <tr> <td>1 lozenge every 1 to 2 hours</td> <td>1 lozenge every 2 to 4 hours</td> <td>1 lozenge every 4 to 8 hours</td> </tr> </tbody> </table> <ul style="list-style-type: none"> ● nicotine lozenge is a medicine and must be used a certain way to get the best results ● place the lozenge in your mouth and allow the lozenge to slowly dissolve (about 20 to 30 minutes). Minimize swallowing. Do not chew or swallow lozenge. ● you may feel a warm or tingling sensation 	Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12	1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ● occasionally move the lozenge from one side of your mouth to the other until completely dissolved (about 20 to 30 minutes) ● do not eat or drink 15 minutes before using or while the lozenge is in your mouth ● to increase your chances of quitting, use at least 9 lozenges per day for the first 6 weeks ● do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you dizziness, heartburn, nausea or other side effects ● do not use more than 5 lozenges in 8 hours. Do not use more than 20 lozenges per day. ● it is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider. <p>Other information</p> <ul style="list-style-type: none"> ● each lozenge contains sodium 14 mg ● store at 20° to 25° (68° to 77°) ● keep bottle tightly closed and protect from light <p>Inactive Ingredients acesulfame potassium, colloidal silicon dioxide, flavors, hydroxypropyl cellulose, magnesium stearate, mannitol, potassium bicarbonate, povidone, sodium alginate, sodium carbonate, sucralose, xanthan gum</p> <p>Questions or comments? Call 1-877-733-8899 Monday-Friday 9AM-5PM EST</p> <p><small>*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Lozenge.</small></p> <p>TAMPER EVIDENT FEATURE: DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING.</p> <p>RETAIN OUTER CARTON FOR FULL PRODUCT USE, DIRECTIONS AND WARNINGS.</p> <p>not for sale to those under 18 years of age ● proof of age required ● not for sale in vending machines or from any source where proof of age cannot be verified</p>	<p>GOOD NEIGHBOR PHARMACY®</p> <p>Compare to Nicorette® Lozenge active ingredient*</p> <p>NDC 46122-732-00 Includes User's Guide</p> <p>4 mg</p> <p>Nicotine Lozenge nicotine polacrilex lozenge, 4 mg (nicotine)</p> <p>Stop Smoking Aid</p> <p>FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP. If you smoke your first cigarette MORE THAN 30 MINUTES after waking up, use 2 mg Nicotine Polacrilex Lozenge</p> <p>72 Lozenges, 4 mg Each (3 Bottles of 24)</p> <p>Mint Flavor</p> <p>actual 8009</p>	<p>TO INCREASE YOUR SUCCESS IN QUITTING:</p> <ol style="list-style-type: none"> 1. You must be motivated to quit. 2. Use Enough - Use at least 9 Nicotine Polacrilex Lozenges per day during the first six weeks. 3. Use Long Enough - Use Nicotine Polacrilex Lozenges for the full 12 weeks. 4. Use With a Support Program as directed in the enclosed User's Guide. <p>Distributed by: AmeriSourceBergen 1 West First Avenue Conshohocken, PA 19428 Questions or Concerns? Visit us at www.mygnp.com</p> <p>100% SATISFACTION GUARANTEED</p>
Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12							
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours							



Product of Switzerland
 P.L.D. A035A FZ00295
 Lot No:
 Exp. Date:

GOOD NEIGHBOR PHARMACY Nicotine Lozenge Mint Flavor

NICOTINE nicotine polacrilex lozenge

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:46122-732
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	4 mg

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)	

SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MANNITOL (UNII: 3OWL53L36A)	
POTASSIUM BICARBONATE (UNII: HM5Z15LEBN)	
POVIDONE (UNII: FZ989GH94E)	
SODIUM ALGINATE (UNII: C269C4G2ZQ)	
SODIUM CARBONATE (UNII: 45P3261C7T)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	white (off white)	Score	no score
Shape	ROUND	Size	16mm
Flavor	MINT	Imprint Code	A754
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46122-732-08	3 in 1 BOX	02/24/2023	
1		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:46122-732-62	1 in 1 BOX	02/24/2023	
2		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209206	02/24/2023	

Labeler - AmerisourceBergen (Good Neighbor Pharmacy) 46122 (007914906)