

**NIGHTTIME COLD AND FLU MAXIMUM STRENGTH- acetaminophen,
diphenhydramine hci, phenylephrine hci liquid
TARGET Corporation**

Drug Facts

Active ingredients (in each 20 mL)

Acetaminophen 650 mg

Diphenhydramine HCl 25 mg

Phenylephrine HCl 10 mg

Purpose

Pain reliever/fever reducer

Antihistamine / cough suppressant

Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms
 - cough
 - nasal congestion
 - minor ache and pains
 - sore throat
 - headache
 - runny nose
 - sneezing
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other drug containing diphenhydramine even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- you are taking the blood thinning drug warfarin
- you are taking sedatives or tranquilizers

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with fever, rash, or headache that lasts These could be

signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses in any 24 hours period
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years and older :
 - 20 mL every 4 hours while symptoms last
- children under 12 years of age do not use

Other information

- **each 20 mL contains:** sodium 12 mg
- store between 20-25°C (68-77F). Do not refrigerate

Inactive ingredients

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call 1-800-910-6874

Principal Display Panel

Compare to active ingredients in Maximum Strength Mucinex® Fast-Max® NightTime Cold & Flu*

maximum strength

Nighttime

cold and Flu

acetaminophen

(pain reliever / fever reducer)

diphenhydramine HCl

(antihistamine / cough suppressant)

phenylephrine HCl

(nasal decongestant)

relieves aches, fever and sore throat

controls cough

relieves nasal congestion

relieves runny nose & sneezing

AGES 12 + YEARS

FL OZ (mL)

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex®Fast-Max® Night Time Cold & Flu.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

Dist. by Target Corp.

Mpls., MN 55403

Product Label

NDC 11673-460-06

Compare to active ingredients in Maximum Strength Mucinex® Fast-Max® Night Time Cold & Flu*

maximum strength nighttime cold and flu

acetaminophen

(pain reliever/fever reducer)
diphenhydramine HCl
(antihistamine/cough suppressant)
phenylephrine HCl (nasal decongestant)

relieves aches, fever and sore throat
controls cough
relieves nasal congestion
relieves runny nose and sneezing



AGES
12+
YEARS

6 FL OZ (177 mL)

PLD- B377D LB004229

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by Peckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Night Time Cold & Flu.

094 14 0185
C-000274-01-049
Dist. by Target Corp.
Mpls., MN 55403
TM & ©2022 Target Brands, Inc.

PLD-B377D
LB004230

Drug Facts

Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg	Pain reliever/fever reducer
Diphenhydramine HCl 25 mg	Antihistamine/cough suppressant
Phenylephrine HCl 10 mg	Nasal decongestant

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

Uses ■ temporarily relieves these common cold and flu symptoms
■ cough ■ nasal congestion
■ minor aches and pains ■ sore throat
■ headache ■ runny nose ■ sneezing
■ temporarily reduces fever
■ controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
■ with any other drug containing acetaminophen

Drug Facts (continued)

(prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ with any other drug containing diphenhydramine, even one used on the skin
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ for children under 12 years of age

Ask a doctor before use if you have
■ liver disease ■ heart disease
■ high blood pressure ■ thyroid disease
■ diabetes ■ glaucoma
■ trouble urinating due to an enlarged prostate gland
■ a breathing problem such as emphysema or chronic bronchitis
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if
■ you are taking the blood thinning drug warfarin
■ you are taking sedatives or tranquilizers

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

When using this product
■ do not use more than directed
■ excitability may occur, especially in children
■ marked drowsiness may occur
■ avoid alcoholic drinks
■ alcohol, sedatives, and tranquilizers may increase drowsiness
■ be careful when driving a motor vehicle or operating machinery
Stop use and ask a doctor if
■ nervousness, dizziness, or sleeplessness occur
■ pain, nasal congestion, or cough gets worse, or lasts more than 7 days
■ fever gets worse, or lasts more than 3 days
■ redness or swelling is present
■ new symptoms occur
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

■ do not take more than directed (see Overdose warning)
■ do not take more than 6 doses in any 24-hour period
■ measure only with dosing cup provided. Do not use any other dosing device.
■ mL = milliliter ■ keep dosing cup with product
■ dose as follows or as directed by a doctor
■ adults and children 12 years and older: 20 mL every 4 hours while symptoms last
■ children under 12 years of age: do not use

Other information

■ each 20 mL contains: sodium 12 mg
■ store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call 1-800-910-6874

TARGET Maximum Strength Nigh Time Cold and Flu

NIGHTTIME COLD AND FLU MAXIMUM STRENGTH

acetaminophen, diphenhydramine hci, phenylephrine hci liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-460	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	650 mg in 20 mL	
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)		DIPHENHYDRAMINE HYDROCHLORIDE	25 mg in 20 mL	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)		PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL	
Inactive Ingredients				
Ingredient Name			Strength	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
GLYCERIN (UNII: PDC6A3C0OX)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
SORBITOL (UNII: 506T60A25R)				
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
PROPYL GALLATE (UNII: 8D4SNN7V92)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
SODIUM CITRATE (UNII: 1Q73Q2JULR)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
XANTHAN GUM (UNII: TTV12P4NEE)				
EDETATE CALCIUM DISODIUM (UNII: 25IH6R4SGF)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-460-06	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/31/2016	08/31/2025
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M012	08/31/2016	08/31/2025	

Labeler - TARGET Corporation (006961700)