MUCOSA DM- dextromethorphan hydrobromide, guaifenesin tablet Magno-Humphries, Inc.

Mucosa DM

MHL NDC 54257-277-01 immediate Acting

Mucosa DM

EXPECTORANT/COUGH SUPPRESSANT

Guaifenesin 400 mg

Dextromethorphan HBr 20 mg

50 TABLETS

- Soothes Cough
- Loosens Congestion

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGE

Active ingredients (in each tablet)

Dextromethorphan HBr 20mg Guaifenesin 400 mg

Purpose

Dextromethorphan HBr	20mgCough	h Suppressant Guaifenesin 4	00
mg	Expectorant		

Uses

helps loosen phlegm (mucus), helps thin bronchial secretions to make coughs more productive, temporarily relieves cough due to minor throat and bronchial irritation associated with the common cold.

Warnings

a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for two weeks after stopping the MAOI drug. If you do not know if your prescription drug contains MAOI, ask a doctor or pharmacist before taking this product. **Do not use for children under 12. Do not use if you are now taking**

cougha ccompanied with excessive phlegm (mucus), persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema. **Ask a doctor before use if you have**

do not use for more than 7 days. When using this product do not exceed recommended dosage

cough lasts for more than 7 days, recurs, or is accompanied with fever, rash, or persistent headache. These could be signs of a serious condition. **Stop use and ask a doctor if**

ask a health professional before use. If pregnant or breast-feeding,

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

adults and children 12 years & older - take one tablet every 4 hours with a full glass of water while symptoms persist. Do not exceed 6 doses in 24 hours.

children under 12 years of age - do not use

Other information

• store at controlled room temperature 15° to 30°C (59° to 86°F) Keep in a dry place and do not expose to excessive heat.

Inactive ingredients

Croscarmellose sodium, maltodextrin, microcrystalline cellulose, povidone, silicon dioxxide, stearic acid.

Questions?

call toll-free 1-800-935-6737

Distributed by: Magno-Humphries, Inc., OR 97223 U.S.A. Visit our website: www.magno-humphries.com

Tigard, OR 97223 U.S.A. Distributed by: Magno-Humphries, Inc.

Visit our website:

www.magno-humphries.com

Package Labeling:





Drug Facts (cont'd) Directions children under 12 years of age | do not use Other information • store at controlled room temperature 15° to 30°C (59° to 86°F) Keep in a dry place and do not expose to excessive heat. Inactive ingredients Croscarmellose sodium, maltodextrin, microcrystalline cellulose, povidone, silicon dioxide, stearic acid. Questions? call toll-free 1-800-935-6737 Distributed by: Magno-Humphries Labs Tigard, OR 97223 U.S.A. Visit our website: www.magno-humphries.com

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED Drug Facts Active ingredients (in each tablet) Purpose Dextromethorphan HBr 20 mg.......Cough Suppressant Guaifenesin 400 mg **USES** helps loosen phlegm (mucus), helps thin bronchial secretions to make coughs more productive, temporarily relieves cough due to minor throat and bronchial irritation associated with the common cold. Warnings Do not use for children under 12. Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for two weeks after stopping the MAOI drug. If you do not know if your prescription drug contains MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have cough accompanied with excessive phlegm (mucus), persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema. When using this product do not exceed recommended dosage do not use for more than 7 days. Stop use and ask a doctor if cough lasts for more than 7 days, recurs, or is accompanied with fever, rash, or persistent headache. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Directions adults and take one tablet every 4 hours with a children 12 full glass of water while symptoms years & older persist. Do not exceed 6 doses in 24 hours.

MUCOSA DM

dextromethorphan hydrobromide, quaifenesin tablet

PEEL FOR DIRECTIONS G7752-050-01-1

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:54257-277

Route of Administration ORAL

Active Ingredient/Active Moiety Ingredient Name Basis of Strength DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ) GUAIFENESIN (400 mg

Inactive Ingredients		
Ingredient Name	Strength	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)		
MALTODEXTRIN (UNII: 7CVR7L4A2D)		
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)		
POVIDONE, UNSPECIFIED (UNII: FZ 989GH94E)		
SILICON DIOXIDE (UNII: ETJ7Z 6XBU4)		
STEARIC ACID (UNII: 4ELV7Z65AP)		

Product Characteristics			
Color	white	Score	2 pieces
Shape	OVAL	Size	17mm
Flavor		Imprint Code	AZ 345
Contains			

ı	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:54257-277- 01	50 in 1 BOTTLE; Type 0: Not a Combination Product	04/04/2017	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/04/2017	

Labeler - Magno-Humphries, Inc. (063251433)

Revised: 10/2023 Magno-Humphries, Inc.