

CHILDRENS TYLENOL- acetaminophen suspension
Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Children's TYLENOL[®]

Drug Facts

Active ingredient (in each 5 mL = 1 teaspoonful)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use.**
- **do not give more than directed (see overdose warning)**
- **shake well before using**
- mL = milliliter; tsp = teaspoonful
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL or tsp) *
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL (1 tsp)
36-47	4-5 years	7.5 mL (1½ tsp)
48-59	6-8 years	10 mL (2 tsp)
60-71	9-10 years	12.5 mL (2½ tsp)
72-95	11 years	15 mL (3 tsp)

* or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- **each 5 mL (1 tsp) contains:** sodium 2 mg
- store between 20-25°C (68-77°F)
- **do not use if carton tape imprinted "TYLENOL" or bottle wrap imprinted "TYLENOL" is broken or missing**

Inactive ingredients

anhydrous citric acid, butylparaben, D&C red no. 33, FD&C blue no. 1, flavors, glycerin, high fructose corn syrup, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified

water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

Questions or comments?

call **1-800-458-1635** (toll-free) or **215-273-8755** (collect)

PRINCIPAL DISPLAY PANEL

NDC 50580-296-50

See New Warning

Children's
TYLENOL[®]

Acetaminophen Oral Suspension
Pain Reliever-Fever Reducer

Pain + Fever

Ages 2-11

Ibuprofen Free

Alcohol Free

Aspirin Free

4 fl oz (120 ml)

160 mg per 5 ml

Grape

Flavor



SNAP THIS
CODE OR
TEXT TYKID
TO 87715

MSG & DATA RATES
MAY APPLY
TEXT HELP FOR HELP.
TEXT STOP FOR STOP.



How can we help?
☎ 1-800-458-1635

NDC 50580-296-50

See New Warning

Children's TYLENOL®

Acetaminophen Oral Suspension
Pain Reliever-Fever Reducer

Pain + Fever
Ages 2-11

Ibuprofen Free
Alcohol Free
Aspirin Free



4 fl oz (120 ml)
160 mg per 5 ml

Grape
Flavor

Children's TYLENOL®

- ✔ **Pediatricians recommend Children's TYLENOL® for pain relief**
- ✔ **Relied on by moms for over 50 years**

DIRECTIONS:

STEP 1.
Shake well
before using



STEP 2.
Remove child
protective
cap




STEP 3.
Squeeze your
child's dose
into the
dosing cup



IMPORTANT:

Please see dosing chart or ask
a doctor for your child's correct dose

USA
995271
905



Drug Facts

Active ingredient
(in each 5 mL = 1 teaspoonful)
Acetaminophen 160 mg.....Pain reliever/fever reducer

Purpose

Uses temporarily:

- reduces fever
- relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount

- with other drugs containing acetaminophen

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- this product does not contain directions or complete warnings for adult use.
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter; tsp = teaspoonful
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL or tsp) *
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL (1 tsp)
36-47	4-5 years	7.5 mL (1½ tsp)
48-59	6-8 years	10 mL (2 tsp)
60-71	9-10 years	12.5 mL (2½ tsp)
72-95	11 years	15 mL (3 tsp)

* or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL (1 tsp) contains: sodium 2 mg
- store between 20-25°C (68-77°F)
- do not use if carton tape imprinted "TYLENOL" or bottle wrap imprinted "TYLENOL" is broken or missing

Inactive ingredients


anhydrous citric acid, butylparaben, D&C red no. 33, FD&C blue no. 1, flavors, glycerin, high fructose corn syrup, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

Questions or comments?

call 1-800-458-1635 (toll-free) or 215-273-8755 (collect)

Distributed by: 30027299/995271
McNeil Consumer Healthcare
Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA
©McNEIL-PPC, INC. 2014
Visit us at www.tylenol.com





CHILDRENS TYLENOL

acetaminophen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50580-296
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Acetaminophen (UNII: 362O9ITL9D) (acetaminophen - UNII:362O9ITL9D)	Acetaminophen	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
anhydrous citric acid (UNII: XF417D3PSL)	
butylparaben (UNII: 3QP1U3FV8)	
D&C red No. 33 (UNII: 9DBA0SBB0L)	
FD&C blue No. 1 (UNII: H3R47K3TBD)	
glycerin (UNII: PDC6A3C0OX)	
high fructose corn syrup (UNII: XY6UN3QB6S)	
microcrystalline cellulose (UNII: OP1R32D61U)	
carboxymethylcellulose sodium, unspecified form (UNII: K679OBS311)	
propylene glycol (UNII: 6DC9Q167V3)	
water (UNII: 059QF0K00R)	
sodium benzoate (UNII: OJ245FE5EU)	
sorbitol (UNII: 506T60A25R)	
sucralose (UNII: 96K6UQ3ZD4)	
xanthan gum (UNII: TTV12P4NEE)	

Product Characteristics

Color	PURPLE	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

		Marketing Start	Marketing End
--	--	-----------------	---------------

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-296-50	1 in 1 CARTON	11/26/2012	
1		120 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package		
2	NDC:50580-296-51	2 in 1 PACKAGE	01/16/2013	
2		1 in 1 CARTON		
2		120 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part343	07/01/2004	

Labeler - Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division (878046358)

Revised: 11/2016

Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division