

TOXIBAN- activated charcoal granule
LLOYD, Inc. of Iowa

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

ToxiBan®

INDICATION:

For use as a decontamination agent via adsorption of organic chemical toxicants in domestic animals.

In all cases of suspected poisonings, consult your veterinarian for complete diagnosis and specific treatment.

DOSAGE AND ADMINISTRATION:

Large Animals - The recommended dosage is 0.75 to 2.0 grams per kg (0.35 to 0.9 grams per lb) body weight. One pound (454 grams) can normally treat an animal weighing 225 to 600 kg (500 to 1,300 lb). Prepare a suspension by mixing one part of ToxiBan Granules with 5 to 7 parts (volume/volume measures) of cold water. Shake or stir vigorously for 10 seconds. Under the direction of a veterinarian, deliver into stomach or rumen, preferably through a stomach tube and funnel or stomach pump. Rinse container and equipment with cold water and administer to the animal before removing the stomach tube.

Small Animals - The recommended dosage is 2 to 4 grams per kg (1 to 2 grams per lb) body weight. Prepare a suspension by mixing one part of ToxiBan Granules with 5 to 7 parts (volume/volume measures) of cold water. Shake or stir vigorously for 10 seconds. Give orally by encouraging the animal to drink the calculated dose. Consult a veterinarian should administration by stomach tube be needed. ToxiBan Suspension or ToxiBan Suspension with Sorbitol may be more convenient for use in small animals.

Adsorptive Power: Per mg

Each mg of ToxiBan Granules can adsorb at least 0.24 mg strychnine sulfate.

1 level teaspoonful 2 grams

1 level tablespoonful 6 grams

Caution: ToxiBan should not be given in conjunction with oral medications as the charcoal may adsorb the therapeutic agent.

KEEP OUT OF REACH OF CHILDREN

LLOYD, Inc.

Shenandoah, IA 51601 USA

List No. 0422

Rev. 0623

PRINCIPAL DISPLAY PANEL - 5 kg Pail Label

NDC 11789-042-20

ToxiBan® Granules
Water Miscible Charcoal-Sorbitol-Kaolin

INDICATION:

For use as a decontamination agent via adsorption of organic chemical toxicants in domestic animals.

In all cases of suspected poisonings, consult your veterinarian for complete diagnosis and specific treatment.

LLOYD®

Net Weight: 5 kg (11.02 lb)

NDC 11789-042-20

ToxiBan® Granules

Water Miscible Charcoal-Sorbitol-Kaolin

INDICATION:
For use as a decontamination agent via adsorption of organic chemical toxicants in domestic animals.

In all cases of suspected poisonings, consult your veterinarian for complete diagnosis and specific treatment.

LLOYD

Net Weight: 5 kg (11.02 lb)

DOSAGE AND ADMINISTRATION:

Large Animals – The recommended dosage is 0.75 to 2.0 grams per kg (0.35 to 0.9 grams per lb) body weight. One pound (454 grams) can normally treat an animal weighing 225 to 600 kg (500 to 1,300 lb). Prepare a suspension by mixing one part of ToxiBan Granules with 5 to 7 parts (volume/volume measures) of cold water. Shake or stir vigorously for 10 seconds. Under the direction of a veterinarian, deliver into stomach or rumen, preferably through a stomach tube and funnel or stomach pump. Rinse container and equipment with cold water and administer to the animal before removing the stomach tube.

Small Animals – The recommended dosage is 2 to 4 grams per kg (1 to 2 grams per lb) body weight. Prepare a suspension by mixing one part of ToxiBan Granules with 5 to 7 parts (volume/volume measures) of cold water. Shake or stir vigorously for 10 seconds. Give orally by encouraging the animal to drink the calculated dose. Consult a veterinarian should administration by stomach tube be needed. ToxiBan Suspension or ToxiBan Suspension with Sorbitol may be more convenient for use in small animals.

Adsorptive Power: _____ Per mg
Each mg of ToxiBan Granules can adsorb at least 0.24 mg strychnine sulfate.

1 level teaspoonful 2 grams
1 level tablespoonful 6 grams

Caution: ToxiBan should not be given in conjunction with oral medications as the charcoal may adsorb the therapeutic agent.

KEEP OUT OF REACH OF CHILDREN

LLOYD, Inc.
Shenandoah, IA 51601 USA

List No. 0422

Rev. 0623



TOXIBAN

activated charcoal granule

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:11789-042
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACTIVATED CHARCOAL (UNII: 2P3VWU3H10) (ACTIVATED CHARCOAL - UNII:2P3VWU3H10)	ACTIVATED CHARCOAL	240 [arb'U] in 1 g

Inactive Ingredients

Ingredient Name	Strength
KAOLIN (UNII: 24H4NWX5CO)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	BLACK (BLACK GRANULATED POWDER)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11789-042-30	453.6 g in 1 BOTTLE		
2	NDC:11789-042-20	5000 g in 1 PAIL		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
UNAPPROVED DRUG OTHER		07/07/2023	

Labeler - LLOYD, Inc. of Iowa (962286535)

Registrant - LLOYD, Inc. of Iowa (007281942)

Establishment

Name	Address	ID/FEI	Business Operations
LLOYD, Inc. of Iowa		962286535	LABEL, API MANUFACTURE, PACK, MANUFACTURE

Establishment

Name	Address	ID/FEI	Business Operations
LLOYD, Inc. of Iowa		007281942	ANALYSIS

Revised: 7/2023

LLOYD, Inc. of Iowa