

**DG HEALTH CHILDRENS PAIN RELIEF- acetaminophen suspension
Dolgenercorp, LLC**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Dolgenercorp, LLC Children's Pain Relief Drug Facts

Active ingredient (in each 5 mL = 1 teaspoonful)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter; tsp = teaspoonful
- find right dose on chart. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL or tsp)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL (1 tsp)
36-47	4-5 years	7.5 mL (1 ½ tsp)
48-59	6-8 years	10 mL (2 tsp)
60-71	9-10 years	12.5 mL (2 ½ tsp)
72-95	11 years	15 mL (3 tsp)

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL (1 tsp) contains: sodium 3 mg
- store at 20-25°C (68-77°F)
- do not use if printed neckband is broken or missing

Inactive ingredients

anhydrous citric acid, butylparaben, calcium sulfate, carrageenan, D&C red #33, FD&C blue #1, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution,

tribasic sodium phosphate

Questions or comments?

1-888-309-9030

Principal Display Panel

For Ages 2 to 11

Children's Pain Relief

Acetaminophen

Oral Suspension

Fever Reducer – Pain Reliever

Compare to active ingredient of Children's Tylenol® Oral Suspension

Alcohol Free

Ibuprofen Free

Aspirin Free

Fast, Effective Relief of Children's Fever and Pain

Grape Flavor

160 mg per 5 mL

4 FL OZ (118 mL)



For Ages 2 to 11

Children's Pain Relief

Acetaminophen
Oral Suspension



For Ages 2 to 11

Children's Pain Relief

Acetaminophen
Oral Suspension

Fever Reducer • Pain Reliever

Compare to active ingredient of
Children's Tylenol® Oral Suspension**

- Alcohol Free
- Ibuprofen Free
- Aspirin Free

Fast, Effective
Relief of
Children's Fever
and Pain



Grape Flavor



160 mg per 5 mL

4 FL OZ (118 mL)



For Ages 2 to 11

Children's Pain Relief

Acetaminophen
Oral Suspension

Fever Reducer • Pain Reliever

Compare to active ingredient of
Children's Tylenol® Oral Suspension**

- Alcohol Free
- Ibuprofen Free
- Aspirin Free

Fast, Effective
Relief of
Children's Fever
and Pain



Grape Flavor



160 mg per 5 mL

4 FL OZ (118 mL)

A0216



LOT NO.

EXP.

: 13026 VT C2





Drug Facts

Active ingredient Purpose
(in each 5 mL = 1 teaspoonful)

Acetaminophen 160 mg...Pain reliever/fever reducer

Uses temporarily:

- reduces fever
- relieves minor aches and pains due to:
 - the common cold ■ flu ■ headache
 - sore throat ■ toothache

Warnings

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes
- more than 5 doses in 24 hours, which is the maximum daily amount
 - with other drugs containing acetaminophen
- Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter; tsp = teaspoonful
- find right dose on chart. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL or tsp)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL (1 tsp)
36-47	4-5 years	7.5 mL (1 ½ tsp)
48-59	6-8 years	10 mL (2 tsp)
60-71	9-10 years	12.5 mL (2 ½ tsp)
72-95	11 years	15 mL (3 tsp)

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL (1 tsp) contains: sodium 3 mg
- store at 20-25°C (68-77°F)
- do not use if printed neckband is broken or missing

Inactive ingredients anhydrous citric acid, butylparaben, calcium sulfate, carrageenan, D&C red #33, FD&C blue #1, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions or comments?

1-888-309-9030

**This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Oral Suspension.

DISTRIBUTED BY DOLGENCORP, LLC
100 MISSION RIDGE
GOODLETTSVILLE, TN 37072 USA

100%
Quality
Guaranteed

1-888-309-9030



DG HEALTH CHILDRENS PAIN RELIEF

acetaminophen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:559 10-302
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
BUTYLPARABEN (UNII: 3QP11U3FV8)	
CALCIUM SULFATE (UNII: WAT0DDB505)	
CARRAGEENAN (UNII: 5C69YCD2YJ)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)	

Product Characteristics

Color	PURPLE (viscous)	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:559 10-302-26	1 in 1 CARTON	09/09/2012	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	09/09/2012	

Labeler - Dolgencorp, LLC (068331990)

Revised: 12/2017

Dolgencorp, LLC