

MILKHOUSE BRAND FORTREX 1% IODINE TEAT DIP- iodine liquid **Stearns Packaging Corporation**

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Fortrex 1% Iodine Teat Dip

TEAT DIPPING is an important part of a MASTITIS CONTROL PLAN:

1. Dip teats immediately after milking.
2. Use good milking and environmental management procedures.
3. Use properly functioning milking equipment.
4. Treat every quarter of every cow at dry off.
5. Identify and treat or manage clinical cases promptly.
6. Cull chronically infected cows.

GREAT FOR FOAMING

TEAT DIPPING IN COLD WEATHER:

When the temperature falls below 10 degrees F, or when wind chill is significant, special precautions should be taken to avoid chapped and frozen teats.

1. Teats should be dry before turning cows out into cold weather.
2. When teats are dipped after milking, allow 30 seconds contact, and blot off any excess teat dip with a single service towel.
3. Warm the teat dip to reduce drying time.
4. Provide windbreaks in outside areas for cows.
5. Monitor fresh cows with swollen udders and teats since they are more susceptible to chapped and frozen teats.

DIRECTIONS FOR USE:

- DO NOT DILUTE.
- Dip teats immediately after milking as a routine part of milking procedures.
- Dip to THOROUGHLY COVER at least the lower one-third of the teat (preferably to the base of the udder), IMMEDIATELY after every milking.

TAKE TIME

OBSERVE

LABEL DIRECTIONS

- Empty and clean dip cups after each milking, or during milking, if they become contaminated.
- Do not pour use dip back into original container.
- KEEP FROM FREEZING
- Keep container closed to prevent contamination.
- Store container in an upright position.

FIRST-AID:

EYES: If contact with eyes occurs, flush with plenty of cool water for 15 minutes. Consult a physician.

INTERNAL: May be harmful if swallowed. If ingested, drink large amounts of water. DO NOT induce vomiting. Get medical attention immediately.

24 HOUR EMERGENCY CONTACT

CHEM-TEL, 1-800-255-3924

MILKHOUSE BRAND

Fortrex

1% Iodine Teat Dip

NPE-Free

An Aid in Reducing the Spread of Organisms Which May Cause Mastitis

ACTIVE INGREDIENTS: Concentrated iodophor, provides 1% titratable iodine, equivalent to 10,000 ppm titratable iodine.

Emollient system: Contains 10% glycerin.

KEEP OUT OF THE REACH OF CHILDREN

See side panel for precautionary and First-Aid Statements.

NET CONTENTS: 1 U.S. FL. GALLON (3.78 LITERS)

STEARNS PACKAGING CORPORATION

4200 Sycamore Ave. (53714), PO Box 3216, Madison, WI 53704
(800) 655-5008

DIRECTIONS FOR USE:

- DO NOT DILUTE.
- Dip teats immediately after milking as a routine part of milking procedures.
- Dip the THERMOCHEK COVER at least the lower one-third of the teat (preferably to the base of the udder), IMMEDIATELY after every milking.
- Empty and clean dip cups after each milking, or during milking, if they become contaminated.
- Do not reuse teats back into original container.
- KEEP FROM FREEZING
- Keep container closed to prevent contamination.
- Store container in an upright position.

TEAT DIPPING IS AN IMPORTANT PART OF A MASTITIS CONTROL PLAN:

1. Dip teats immediately after milking.
2. Use good milking and environmental management procedures.
3. Use properly functioning milking equipment.
4. Treat every quarter of every cow at dry off.
5. Identify and treat or manage clinical cases promptly.
6. Cull chronically infected cows.

TEAT DIPPING IN COLD WEATHER:

- When the temperature falls below 10°F, or when wind chill is significant, special precautions should be taken to avoid chapped and frozen teats.
1. Teats should be dry before turning cows out in cold weather.
 2. When teats are dipped after milking, allow 30 seconds contact, and blot off any excess teat dip with a single service towel.
 3. Warm the teat dip to reduce drying time.
 4. Provide windbreaks in outside areas for cows.
 5. Monitor fresh cows with swollen udders and teats since they are more susceptible to chapped and frozen teats.



GREAT FOR FOAMING!



An Aid in Reducing the Spread of Organisms Which May Cause Mastitis

ACTIVE INGREDIENTS: Concentrated iodophor, provides 1% titratable iodine, equivalent to 10,000 ppm titratable iodine.

EMOLLIENT SYSTEM: Contains 10% glycerin.

KEEP OUT OF THE REACH OF CHILDREN

See side panel for Precautionary Statements.

NET CONTENTS: 1 U.S. FL. GALLON (3.78 LITERS)



STEARNS PACKAGING CORPORATION
4200 Sycamore Ave. (53714), PO Box 3216, Madison, WI 53704
(800) 655-5008

FIRST-AID:

EYES: If contact with eyes occurs, flush with plenty of cool water for 15 minutes. Consult a physician.

INTERNAL: May be harmful if swallowed. If ingested, drink large amounts of water. DO NOT induce vomiting. Get medical attention immediately.

PRIMEROS AUXILIOS:

OJOS: Si ocurre contacto con los ojos, lavar con abundante agua fría durante 15 minutos. Consulte a un médico.

INTERNO: Puede ser nocivo en caso de ingestión. Beba grandes cantidades de agua si se ingiere este producto. NO induzca el vómito. Solicite atención médica inmediatamente.

24 HOUR EMERGENCY CONTACT

24 horas servicio telefónico de repuesta a emergencias:
1-800-255-3924

#1202247

(ST-224)

REGISTER NUMBER



LOT NO.:

EXP. DATE:

INSTRUCCIONES POR EL USO:

- NO le diluir.
- Sumerja los pezones inmediatamente después del ordeño como parte rutinnaria de procedimiento de ordeño.
- Sumerja para cubrir a fondo el menor el tercio inferior de la tetina (preferiblemente a la base de la ubra); inmediatamente después de cada ordeño.
- Vade y limpie las tetinas sumergiendo después de cada ordeño, o durante el ordeño, si contaminan.
- No vuelva a bañar pezones u/teatado volver a la receptacle de ordeño.
- NO F.E.E. QUE SE CONGELAN.
- Mantenga el recipiente bien cerrado para evitar la contaminación.
- Mantenga el recipiente en una posición vertical.

BAÑO DE PEZONES ES UNA PARTE IMPORTANTE DE UN PLAN PARA CONTROL DE LA MASTITIS:

1. Sumerja los pezones inmediatamente después del ordeño.
2. Utilice buenas prácticas de gestión para el ordeño y medio ambiente.
3. Utilice equipo de ordeño que funcione correctamente.
4. Trate a cada cuarto de la ubra de cada vaca a dentro de hora de sacado.
5. Identifique y trate a manjea casos clínicos con prontitud.
6. Entreeque las vacas que están infectadas crónicamente.

BAÑO DE PEZONES EN TIEMPO FRÍO:

- Cuando la temperatura cae por debajo de 10°F (-12°C), o cuando el factor de enfriamiento del viento es importante tome precauciones especiales para evitar los pezones agrietados y congelados.
1. Los pezones deben estar secos antes de zollar las vacas en el frío.
 2. Cuando los pezones se sumergen después del ordeño, espere 30 segundos de contacto, y posteriormente sacelos el exceso de baño de pezones con una toalla desechable.
 3. Caliente el baño de pezones para reducir el tiempo de sacado.
 4. Proporcione protección contra el viento en las áreas fuera para las vacas.
 5. Vigile las vacas frescas con ubras hinchadas y pezones ya que son más susceptibles a los pezones agrietados y congelados.



LABEL 60282-1224

Rev 03/16

MILKHOUSE BRAND FORTREX 1% IODINE TEAT DIP

iodine liquid

Product Information

Product Type

OTC ANIMAL DRUG

Item Code (Source)

NDC:60282-1224

Route of Administration

TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	1 kg in 100 kg

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	10 kg in 100 kg

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:60282-1224-1	3.96 kg in 1 BOTTLE, PLASTIC		
2	NDC:60282-1224-2	19.80 kg in 1 PAIL		
3	NDC:60282-1224-3	59.39 kg in 1 DRUM		
4	NDC:60282-1224-4	217.77 kg in 1 DRUM		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		04/13/2016	

Labeler - Stearns Packaging Corporation (006069256)**Registrant** - Stearns Packaging Corporation (006069256)**Establishment**

Name	Address	ID/FEI	Business Operations
Stearns Packaging Corporation		006069256	manufacture, api manufacture

Revised: 10/2021

Stearns Packaging Corporation