

CLEARLAX- polyethylene glycol 3350 powder, for solution
STRATEGIC SOURCING SERVICES LLC

SUNMARK PEG

Active ingredient (in each dose)

Polyethylene Glycol 3350, 17 grams (cap filled to the indicated "17 GRAMS" line)

Purpose

Osmotic Laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose , get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed unless advised by your doctor
- the bottle top is a measuring cup marked to contain 17 grams of powder when filled

to the indicated "17 GRAMS" line

- adults and children 17 years of age and older:
- use once a day
- fill to indicated "17 GRAMS" line in cap which is marked to indicate the correct dose 17 grams
- stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- do not combine with starch-based thickeners used for difficulty swallowing
- ensure that the powder is fully dissolved before drinking
- do not drink if there are any clumps
- do not use more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if foil seal under cap printed with "SEALED for YOUR PROTECTION" is missing, open or broken.


Inactive ingredients

none

Questions or comments?

call **833-358-6431** Monday to Friday 9:00am to 7:00pm EST

Package label



Sugar-Free
*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Miralax®

Distributed by
McKesson Corp., via SSSL
Memphis, TN 38141
Money Back Guarantee
www.sunmarkbrand.com
© 2003 McKesson Corporation



Lot No.:
Exp. Date

Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool.
Dissolves In Any Beverage

Drug Facts	Purpose	Drug Facts (continued)
Active ingredient (in each dose) Polyethylene Glycol 3350, 17 grams (cap filled to the indicated "17 GRAMS" line) ... Osmotic Laxative		Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (800) 222-1222.
Use ■ relieves occasional constipation (irregularity) ■ generally produces a bowel movement in 1 to 3 days		Directions ■ do not take more than directed unless advised by your doctor ■ the bottle cap is a measuring cup marked to contain 17 grams of powder when filled to the indicated "17 GRAMS" line ■ adults and children 17 years of age and older: ■ use once a day ■ fill to indicated "17 GRAMS" line in cap which is marked to indicate the correct dose of 17 grams ■ stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink ■ do not combine with starch-based thickeners used for difficulty swallowing ■ ensure that the powder is fully dissolved before drinking ■ do not drink if there are any clumps ■ do not use more than 7 days ■ children 16 years of age or under: ask a doctor
Warnings Allergy alert: Do not use if you are allergic to polyethylene glycol Do not use if you have kidney disease, except under the advice and supervision of a doctor Ask a doctor before use if you have ■ nausea, vomiting or abdominal pain ■ a sudden change in bowel habits that lasts over 2 weeks ■ irritable bowel syndrome Ask a doctor or pharmacist before use if you are taking a prescription drug When using this product you may have loose, watery, more frequent stools Stop use and ask a doctor if ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. ■ you get diarrhea ■ you need to use a laxative for longer than 1 week If pregnant or breast-feeding, ask a health professional before use.		Other information ■ store at 20° - 25°C (68° - 77°F) ■ tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken Inactive ingredients none
		Questions or comments? Call 833-358-6431 Monday to Friday 9:00am to 7:00pm EST

7116-0510-Q3-BL Rev 01/2022

CLEARLAX

polyethylene glycol 3350 powder, for solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49348-143
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL 3350 - UNII:G2M7P15E5P)	POLYETHYLENE GLYCOL 3350	17 g in 17 g

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49348-143-70	238 g in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2016	
2	NDC:49348-143-92	510 g in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090812	07/01/2016	

Labeler - STRATEGIC SOURCING SERVICES LLC (116956644)**Registrant** - Geri-Care Pharmaceutical Corp (611196254)