XTRA BLU PART A- lactic acid solution solution AgroChem USA, LLC

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Xtra BLU Part A

XTRA BLU (Part A)
ClO2 Teat Dip Part A

WARNING

KEEP OUT OF REACH OF CHILDREN

NOT FOR HUMAN USE.

AVOID FREEZING.

KEEP OUT OF DIRECT SUNLIGHT.

CAUSES SERIOUS EYE IRRITATION.

FOR FARM AND INDUSTRIAL USE ONLY.

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal

plant

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

NDC: 17307-3020-3

NET CONTENTS: 55 GALLON (208.2L)

LOT:

EXP. DATE:

XTRA BLU is a two-part Chlorine Dioxide teat dip that can be effectively used to reduce the spread of mastitis causing organisms. Recommended for pre and post milking applications.

DIRECTIONS:

IMPORTANT- MUST BE MIXED PRIOR TO USE. Mix PART A & PART B in equal parts. Mix by volume. CREATES CHLORINE DIOXIDE. Reaches minimum Chlorine Dioxide levels within 15 minutes of mixing. Will maintain minimum Chlorine Dioxide levels for up to 24 hours. It is recommended to replace with freshly mixed solution after 24 hours.

General Use

Once mixed according to directions, do not mix product with any other teat dips, or other product.

Use a teat dip cup to apply. Clean dip cup just prior to use.

If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.

If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

This product can also be foamed or sprayed using recommended equipment.

Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels

Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product.

Allow product to stand for 30 seconds.

Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product.

Dip each teat full-length into the teat dip cup containing product. Allow to

air dry. Do not wipe.

If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking.

Dipping should be resumed at least twice daily one week before calving.

Manufactured By:

AgroChem USA, LLC 26 Freedom Way • Saratoga Springs, NY 12866 • (518)226-4850

XTRA BLU (PART A)

CIO₂ TEAT DIP PART A

WARNING

NOT FOR HUMAN USE.
AVOID FREEZING.
KEEP OUT OF DIRECT SUNLIGHT.
CAUSES SERIOUS EYE IRRITATION.
KEEP OUT OF REACH OF CHILDREN.
FOR FARM AND INDUSTRIAL USE ONLY.

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/face protection.

FIRSTAID/ RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

STORAGE: Store locked up.

 $\ensuremath{\mathsf{DISPOSAL}}$: Dispose of contents/ container to an approved waste disposal plant.

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

ACTIVE INGREDIENT: Lactic Acid 2.64% EMOLLIENT PACKAGE: 12 %

NDC: 17307-3020-3

NET CONTENTS: 55 GALLON (208.2L)

LOT: EXP:







XTRA BLU is a two-part Chbrine Dioxide teat dip that can be effectively used to reduce the spread of mastitis causing organisms. Recommended for pre and post milking applications.

DIRECTIONS:

IMPORTANT- MUST BE MIXED PRIOR TO USE. Mix PART A & PART B in equal parts. Mix by volume. CREATES CHLORINE DIOXIDE. Reaches minimum Chlorine Dioxide levels within 15 minutes of mixing. Will maintain minimum Chlorine Dioxide levels for up to 24 hours. It is recommended to replace with freshly mixed solution after 24 hours.

General Use

Once mixed according to directions, do not mix product, any other teat dips, or other products.

Use a teat dip cup to apply. Clean dip cup just prior to use

If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.

If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product.

Do not reuse or return any unused product to the original container.

This product can also be foamed or sprayed using recommended equipment.

Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis.

Pie-wash teats thoroughly just prior to milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels.

Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product. Allow product to stand for 30 seconds.

Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product.

Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking. Dipping should be resumed at least twice daily one week before calving.

Manufactured By: AgroChemUSA, LLC 26 Freedom Way Saratoga Springs, NY 12866 Phone: 518-226-4850



XTRA BLU PART A

lactic acid solution solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:17307-3020
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
LACTIC ACID (UNII: 33X04XA5AT) (LACTIC ACID - UNII:33X04XA5AT)	LACTIC ACID	0.03 kg in 1 kg	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17307-3020-1	19 kg in 1 CONTAINER		
2	NDC:17307-3020-2	56.8 kg in 1 DRUM		
3	NDC:17307-3020-3	208.2 kg in 1 DRUM		
4	NDC:17307-3020-4	1003.1 kg in 1 CONTAINER		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/18/2023	
other			

Labeler - AgroChem USA, LLC (118336116)

Registrant - AgroChem USA, LLC (118336116)

Establishment			
Name	Address	ID/FEI	Business Operations
AgroChem USA, LLC		118336116	api manufacture, manufacture, analysis, label, pack

Revised: 12/2023 AgroChem USA, LLC