

SALICYLIC ACID- medicated corn removers patch
Foodhold USA

CareOne Medicated Corn Removers

Active ingredient

Salicylic acid 40%

Purpose

Corn remover

Uses

- for the removal of corns
- relieves pain by removing corns

Warnings

For external use only.

Do not use

- if you are diabetic
- have poor blood circulation
- on irritated skin, or any area that is infected or reddened

Stop use and ask a doctor if discomfort persists

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- wash affected area and dry area thoroughly
- if necessary, cut medicated patch to fit corn
- apply adhesive side down of medicated patch onto corn
- cover medicated patch with pad after 48 hours, remove medicated patch
- repeat procedure every 48 hours as needed for up to 14 days (until corn is removed)
- may soak corn in warm water for 5 minutes to assist in removal

Other information

store between 15°C to 30°C (59°F to 86°F)

Inactive ingredients

acrylic adhesive, acrylic polymer, polyethylene, polyvinyl alcohol

Questions?

call 1-866-964-0939

Principal Display Panel

CareOne

MEDICATED CORN REMOVERS

Salicylic Acid

EFFECTIVE CORN REMOVAL TREATMENT

CUSHIONS AGAINST PRESSURE & FRICTION

9 PADS

9 MEDICATED PATCHES

CAREone®

NDC 00000-000-00

Compare to the Active Ingredient in
Dr. Scholl's® Corn Removers*

LOT #
EXP DATE:

Drug Facts

Active Ingredient Purpose
Salicylic acid 40%.....Corn remover

Uses

- for the removal of corns
- relieves pain by removing corns

Warnings

For external use only.

Do not use

- if you are diabetic
- if you have poor blood circulation
- on irritated skin, on any area that is infected or reddened

Stop use and ask a doctor if discomfort persists

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- wash affected area and dry thoroughly
- if necessary, cut medicated patch to fit corn
- apply adhesive side of medicated patch onto corn
- cover medicated patch with pad
- after 48 hours, remove medicated patch
- repeat procedure every 48 hours as needed for up to 14 days (until corn is removed)
- may soak corn in warm water for 5 minutes to assist in removal

Other information

store between 15° and 30°C (59° and 86°F)

Inactive ingredients

acrylic adhesive, acrylic polymer, polyethylene, polyvinyl alcohol

Questions? Call 1-866-964-0939



*This product is not manufactured or distributed by Bayer Healthcare LLC, owner of the registered trademark Dr. Scholl's®.
DISTRIBUTED BY FOODHOLD U.S.A., LLC
LANDOVER, MD 20785-1-877-846-9949
© 2018 Ahold Licensing Sarl
MADE IN USA WITH U.S. AND IMPORTED PARTS
Quality guaranteed or your money back.

EFFECTIVE CORN REMOVAL TREATMENT
CUSHIONS AGAINST PRESSURE & FRICTION



9 PADS
9 MEDICATED PATCHES

52-003CO-01

SALICYLIC ACID

medicated corn removers patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41520-558
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SALICYLIC ACID (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ)	SALICYLIC ACID	40 mg in 9

Inactive Ingredients

Ingredient Name	Strength
POLYVINYL ACETATE (UNII: 32K497ZK2U)	
VINYL ACETATE (UNII: L9MK238N77)	
HIGH DENSITY POLYETHYLENE (UNII: UG00KM4WR7)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41520-558-09	9 in 1 PACKAGE; Type 1: Convenience Kit of Co-Package	11/27/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M030	11/27/2017	

Labeler - Foodhold USA (809183973)

Revised: 1/2024

Foodhold USA