

NEOSPORIN PLUS BURN RELIEF- bacitracin zinc, neomycin sulfate, polymyxin b sulfate, and pramoxine hydrochloride ointment
Johnson & Johnson Consumer Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

NEOSPORIN[®] + BURN RELIEF

Drug Facts

Active ingredients (in each gram)	Purpose
Bacitracin Zinc (500 units)	First aid antibiotic
Neomycin Sulfate (3.5 mg)	First aid antibiotic
Polymyxin B Sulfate (10,000 units)	First aid antibiotic
Pramoxine HCl (10 mg)	External analgesic

Uses

first aid to help prevent infection and for the temporary relief of pain in minor:

- cuts
- scrapes
- burns

Warnings

For external use only.

Do not use

- if you are allergic to any of the ingredients
- in the eyes
- over large areas of the body

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

Stop use and ask a doctor if

- you need to use longer than 1 week
- condition persists or gets worse
- symptoms persist for more than 1 week, or clear up and occur again within a few days
- rash or other allergic reaction develops

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Centre right away.

Directions

- adults and children 2 years of age and older:
 - clean the affected area
 - apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
 - may be covered with a sterile bandage
- children under 2 years of age: ask a doctor

Other information

- store at 20° to 25°C (68° to 77°F)

Inactive ingredients

Petrolatum

Questions?

call **800-223-0182** or outside the US **215-273-8755** (collect)

Distributed by:

JOHNSON & JOHNSON CONSUMER INC.

Skillman, NJ 08558

PRINCIPAL DISPLAY PANEL - 14.2 g Tube Carton

#1 **DOCTOR**
RECOMMENDED
BRAND

NEOSPORIN®

+

BURN RELIEF
Maximum Strength
Pain Relief to
Soothe Minor Burns

24-Hour Infection
Protection

Triple Antibiotic
Formula

NO STING

OINTMENT
FIRST AID ANTIBIOTIC/ PAIN RELIEVING OINTMENT

Bacitracin Zinc-Neomycin Sulfate-Polymyxin B Sulfate-
Pramoxine HCl

NET WT. 0.5 OZ (14.2 g)

30047867



3 12547 23836 6

The makers of NEOSPORIN® do not make store brand products.

how2recycle.info



NEOSPORIN®

#1 DOCTOR RECOMMENDED BRAND



BURN RELIEF



Maximum Strength
Pain Relief to
Soothe Minor Burns

24-Hour Infection
Protection

Triple Antibiotic
Formula

NO STING



OINTMENT

FIRST AID ANTIBIOTIC/PAIN RELIEVING OINTMENT
Bacitracin Zinc - Neomycin Sulfate - Polymyxin B Sulfate -
Pramoxine HCl

NET WT. 0.5 OZ (14.2 g)



Every Cut.
Every Time.
Use with
BAND-AID®
Brand
Bandages



Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
Skillman, NJ 08558
© J&JCI 2020 www.neosporin.com

Drug Facts

Active ingredients (in each gram)

Bacitracin Zinc (500 units).....First aid antibiotic
Neomycin Sulfate (3.5 mg).....First aid antibiotic
Polymyxin B Sulfate (10,000 units).....First aid antibiotic
Pramoxine HCl (10 mg).....External analgesic

Uses first aid to help prevent infection and for temporary relief of pain or discomfort in minor:

Drug Facts (continued)

■ cuts ■ scrapes ■ burns

Warnings

For external use only.

Do not use

■ if you are allergic to any of the ingredients
■ in the eyes ■ over large areas of the body

Drug Facts (continued)

Ask a doctor before use if you have

■ deep or puncture wounds ■ animal bites ■ serious burns

Stop use and ask a doctor if

■ you need to use longer than 1 week
■ condition persists or gets worse
■ symptoms persist for more than 1 week, or clear up and occur again within a few days
■ rash or other allergic reaction develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

■ adults and children 2 years of age and older:
■ clean the affected area ■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily ■ may be covered with a sterile bandage
■ children under 2 years of age: ask a doctor

Other information ■ Store at 20° to 25°C (68° to 77°F)

Inactive ingredient Petrolatum

Questions? Call 800-223-0182 or 215-273-8755 (collect)

EXP
LOT

CODE AREA

NEOSPORIN®

+ BURN RELIEF

OINTMENT

NEOSPORIN PLUS BURN RELIEF

bacitracin zinc, neomycin sulfate, polymyxin b sulfate, and pramoxine hydrochloride ointment

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69968-0518
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BACITRACIN ZINC (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN	500 [iU] in 1 g
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN SULFATE	3.5 mg in 1 g
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B	10000 [iU] in 1 g
PRAMOXINE HYDROCHLORIDE (UNII: 88AYB867L5) (PRAMOXINE - UNII:068X84E056)	PRAMOXINE HYDROCHLORIDE	10 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
PETROLATUM (UNII: 4T6H12BN9U)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69968-0518-1	1 in 1 CARTON	02/01/2019	
1		14.2 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333B	02/01/2019	

Labeler - Johnson & Johnson Consumer Inc. (118772437)