MAXIMUM STRENGTH ANTACID MINT- aluminum hydroxide, magnesium hydroxide, dimethicone liquid CVS

CVS ANTACID MINT MAX

Active ingredients (in each 5 mL teaspoonful)

Aluminum hydroxide 400 mg (equivalent to dried gel, USP) Magnesium hydroxide 400 mg Simethicone 40 mg

Purposes

Antacid

Antigas

Uses

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are taking a

prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Directions

- shake well before use
- adults and children 12 years of age and older: take 2 to 4 teaspoonfuls between meals, at bedtime or as directed by a doctor
- do not take more than 12 teaspoonfuls in 24 hours or use the maximum dosage for more than 2 weeks.

• children under under 12 years: ask a doctor

Other information

- each 5 mL teaspoonful contains: magnesium 165 mg, sodium 1 mg
- protect from freezing
- store at room temperature
- keep tightly closed

Inactive ingredients

benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor, hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

package label

CVS HEALTH

compare to active ingredients in Maalox® Maximum Strength*

Maximum Strength

antacid

liquid

Alumina, Magnesia Simethicone


antacid/antigas Fast-acting

Soothing relief of

- heartburn
- acid indigestion
- pressure & bloating

Mint Flavor

12 FL OZ (355 mL)



Drug Facts

TAMPER EVIDENT: DO NOT USE IF BREAKAWAY BAND ON BOTTLE CAP IS MISSING OR BROKEN.

Active ingredients (in each 5 mL teaspoonful)

Aluminum hydroxide (equivalent to dried gel, USP) 400 mg ... Antacid

Magnesium hydroxide 400 mg Simethicone 40 mg

Uses relieves: • heartburn

- sour stomach
- · acid indigestion · symptoms referred to as gas

Ask doctor before use if you have

kidney disease • a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids may interact with certain prescription drugs. Stop use and ask a doctor if symptoms last for more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Directions • shake well before use

- adults and children 12 years of age and older: take 2 to 4 teaspoonfuls between meals, at bedtime or as directed by a doctor
- do not take more than 12 teaspoonfuls in 24 hours or use the maximum dosage for more than 2 weeks.
- children under 12 years of age: ask a doctor

Other information • each 5 mL teaspoonful contains: magnesium 165 mg, sodium 1 mg • store at room temperature • protect from freezing

- keep tightly closed

Inactive ingredients benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor, hypromellose, microcrystalline cellulose, propytparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments? 1-800-540-3765

*This product is not manufactured or distributed by Johnson & Johnson Merck, distributor of Mylantae. #69621

Distributed by: CVS Pharmacy, Inc. One CVS Drive, Woonsocket, RI 02895 C2015 CVS/Pharmacy CVS.com® 1-800-SHOP CVS

V-13966

905-06242-3 REV GC-0615





MAXIMUM STRENGTH ANTACID MINT

aluminum hydroxide, magnesium hydroxide, dimethicone liquid

Product Information

Product Type HUMAN OTC DRUG NDC:69842-296 Item Code (Source)

ORAL Route of Administration

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	400 mg in 5 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	400 mg in 5 mL
DIMETHICONE (UNII: 92RU3N3Y10) (DIMETHICONE - UNII:92RU3N3Y10)	DIMETHICONE	40 mg in 5 mL

Inactive Ingredients				
Ingredient Name	Strength			
HYPROMELLOSES (UNII: 3NXW29V3WO)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
BUTYLPARABEN (UNII: 3QPI1U3FV8)				
PROPYLPARABEN (UNII: Z8IX2SC1OH)				
WATER (UNII: 059QF0KO0R)				
SACCHARIN SODIUM (UNII: SB8ZUX40TY)				
SORBITOL (UNII: 506T60A25R)				
BENZYL ALCOHOL (UNII: LKG8494WBH)				
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K6790BS311)				

Product Characteristics			
Color		Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

ı	Packaging					
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
	1	NDC:69842-296- 73	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2015		

Marketing Information				
Marketing Application Number or Monograph Category Citation		Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M001	06/01/2015		

Labeler - CVS (062312574)

Registrant - GCP Laboratories (965480861)

Establishment					
Name	Address	ID/FEI	Business Operations		
GCP Laboratories		965480861	manufacture(69842-296)		

Revised: 10/2023 CVS