

## **ACID GONE ANTACID- aluminum hydroxide and magnesium carbonate liquid**

### **Major Pharmaceuticals**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **MAJOR ACID GONE**

#### **Active ingredient (in each 15mL tablepoonful)**

Aluminum hydroxide 95mg

Magnesium carbonate 358mg

#### **Purpose**

Antacid

#### **Uses**

relieves

- heartburn
- acid indigestion
- sour stomach
- upset stomach associated with these symptoms

#### **Warnings**

##### **Ask a doctor before use if you have**

- kidney disease
- a magnesium-restricted diet
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are** taking a prescription drug.

Antacids may interact with certain prescription drugs.

##### **When using this product**

- do not take more than 8 tablepoonfuls in 24 hours
- do not use the maximum dosage for more than 2 weeks
- laxative effect may occur

**If pregnant or breast-feeding,** ask a health professional before use.

##### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

- shake well before use
- adults and children 12 years and older: take 1 to 2 tablepoonfuls four times a day or as directed by a doctor
- children under 12 years: ask a doctor

- take after meals and at bedtime
- dispense product only by spoon or other measuring device

**Other information**

- **each 15 mL tablespoonful contains:** magnesium 105mg, sodium 39mg
- store at up to 25°C (77°F)
- avoid freezing
- keep tightly closed
- **TAMPER-EVIDENT:** Do not use this product if inner foil seal over the mouth of the bottle is broken or missing.

**Inactive ingredients (Regular Strength)**

benzyl alcohol, D&C yellow #10, edetate disodium, FD&C blue #1, flavor, glycerin, purified water, saccharin sodium, sodium alginate, sorbitol solution, xanthan gum

**Questions or comments?**

1-800-540-3765

**Principal Display Panel**

major

NDC 0904-7727-14

**Acid Gone Antacid**

**Alumina and Magnesium Carbonate**

**Oral Suspension USP**

Relieves Heartburn  
& Acid Indigestion

Compare to the active ingredients in Gaviscon® \*

12 FL OZ (355 mL)

MAJOR<sup>®</sup>

NDC 0904-7727-14

# Acid Gone<sup>™</sup>

## ANTACID

Alumina and Magnesium Carbonate  
Oral Suspension USP

Relieves Heartburn  
& Acid Indigestion



Compare to the active ingredients in Gaviscon<sup>®</sup>\*

12 FL OZ (355 mL)

226-15212-5  
Rev. 03/17

### Drug Facts

Active ingredients (in each 15 mL tablespoonful)	Purpose
Aluminum hydroxide 95 mg	Antacid
Magnesium carbonate 358 mg	Antacid

**Uses** relieves • acid indigestion • heartburn • sour stomach  
• upset stomach associated with these symptoms

### Warnings

**Ask a doctor before use if you have** • kidney disease  
• a magnesium-restricted diet • a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are** taking a prescription drug.  
Antacids may interact with certain prescription drugs.

### When using this product

- do not take more than 8 tablespoonfuls in 24 hours
- do not use the maximum dosage for more than 2 weeks
- laxative effect may occur

**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help  
or contact a Poison Control Center right away.

### Directions

- shake well before use
- adults and children 12 years and older: take 1 to 2 tablespoonfuls  
four times a day or as directed by a doctor
- children under 12 years: ask a doctor
- take after meals and at bedtime
- dispense product only by spoon or other measuring device

### Other information

- each 15 mL tablespoonful contains: magnesium 105 mg, sodium 39 mg
- store at up to 25°C (77°F) • avoid freezing • keep tightly closed
- **TAMPER-EVIDENT:** Do not use this product if inner foil seal over the  
mouth of the bottle is broken or missing.

### Inactive ingredients

benzyl alcohol, D&C yellow no. 10,  
edetate disodium, FD&C blue no. 1, flavor, glycerin, purified  
water, saccharin sodium, sodium alginate, sorbitol solution,  
xanthan gum

### Questions or comments? 1-800-540-3765

\*This product is not manufactured or distributed by the owner of the registered  
trademark Gaviscon<sup>®</sup>. 926-15212-5

Re-Order No. 264978

Distributed by MAJOR<sup>®</sup> PHARMACEUTICALS  
17177 N Laurel Park Drive, Suite 233, Livonia, MI 48152

Rev 03/17 M-99



## ACID GONE ANTACID

aluminum hydroxide and magnesium carbonate liquid

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0904-7727
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDRO XIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDRO XIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDRO XIDE	95 mg in 15 mL
MAGNESIUM CARBO NATE (UNII: 0E53J927NA) (CARBONATE ION - UNII:7UJQ5OPE7D)	MAGNESIUM CARBONATE	358 mg in 15 mL

## Inactive Ingredients

Ingredient Name	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM ALGINATE (UNII: C269C4G2ZQ)	
SORBITOL (UNII: 506T60A25R)	
WATER (UNII: 059QF0K00R)	
XANTHAN GUM (UNII: TTV12P4NEE)	

## Product Characteristics

Color	green	Score	
Shape		Size	
Flavor	SPEARMINT (cool mint)	Imprint Code	
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-7727-14	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2017	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part331	12/30/2004	

**Labeler** - Major Pharmaceuticals (191427277)

**Registrant** - GCP Laboratories (965480861)

## Establishment

Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(0904-7727)