

MUCINEX FAST-MAX CONGESTION AND HEADACHE- acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride capsule, liquid filled
RB Health (US) LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Mucinex® Fast-Max®
Congestion & Headache

Drug Facts

<i>Active ingredients (in each liquid gel)</i>	<i>Purposes</i>
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - nasal congestion
 - cough
 - headache
 - minor aches and pains
 - sinus congestion and pressure
- temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and older: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredients

FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, lecithin (soy), mineral oil, polyethylene glycol, povidone, propylene glycol, shellac, sorbitol sorbitan solution, titanium dioxide, water

Questions?

1-866-MUCINEX (1-866-682-4639) You may also report side effects to this phone number.

Dist. by: RB Health (US)

Parsippany, NJ 07054-0224
Made in China

PRINCIPAL DISPLAY PANEL - 16 Liquid Gel Blister Pack Carton

Fast Dissolving Liquid Gels!

MAXIMUM STRENGTH
NDC 63824-599-16

Mucinex®
FAST-MAX®

**CONGESTION &
HEADACHE**

Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Phenylephrine HCl – Nasal Decongestant

- ▣ **Relieves Headache & Fever**
- ▣ **Controls Cough**
- ▣ **Relieves Nasal Congestion**

16
LIQUID GELS
(Liquid Filled Capsules)
DAY TIME
FOR AGES 12+

MAXIMUM STRENGTH



CONGESTION & HEADACHE

Tamper evident: Do not use if carton is damaged or if printed seal on blister is broken or missing.

16 LIQUID GELS

Fast Dissolving Liquid Gels!

NDC 63824-599-16

MAXIMUM STRENGTH



CONGESTION & HEADACHE

Also try the Max Strength, Multi-Symptom Mucinex® Fast-Max® Night Time products for night time cold & flu symptom relief

CONGESTION & HEADACHE



MAXIMUM STRENGTH

Maximum Strength per 4-hour dose

Do not take more than a total of 12 liquid gels in a 24-hour period.

Take only as directed.

Keep carton for complete product information.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Acetaminophen – Pain Reliever/Fever Reducer

Dextromethorphan HBr – Cough Suppressant

Phenylephrine HCl – Nasal Decongestant

- ✓ Relieves Headache & Fever
- ✓ Controls Cough
- ✓ Relieves Nasal Congestion



16 LIQUID GELS (Liquid Filled Capsules)

DAY TIME FOR AGES 12+



HEALTH • HYGIENE • HOME

www.mucinex.com

Patents: www.rb.com/patents

Dist. by: RB Health (US)
 Parsippany, NJ 07054-0224

Made in China
 ©2019 RB Health

022519 8357063



3 63824 58716 5

Drug Facts

Active Ingredients (in each liquid gel)


Acetaminophen 325 mg.....Pain reliever/fever reducer

Purposes

Drug Facts (continued)

- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are



MUCINEX FAST-MAX CONGESTION AND HEADACHE

acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63824-599
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Acetaminophen (UNII: 362O9ITL9D) (Acetaminophen - UNII:362O9ITL9D)	Acetaminophen	325 mg
Dextromethorphan hydrobromide (UNII: 9D2RTI9KYH) (Dextromethorphan - UNII:7355X3ROTS)	Dextromethorphan hydrobromide	10 mg
Phenylephrine hydrochloride (UNII: 04JA59TNSJ) (Phenylephrine - UNII:1WS297W6MV)	Phenylephrine hydrochloride	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C red no. 40 (UNII: WZB9127XOA)	
FD&C yellow no. 6 (UNII: H77VEI93A8)	
gelatin, unspecified (UNII: 2G86QN327L)	
glycerin (UNII: PDC6A3C0OX)	
lecithin, soybean (UNII: 1DI56QDM62)	
mineral oil (UNII: T5L8T28FGP)	
polyethylene glycol, unspecified (UNII: 3WJQ0SDW1A)	
povidone, unspecified (UNII: FZ989GH94E)	
propylene glycol (UNII: 6DC9Q167V3)	
shellac (UNII: 46N107B71O)	
sorbitol (UNII: 506T60A25R)	
titanium dioxide (UNII: 15FIX9V2JP)	
water (UNII: 059QF0KO0R)	

Product Characteristics

Color	RED	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	PC9
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63824-599-16	2 in 1 CARTON	11/27/2015	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	11/27/2015	

Labeler - RB Health (US) LLC (081049410)

Revised: 10/2019

RB Health (US) LLC