

**CHILDRENS MULTI SYMPTOM FEVER AND COLD- acetaminophen,  
chlorpheniramine maleate, dextromethorphan hbr, phenylephrine  
hcl suspension  
THE KROGER CO.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Children's Multi-Symptom Fever & Cold 4 FL OZ (118 mL) Grape Flavor**

***Active ingredients (in each 5 mL)***

Acetaminophen 160 mg  
Chlorpheniramine maleate 1 mg  
Dextromethorphan HBr 5 mg  
Phenylephrine HCl 2.5 mg

***Purposes***

Pain reliever/fever reducer  
Antihistamine  
Cough suppressant  
Nasal decongestant

***Uses***

- temporarily relieves
  - minor aches and pains
  - headache
  - minor sore throat pain
  - runny nose
  - itchy nose or throat
  - sneezing
  - itchy, watery eyes due to hay fever
  - nasal and sinus congestion
  - cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily reduces fever

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- in a child under 4 years of age
- if your child is allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for the purpose of making your child sleepy

### **Ask a doctor before use if the child has**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- a breathing problem such as chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- chronic cough that lasts, such as occurs with asthma

### **Ask a doctor or pharmacist before use if the child is**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- **do not exceed recommended dosage**
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness
- excitability may occur, especially in children

## Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Prompt medical attention is critical even if you do not notice any signs or symptoms.

## Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- **shake well before use**
- may be given every 4 hours, while symptoms persist. Do not give more than 5 doses in 24 hours unless directed by a doctor.
- use enclosed dosing cup only. Keep for use with this product. Do not use any other dosing device.
- mL = milliliter

Age	Dose
<b>children under 4 years</b>	<b>do not use</b>
children 4 to under 6 years of age	do not use unless directed by a doctor
children 6 to under 12 years of age	10 mL

## Other information

- each 5 mL contains: **sodium 5 mg**
- **Tamper-evident: do not use if printed inner seal under cap is torn or missing**
- store at room temperature. Protect from light.
- contains no aspirin
- close cap tightly

## Inactive ingredients

citric acid, D&C Red #33, edetate disodium, FD&C Blue #1, FD&C Red #40, flavor, glycerin, microcrystalline cellulose, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium carboxymethylcellulose, sorbitol, sucrose, xanthan gum

**Questions or comments?**

**1-866-467-2748**

**Principal Display Panel**

**NDC 30142-988-04**

**Compare to the active ingredients in Children's Triaminic® Suspension Multi-Symptom Fever & Cold\***

Children's Multi-Symptom

**Fever & Cold**

**ACETAMINOPHEN /**

Pain Reliever/Fever Reliever

**CHLORPHENIRAMINE MALEATE /**

Antihistamine

**DEXTROMETHORPHAN HBR /**

Cough Suppressant

**PHENYLEPHRINE HCL /**

Nasal Decongestant

- **Cold & Flu Symptoms Relief**
- **Fever**
- **Runny & Stuffy Nose**
- **Aches & Pains**
- **Sore Throat**
- **Cough**

**Grape Flavor**

Ages 6-11 Years

Naturally and artificially flavored

4 FL OZ (118 mL)

**IMPORTANT: Keep this carton  
for future reference on full  
labeling**

\*This product is not manufactured or distributed by GSK Consumer Healthcare, owner of the registered trademark Children's Triaminic Suspension Multi-Symptom Fever & Cold.

**Distributed by:**



# CHILDRENS MULTI SYMPTOM FEVER AND COLD

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr, phenylephrine hcl suspension

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:30142-988
<b>Route of Administration</b>	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	1 mg in 5 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 5 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 5 mL

Inactive Ingredients	
Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>D&amp;C RED NO. 33</b> (UNII: 9DBA0SBB0L)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>PROPYL GALLATE</b> (UNII: 8D4SNN7V92)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED</b> (UNII: K679OBS311)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

Product Characteristics			
<b>Color</b>	PURPLE	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	GRAPE	<b>Imprint Code</b>	
<b>Contains</b>			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30142-988-04	1 in 1 CARTON	09/03/2020	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	09/03/2020	

**Labeler** - THE KROGER CO. (006999528)

Revised: 8/2023

THE KROGER CO.