

TIGER SUPPLY INC TOPICAL ANESTHETIC- benzocaine gel
Tiger Supply Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Active Ingredient

Benzocaine 200mg (in each g)

Purpose

Oral Anesthetic

Use

For oral mucosa use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy Alert: Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Do not use

- for more than 7 days unless directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.
- for teething
- in children under 2 years of age

When using this product Avoid contact with eyes. If contact occurs, flush with water.

Do not exceed recommended dosage. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breast feeding, ask a physician before use.

Keep out of reach of children.

Directions

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

Other Information

Store at 59° - 86°F (15°-30°C). Protect from freezing.

Inactive Ingredients

flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

Questions or comments?


631-293-2700 or www.tigersupplyinc.com

PEEL BACK to review drug facts

Manufactured for:



99 Sherwood Avenue
Farmingdale, NY 11735
Tel: 631.293.2700
www.tigersupplyinc.com



TOPICAL ANESTHETIC GEL
BENZOCAINE 20 %

Net Content:
1 oz (30 g) **GLUTEN FREE**

<p>directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.</p> <ul style="list-style-type: none"> • for teething • in children under 2 years of age <p>When using this product Avoid contact with eyes. If contact occurs, flush with water.</p> <p>Do not exceed recommended dosage. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.</p> <p>If pregnant or breast feeding, ask a physician before use.</p> <p>Keep out of reach of children.</p>	<p>Directions</p> <ul style="list-style-type: none"> • Apply only amount needed to the oral mucosa to prevent or relieve pain. • children under 2 years of age: do not use <p>Other information Store at 59° - 86°F (15 - 30°C). Protect from freezing.</p> <p>Inactive ingredients flavoring, PEG 3350, PEG 400, sodium saccharin, water. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.</p> <p>Questions or comments? 631-293-2700 or tigersupplyinc.com</p>	<p>99292251 Rev 11/2018</p>
--	---	-----------------------------

<p>Drug Facts</p> <p>Active ingredients Benzocaine 200 mg (in each g)</p> <p>Purpose Oral Anesthetic</p> <p>Use For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.</p> <p>Warnings Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:</p> <ul style="list-style-type: none"> • pale, gray or blue colored skin (cyanosis) • headache • rapid heart rate • shortness of breath • dizziness or lightheadedness • fatigue or lack of energy <p>Allergy Alert: Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.</p> <p>Do not use</p> <ul style="list-style-type: none"> • for more than 7 days unless 		
---	--	--

TIGER SUPPLY INC TOPICAL ANESTHETIC

benzocaine gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69484-003
Route of Administration	DENTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
FD&C YELLOW NO. 5 (UNII: I753WB2F1M)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
D&C GREEN NO. 5 (UNII: 8J6RDU8L9X)	

Product Characteristics

Color	red	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69484-003-30	30 g in 1 JAR; Type 0: Not a Combination Product	11/01/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	05/08/2013	

Labeler - Tiger Supply Inc. (168720709)

Revised: 1/2022

Tiger Supply Inc.