

**IBUPROFEN- ibuprofen tablet**  
**L.N.K. International, Inc.**

-----  
**Sound Body 44-291**

***Active ingredient (in each brown tablet)***

Ibuprofen USP, 200 mg (NSAID)\*  
\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
  - headache
  - the common cold
  - menstrual cramps
  - backache
  - toothache
  - muscular aches
  - minor pain of arthritis
- temporarily reduces fever

***Warnings***

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- are age 60 or older
- take more of for a longer time than directed
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart

failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you have problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if you are**

- taking any other drug
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - have bloody or black stools
  - vomit blood
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - slurred speech
  - leg swelling
  - trouble breathing
  - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- **do not take more than directed**

- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 tablet, 2 tablets may be used
  - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

#### ***Other information***

- use by expiration date on package
- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)

#### ***Inactive ingredients***

carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

#### ***Questions or comments?***

**Call 1-800-426-9391** 8:30 AM-4:00 PM ET, Monday - Friday

#### **Principal Display Panel**

**SOUNDBODY™**

†**Compare to the active ingredient in Advil® Tablets**

NDC 50844-921-14

#### **Ibuprofen**

Ibuprofen Tablets USP, 200 mg

Pain Reliever/Fever Reducer (**NSAID**)

**500 TABLETS**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS  
BROKEN OR MISSING**

†This product is not manufactured or distributed by Wyeth LLC, owner of the registered trademark Advil® Tablets.

50844 ORG061829114

Manufactured of Big Lots Stores, Inc.

by **LNK INTERNATIONAL, INC.**

60 Arkay Drive, Hauppauge, NY 11788 USA

V#733000 ITEM#022729114



<sup>1</sup>Compare to the active ingredient in Advil® Tablets

NDC 50844-921-14

# Ibuprofen

Ibuprofen Tablets USP, 200 mg  
Pain Reliever/Fever Reducer  
(NSAID)

500 TABLETS

### Drug Facts

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

**Active ingredient (in each brown tablet)**  
Ibuprofen USP, 200 mg (NSAID)<sup>1</sup> .....Pain reliever/fever reducer  
<sup>1</sup>nonsteroidal anti-inflammatory drug

### Purpose

**Uses** ■ temporarily relieves minor aches and pains due to: ■ headache ■ the common cold  
■ menstrual cramps ■ backache ■ toothache  
■ muscular aches ■ minor pain of arthritis  
■ temporarily reduces fever

### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
■ hives ■ facial swelling ■ asthma (wheezing)  
■ shock ■ skin reddening ■ rash ■ blisters  
If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you  
■ take a blood thinning (anticoagulant) or steroid drug  
■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ are age 60 or older  
■ take more or for a longer time than directed  
■ have had stomach ulcers or bleeding problems  
■ have 3 or more alcoholic drinks every day while using this product  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, ■

### Drug Facts (continued)

and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.  
**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer  
■ right before or after heart surgery

### Ask a doctor before use if

■ stomach bleeding warning applies to you  
■ you have a history of stomach problems, such as heartburn ■ you are taking a diuretic  
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke  
■ you have problems or serious side effects from taking pain relievers or fever reducers

### Ask a doctor or pharmacist before use if you are

■ taking any other drug  
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin  
■ under a doctor's care for any serious condition

### When using this product

■ take with food or milk if stomach upset occurs

### Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding: ■ feel faint  
■ have bloody or black stools ■ vomit blood  
■ have stomach pain that does not get better  
■ you have symptoms of heart problems or stroke:  
■ chest pain ■ slurred speech  
■ leg swelling ■ trouble breathing  
■ weakness in one part or side of body  
■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present in the painful area  
■ any new symptoms appear

### Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### Directions

■ do not take more than directed  
■ the smallest effective dose should be used  
■ adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist  
■ if pain or fever does not respond to 1 tablet, 2 tablets may be used  
■ do not exceed 6 tablets in 24 hours, unless directed by a doctor  
■ children under 12 years: ask a doctor

### Other information

■ use by expiration date on package  
■ store between 20°-25°C (68°-77°F)  
■ avoid excessive heat 40°C (104°F)

### Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

**Questions or comments?** Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday



This product is not manufactured or distributed by Wyeth, LLC, owner of the registered trademark Advil® Tablets. 50844-921-14  
Manufactured for: LWK International, Inc.  
60 Arden Drive, Hightstown, NJ 08520 USA

44-291

## IBUPROFEN

ibuprofen tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50844-921
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

### Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

**Product Characteristics**

<b>Color</b>	BROWN	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;291
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-921-12	1 in 1 CARTON	05/24/1988	
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:50844-921-06	200 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/24/1988	05/02/2019
3	NDC:50844-921-14	500 in 1 BOTTLE; Type 0: Not a Combination Product	05/24/1988	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	05/24/1988	

**Labeler** - L.N.K. International, Inc. (038154464)**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(50844-921)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(50844-921)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	PACK(50844-921)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	PACK(50844-921)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	PACK(50844-921)

