

XT- 22C CONCENTRATE (PART B)- sodium chlorite solution solution
AgroChem Inc

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

XT- 22C Concentrate (Part B)

XT- 22C Concentrate (Part B)

DANGER

AVOID FREEZING.

CAUSES EYE IRRITATION.

KEEP OUT OF DIRECT SUNLIGHT.

KEEP OUT OF REACH OF CHILDREN.

FOR FARM AND INDUSTRIAL USE ONLY.

INGREDIENTS:

Sodium Chlorite - 3.5%

XT- 22C Concentrate is a two-part chlorine dioxide teat dip that can be effectively used to reduce the spread of mastitis causing organisms. Recommended for post milking applications.

Directions:

IMPORTANT THIS IS A TWO-PART CONCENTRATE - MUST BE MIXED PRIOR TO USE. MIX RATIO is (20 Water : 1 Part A : 1 Part B). In a clean container, add (20) parts water, then add (1) part XT- 22C Concentrate Part A, next add (1) part XT- 22C Concentrate Part B. FOLLOW THE DIRECTIONS. DO NOT MIX PART A AND B TOGETHER. CREATES CHLORINE DIOXIDE. Reaches minimum Chlorine Dioxide levels (> 1000 ppm) within 15 minutes of mixing.

General Use

Teat Sprays can also be foamed or sprayed using recommended equipment.

If using a teat dip cup to apply. Clean dip cup just prior to use. If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip. If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product. If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

Pre Dipping Use

Perform fore stripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels. Just before each cow is milked, dip, foam, or spray, the full length of each teat with the product.

Allow product to stand for 30 seconds. Wipe teats dry after application, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product. Dip, foam, or spray, the full length of each teat. Allow to air dry. Do not wipe.

If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the product should be applied to the teats, once a day for 3-4 days after the last milking.

Applications should resume at least twice daily one week before calving.

XT-22C CONCENTRATE | PART B

TEAT DIP

UN 1908 Chlorite solution, 8, III

DANGER



AVOID FREEZING.
CAUSES EYE IRRITATION.
KEEP OUT OF DIRECT SUNLIGHT.
KEEP OUT OF REACH OF CHILDREN.
FOR FARM AND INDUSTRIAL USE ONLY.

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.
FIRST AID / RESPONSE:
IF SWALLOWED: Rinse mouth. Do NOT induce vomiting.
IF ON SKIN (or hair): Take off immediately all contaminated clothing. Rinse skin with water/shower. If
INHALED: Remove person to fresh air and keep comfortable for breathing. Immediately call a POISON CENTER/doctor. **IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER/doctor. Wash contaminated clothing before reuse.
STORAGE: Store locked up.
DISPOSAL: Dispose of contents/ container to an approved waste disposal plant

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

INGREDIENTS:
Sodium Chlorite - 3.5%

NET CONTENTS:

LOT: EXP:

XT-22C Concentrate is a two-part chlorine dioxide teat dip that can be effectively used to reduce the spread of mastitis causing organisms. Recommended for post milking applications.

Directions:
IMPORTANT THIS IS A TWO-PART CONCENTRATE - MUST BE MIXED PRIOR TO USE. MIX RATIO is (20 Water : 1 Part A : 1 Part B). In a clean container, add (20) parts water, then add (1) part XT-22C Concentrate Part A, next add (1) part XT-22C Concentrate Part B. FOLLOW THE DIRECTIONS. DO NOT MIX PART A AND B TOGETHER. CREATES CHLORINE DIOXIDE. Reaches minimum Chlorine Dioxide levels (> 1000 ppm) within 15 minutes of mixing.

General Use
Teat Sprays can also be foamed or sprayed using recommended equipment.
If using a teat dip cup to apply. Clean dip cup just prior to use. If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip. If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product. If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

Pre Dipping Use
Perform forestripping and inspect teats for clinical mastitis.
Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels. Just before each cow is milked, dip, foam, or spray, the full length of each teat with the product.
Allow product to stand for 30 seconds. Wipe teats dry after application, using single-service towels to avoid contamination of milk.

Post Dipping Use
Immediately after each milking, use product. Dip, foam, or spray, the full length of each teat. Allow to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use
When a cow is being dried off, the product should be applied to the teats, once a day for 3-4 days after the last milking. Applications should resume at least twice daily one week before calving.



AgroChem

26 Freedom Way - Saratoga Springs, NY 12866 - (518)226-4850

XT- 22C CONCENTRATE (PART B)

sodium chlorite solution solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:17307-3390
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
sodium chlorite (UNII: G538EBV4VF) (CHLORITE ION - UNII:Z63H374SB6)	sodium chlorite	0.114 kg in 1 kg

Inactive Ingredients

Ingredient Name	Strength
water (UNII: 059QF0K00R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17307-3390-1	19 kg in 1 CONTAINER		
2	NDC:17307-3390-2	57 kg in 1 DRUM		
3	NDC:17307-3390-3	208 kg in 1 DRUM		
4	NDC:17307-3390-4	1040 kg in 1 DRUM		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		03/16/2021	

Labeler - AgroChem Inc (805427700)

Registrant - AgroChem Inc (805427700)

Establishment

Name	Address	ID/FEI	Business Operations
AgroChem Inc		805427700	api manufacture, manufacture

Revised: 3/2021

AgroChem Inc