

AMOXI-TABS- amoxicillin tablet, film coated
Zoetis Inc.

amoxi♦tabs®
(amoxicillin tablets), USP

Veterinary Tablets

For use in dogs and cats

CAUTION

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION

Amoxi-Tabs (amoxicillin tablets) is a semisynthetic antibiotic with a broad spectrum of activity. It provides bactericidal activity against a wide range of common gram-positive and gram-negative pathogens. Chemically, it is D(-)- α -amino-p-hydroxybenzyl penicillin trihydrate.

CLINICAL PHARMACOLOGY

Amoxi-Tabs is stable in the presence of gastric acid and may be given without regard to meals. It is rapidly absorbed after oral administration. It diffuses readily into most body tissues and fluids with the exception of brain and spinal fluid, except when meninges are inflamed. Most of the amoxicillin is excreted unchanged in the urine.

Amoxicillin is similar to ampicillin in its bactericidal action against susceptible organisms. It acts through the inhibition of biosynthesis of cell wall mucopeptide. *In vitro* and/or *in vivo* studies have demonstrated the susceptibility of most strains of the following gram-positive and gram-negative bacteria: a- and b-haemolytic streptococci, nonpenicillinase-producing staphylococci, *Streptococcus faecalis*, *Escherichia coli*, and *Proteus mirabilis*. Because it does not resist destruction by penicillinase, it is not effective against penicillinase-producing bacteria, particularly resistant staphylococci. All strains of *Pseudomonas* and most strains of *Klebsiella* and *Enterobacter* are resistant.

INDICATIONS AND USAGE

Dogs: Amoxi-Tabs are indicated in the treatment of susceptible strains of the organisms causing the following infections:

Respiratory tract infections (tonsillitis, tracheobronchitis) due to *Staphylococcus aureus*, *Streptococcus* spp., *E. coli*, and *Proteus mirabilis*. Genitourinary tract infections (cystitis) due to *Staphylococcus aureus*, *Streptococcus* spp., *E. coli*, and *Proteus mirabilis*.

Genitourinary tract infections(cystitis) due to *Staphylococcus aureus*,

Streptococcus spp., *E. coli*, and *Proteus mirabilis*.

Gastrointestinal tract infections (bacterial gastroenteritis) due to *Staphylococcus aureus*, *Streptococcus spp.*, *E. coli*, and *Proteus mirabilis*. Bacterial dermatitis due to *Staphylococcus aureus*, *Streptococcus spp.*, and *Proteus mirabilis*.

Bacterial dermatitis due to *Staphylococcus aureus*, *Streptococcus spp.*, and *Proteus mirabilis*.

Soft tissue infections (abscesses, lacerations, and wounds) due to *Staphylococcus aureus*, *Streptococcus spp.*, *E. coli*, and *Proteus mirabilis*.

Cats: Amoxi-Tabs are indicated in the treatment of susceptible strains of the organisms causing the following infections:

Upper respiratory tract infections due to *Staphylococcus aureus*, *Streptococcus spp.*, and *E. coli*.

Genitourinary tract infections (cystitis) due to *Staphylococcus aureus*, *Streptococcus spp.*, *E. coli*, and *Proteus mirabilis*.

Gastrointestinal tract infections due to *E. coli*. Skin and soft tissue infections (abscesses, lacerations, and wounds) due to *Staphylococcus aureus*, *Streptococcus spp.*, *E. coli*, and *Pasteurella multocida*.

As with all antibiotics, appropriate *in vitro* culturing and susceptibility testing of samples taken before treatment should be conducted.

CONTRAINDICATIONS

The use of this drug is contraindicated in animals with a history of an allergic reaction to penicillin.

WARNING

For use in dogs and cats only.

ADVERSE REACTIONS

Amoxicillin is a semisynthetic penicillin and has the potential for producing allergic reactions. If an allergic reaction occurs, administer epinephrine and/or steroids.

DOSAGE AND ADMINISTRATION

DOSAGE AND ADMINISTRATION:

Dogs: The recommended dosage is 5 mg/lb of body weight twice a day.

Cats: The recommended dosage is 50 mg (5–10 mg/lb) once a day.

Dosage should be continued for 5–7 days or 48 hours after all symptoms have subsided. If no improvement is seen in 5 days, review diagnosis and change therapy.

Do Not Store at Temperatures Above 25°C (77°F)

Keep Bottle Tightly Closed.

HOW SUPPLIED

Amoxi-Tabs are supplied in 5 strengths: 50 mg, 100 mg, 150 mg, and 200 mg in bottles of 500 tablets; 400 mg in bottles of 250 tablets.

Approved by FDA under NADA # 055-078

Approved by FDA under NADA # 055-081

zoetis

Distributed by:
Zoetis Inc.
Kalamazoo, MI 49007

P1523357
Revised: January 2020

PRINCIPAL DISPLAY PANEL - 150 mg Tablet Bottle Label

Recommended Dosage:
Dogs: 5 mg/lb of body weight twice a day.
Dosage should be continued for 5–7 days or 48 hours after all symptoms have subsided.

Warning: For use in dogs only.
Do Not Store at Temperatures Above 25°C (77°F)
Keep Bottle Tightly Closed.
Read Insert for Complete Directions

amoxi tabs[®]
(amoxicillin tablets), USP
Veterinary Tablets
For use in dogs only
Each film-coated tablet contains amoxicillin trihydrate equivalent to 150 mg of amoxicillin USP

Caution: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

150 mg 

500 tablets
Approved by FDA under NADA # 055-078

zoetis

P1425206

Product of Spain
Distributed by:
Zoetis Inc.
Kalamazoo, MI 49007
Code: TS/DRUGS/57/2003
Lot
Exp

0 187219 01235 2

PRINCIPAL DISPLAY PANEL - 200 mg Tablet Bottle Label

Recommended Dosage:
Dogs: 5 mg/lb of body weight twice a day.
Dosage should be continued for 5–7 days or 48 hours after all symptoms have subsided.

Warning: For use in dogs only.

Do Not Store at Temperatures Above 25°C (77°F)

Keep Bottle Tightly Closed.

Read Insert for Complete Directions

amoxi♦tabs®
(amoxicillin tablets), USP
Veterinary Tablets

For use in dogs only

Each film-coated tablet contains amoxicillin trihydrate equivalent to 200 mg of amoxicillin USP

Caution: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

P1425207

200 mg 

500 tablets

Approved by FDA under NADA # 055-078

zoetis

Product of Spain
Distributed by:
Zoetis Inc.
Kalamazoo, MI 49007
Code: TS/DRUG S/57/2003
Lot
Exp

0
8721901233
8



PRINCIPAL DISPLAY PANEL - 400 mg Tablet Bottle Label

Recommended Dosage:

Dogs: 5 mg/lb of body weight twice a day. Dosage should be continued for 5–7 days or 48 hours after all symptoms have subsided.

Warning: Not for use in animals which are raised for food production.

Do Not Store at Temperatures Above 25°C (77°F)
Keep Bottle Tightly Closed. Read Insert for Complete Directions



amoxi♦tabs®

(amoxicillin tablets), USP

Veterinary Tablets

For use in dogs only

Each film-coated tablet contains amoxicillin trihydrate equivalent to 400 mg of amoxicillin USP

Caution: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

400 mg



250 tablets

Approved by FDA under NADA # 055-078

P1425208

Product of Spain
Distributed by:
Zoetis Inc.
Kalamazoo, MI 49007
Code: TS/DRUGS/57/2003

Lot
Exp

AMOXI-TABS

amoxicillin tablet, film coated

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:54771-8007
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
(2S,5R,6R)-6-((R)-(-)-2-AMINO-2-(P-HYDROXYPHENYL)ACETAMIDO)-3,3-DIMETHYL-7-OXO-4-THIA-1-AZABICYCL(3.2.0)HEPTANE-2-CARBOXYLIC ACID TRIHYDRATE (UNII: 804826J2HU) (AMOXICILLIN ANHYDROUS - UNII:9EM05410Q9)	AMOXICILLIN ANHYDROUS	150 mg

Product Characteristics

Color	purple	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	BMP208
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54771-8007-7	500 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA055078	02/04/1976	

AMOXI-TABS

amoxicillin tablet, film coated

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:54771-6044
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
(2S,5R,6R)-6-((R)-(-)-2-AMINO-2-(P-HYDROXYPHENYL)ACETAMIDO)-3,3-DIMETHYL-7-OXO-4-THIA-1-AZABICYCL(3.2.0)HEPTANE-2-CARBOXYLIC ACID TRIHYDRATE (UNII: 804826J2HU) (AMOXICILLIN ANHYDROUS - UNII:9EM05410Q9)	AMOXICILLIN ANHYDROUS	200 mg

Product Characteristics

Color	orange (Coral)	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	BMP203
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54771-6044-4	500 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA055078	02/04/1976	

AMOXI-TABS

amoxicillin tablet, film coated

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:54771-6046
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
(2S,5R,6R)-6-((R)-(-)-2-AMINO-2-(P-HYDROXYPHENYL)ACETAMIDO)-3,3-DIMETHYL-7-OXO-4-THIA-1-AZABICYCL(3.2.0)HEPTANE-2-CARBOXYLIC ACID TRIHYDRATE (UNII: 804826J2HU) (AMOXICILLIN ANHYDROUS - UNII:9EM05410Q9)	AMOXICILLIN ANHYDROUS	400 mg

Product Characteristics

Color	green	Score	no score
Shape	ROUND	Size	14mm
Flavor		Imprint Code	BMP196
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54771-6046-6	250 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA055078	02/04/1976	

Labeler - Zoetis Inc. (828851555)