

**AMPICILLIN- ampicillin injection powder, for solution**  
**Methapharm Inc.**

-----

**AMPICILLIN FOR INJECTION, USP**  
**For Intravenous or Intramuscular Injection**  
**Rx only**

To reduce the development of drug-resistant bacteria and maintain the effectiveness of ampicillin and other antibacterial drugs, ampicillin should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.


**DESCRIPTION**

Ampicillin for Injection, USP the monosodium salt of [2S-[2 $\alpha$ , 5 $\alpha$ , 6 $\beta$ (S\*)]]-6-[(aminophenylacetyl)amino]-3,3-dimethyl-7-oxo-4-thia-1-azabicyclo [3.2.0] heptane-2-carboxylic acid, is a synthetic penicillin. It is an antibacterial agent with a broad spectrum of bactericidal activity against both penicillin-susceptible Gram-positive organisms and many common Gram-negative pathogens.

Ampicillin for Injection, USP is a dry, white to off-white powder. The reconstituted solution is clear, colorless and free from visible particulates.

Each vial of Ampicillin for Injection, USP contains ampicillin sodium equivalent to 250 mg, 500 mg, 1 gram or 2 grams ampicillin. Ampicillin for Injection, USP contains 65.8 mg [2.9 mEq] sodium per gram ampicillin.

It has the following molecular structure:



The molecular formula is C<sub>16</sub>H<sub>18</sub>N<sub>3</sub>NaO<sub>4</sub>S, and the molecular weight is 371.39. The pH range of the reconstituted solution is 8 to 10.

**CLINICAL PHARMACOLOGY**

Ampicillin for Injection, USP diffuses readily into most body tissues and fluids. However, penetration into the cerebrospinal fluid and brain occurs only when the meninges are inflamed. Ampicillin is excreted largely unchanged in the urine and its excretion can be delayed by concurrent administration of probenecid. Due to maturational changes in renal function, ampicillin half-life decreases as postmenstrual age (a sum of gestational age and postnatal age) increases for infants with postnatal age of less than 28 days. The active form appears in the bile in higher concentrations than those found in serum. Ampicillin is the least serum-bound of all the penicillins, averaging about 20% compared to approximately 60 to 90% for other penicillins. Ampicillin for Injection, USP is well tolerated by most patients and has been given in doses of 2 grams daily for many weeks without adverse reactions.

## Microbiology

While in vitro studies have demonstrated the susceptibility of most strains of the following organisms, clinical efficacy for infections other than those included in the **INDICATIONS AND USAGE** section has not been demonstrated.

### *Antibacterial Activity*

The following bacteria have been shown in in vitro studies to be susceptible to Ampicillin for Injection, USP:

#### Gram-positive Bacteria

Hemolytic and nonhemolytic streptococci

*Streptococcus pneumoniae*

Nonpenicillinase-producing staphylococci

*Clostridium spp.*

*B. anthracis*

*Listeria monocytogenes*

Most strains of enterococci.

#### Gram-negative Bacteria

*H. influenzae*

*N. gonorrhoeae*

*N. meningitidis*

*Proteus mirabilis*

Many strains of *Salmonella*, *Shigella*, and *E. coli*.

AMPICILLIN does not resist destruction by penicillinase.

### *Susceptibility Testing*

For specific information regarding susceptibility test interpretive criteria and associated test methods and quality control standards recognized by FDA for this drug, please see: <https://www.fda.gov/STIC>.

## INDICATIONS AND USAGE

Ampicillin for Injection, USP is indicated in the treatment of infections caused by susceptible strains of the designated organisms in the following conditions:

**Respiratory Tract Infections** caused by *Streptococcus pneumoniae*, *Staphylococcus aureus* (penicillinase and nonpenicillinase-producing), *H. influenzae*, and Group A beta-hemolytic Streptococci.

**Bacterial Meningitis** caused by *E. coli*, Group B Streptococci, and other Gram-negative bacteria (*Listeria monocytogenes*, *N. meningitidis*). The addition of an aminoglycoside with ampicillin may increase its effectiveness against Gram-negative bacteria.

**Septicemia and Endocarditis** caused by susceptible Gram-positive organisms including *Streptococcus spp.*, penicillin G-susceptible staphylococci, and enterococci. Gram-negative sepsis caused by *E. coli*, *Proteus mirabilis* and *Salmonella spp.* responds to ampicillin. Endocarditis due to enterococcal strains usually respond to intravenous therapy. The addition of an aminoglycoside may enhance the effectiveness of ampicillin when treating streptococcal endocarditis.

**Urinary Tract Infections** caused by sensitive strains of *E. coli* and *Proteus mirabilis*.

**Gastrointestinal Infections** caused by *Salmonella typhi* (typhoid fever), other *Salmonella spp.*, and *Shigella spp.* (dysentery) usually respond to oral or intravenous therapy.

Bacteriology studies to determine the causative organisms and their susceptibility to ampicillin should be performed. Therapy may be instituted prior to obtaining results of susceptibility testing.

It is advisable to reserve the parenteral form of this drug for moderately severe and severe infections and for patients who are unable to take the oral forms. A change to oral ampicillin may be made as soon as appropriate.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Ampicillin for Injection, USP and other antibacterial drugs, Ampicillin for Injection, USP should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

Indicated surgical procedures should be performed

## **CONTRAINDICATIONS**

A history of a previous hypersensitivity reaction to any of the penicillins is a contraindication.

## **WARNINGS**

Serious and occasionally fatal hypersensitivity (anaphylactoid) reactions have been reported in patients on penicillin therapy. Although anaphylaxis is more frequent following parenteral therapy, it has occurred in patients on oral penicillins. These reactions are more apt to occur in individuals with a history of penicillin hypersensitivity and/or a history of sensitivity to multiple allergens.

There have been well-documented reports of individuals with a history of penicillin hypersensitivity reactions who have experienced severe hypersensitivity reactions when treated with a cephalosporin. Before initiating therapy with a penicillin, careful inquiry should be made concerning previous hypersensitivity reactions to penicillins, cephalosporins, and other allergens. If an allergic reaction occurs, the drug should be discontinued and appropriate therapy instituted.

**SERIOUS ANAPHYLACTOID REACTIONS REQUIRE IMMEDIATE EMERGENCY TREATMENT WITH EPINEPHRINE, OXYGEN, INTRAVENOUS STEROIDS, AND AIRWAY MANAGEMENT, INCLUDING INTUBATION, SHOULD ALSO BE ADMINISTERED AS INDICATED.**

*Clostridium difficile* associated diarrhea (CDAD) has been reported with use of nearly all anti-bacterial agents, including ampicillin for injection, and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

*C. difficile* produces toxins A and B which contribute to the development of CDAD.

Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibiotic use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing antibiotic use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibiotic treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated

## **PRECAUTIONS**

### **General**

The possibility of superinfections with mycotic organisms or bacterial pathogens should be kept in mind during therapy. In such cases, discontinue the drug and substitute appropriate treatment.

A high percentage (43 to 100 percent) of patients with infectious mononucleosis who receive ampicillin develop a skin rash. Typically, the rash appears 7 to 10 days after the start of oral ampicillin therapy and remains for a few days to a week after the drug is discontinued. In most cases, the rash is maculopapular, pruritic, and generalized. Therefore, the administration of ampicillin is not recommended in patients with mononucleosis. It is not known whether these patients are truly allergic to ampicillin.

Prescribing ampicillin for injection in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

### **Information for Patients**

Patients should be counseled that antibacterial drugs including ampicillin should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When ampicillin is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may: (1) decrease the effectiveness of the immediate treatment, and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by ampicillin or other antibacterial drugs in the future.

Diarrhea is a common problem caused by antibiotics which usually ends when the antibiotic is discontinued. Sometimes after starting treatment with antibiotics, patients can develop watery and bloody stools (with or without stomach cramps and fever) even as late as two or more months after having taken the last dose of the antibiotic. If this occurs, patients should contact their physician as soon as possible.

### **Laboratory Tests**

As with any potent drug, periodic assessment of organ system function, including renal, hepatic, and hematopoietic, should be made during prolonged therapy.

Transient elevation of serum transaminase has been observed following administration

of ampicillin. The significance of this finding is not known.

### **Drug Interactions**

The concurrent administration of allopurinol and ampicillin increases substantially the incidence of skin rashes in patients receiving both drugs as compared to patients receiving ampicillin alone. It is not known whether this potentiation of ampicillin rashes is due to allopurinol or the hyperuricemia present in these patients.

### **Drug/Laboratory Test Interactions**

With high urine concentrations of ampicillin, false-positive glucose reactions may occur if Clinitest, Benedict's Solution, or Fehling's Solution are used. Therefore, it is recommended that glucose tests based on enzymatic glucose oxidase reactions (such as Clinistix or Tes-Tape) be used.

### **Carcinogenesis, Mutagenesis, and Impairment of Fertility**

No long-term animal studies have been conducted with this drug.

### **Pregnancy - Category B**

Reproduction studies have been performed in laboratory animals at doses several times the human dose and have revealed no evidence of adverse effects due to ampicillin. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

### **Labor and Delivery**

Oral ampicillin-class antibiotics are poorly absorbed during labor. Studies in guinea pigs showed that intravenous administration of ampicillin slightly decreased the uterine tone and frequency of contractions, but moderately increased the height and duration of contractions. However, it is not known whether use of these drugs in humans during labor or delivery has immediate or delayed adverse effects on the fetus, prolongs the duration of labor, or increases the likelihood that forceps delivery or other obstetrical intervention or resuscitation of the newborn will be necessary.

### **Nursing Mothers**

Ampicillin is excreted in trace amounts in human milk. Therefore, caution should be exercised when ampicillin-class antibiotics are administered to a nursing woman.

### **Pediatric Use**

Guidelines for the administration of these drugs to children, including neonates are presented in **DOSAGE AND ADMINISTRATION** section.

## **ADVERSE REACTIONS**

As with other penicillins, it may be expected that untoward reactions will be essentially limited to sensitivity phenomena. They are more likely to occur in individuals who have previously demonstrated hypersensitivity to penicillins and in those with a history of

allergy, asthma, hay fever, or urticaria.

The following adverse reactions have been reported as associated with the use of ampicillin:

### **Gastrointestinal**

Glossitis, stomatitis, black “hairy” tongue, nausea, vomiting, enterocolitis, pseudomembranous colitis, and diarrhea. (These reactions are usually associated with oral dosage forms.)

### **Hypersensitivity Reactions**

Skin rashes and urticaria have been reported frequently. A few cases of exfoliative dermatitis and erythema multiforme have been reported. Linear IgA bullous dermatosis has been reported. Anaphylaxis is the most serious reaction experienced and has usually been associated with the parenteral dosage form.

**Note:** Urticaria, other skin rashes, and serum sickness-like reactions may be controlled with antihistamines and, if necessary, systemic corticosteroids. Whenever such reactions occur, ampicillin should be discontinued, unless, in the opinion of the physician, the condition being treated is life-threatening and amenable only to ampicillin therapy. Serious anaphylactic reactions require the immediate use of epinephrine, oxygen, and intravenous steroids.

**Liver** - A moderate rise in serum glutamic oxaloacetic transaminase (SGOT) has been noted, particularly in infants, but the significance of this finding is unknown. Mild transitory SGOT elevations have been observed in individuals receiving larger (two to four times) than usual and oft-repeated intramuscular injections. Evidence indicates that glutamic oxaloacetic transaminase (GOT) is released at the site of intramuscular injection of ampicillin sodium and that the presence of increased amounts of this enzyme in the blood does not necessarily indicate liver involvement.

**Hemic and Lymphatic Systems** - Anemia, thrombocytopenia, thrombocytopenic purpura, eosinophilia, leukopenia, and agranulocytosis have been reported during therapy with the penicillins. These reactions are usually reversible on discontinuation of therapy and are believed to be hypersensitivity phenomena.

**Central Nervous System** - Seizures

**To report SUSPECTED ADVERSE REACTIONS, contact Methapharm, Inc. at 1-800-287-7686 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

### **OVERDOSAGE**

In cases of overdose, discontinue medication, treat symptomatically, and institute supportive measures as required. In patients with renal function impairment, ampicillin-class antibiotics can be removed by hemodialysis but not peritoneal dialysis.

### **DOSAGE AND ADMINISTRATION**

Infections of the respiratory tract and soft tissues.

Patients weighing 40 kg (88 lbs) or more: 250 mg to 500 mg every 6 hours.

Patients weighing less than 40 kg (88 lbs): 25 to 50 mg/kg/day in equally divided doses

at 6- to 8-hour intervals.

Infections of the gastrointestinal and genitourinary tracts (including those caused by *Neisseria gonorrhoeae* in females).

Patients weighing 40 kg (88 lbs) or more: 500 mg every 6 hours.

Patients weighing less than 40 kg (88 lbs): 50 mg/kg/day in equally divided doses at 6- to 8- hour intervals.

In the treatment of chronic urinary tract and intestinal infections, frequent bacteriological and clinical appraisal is necessary. Smaller doses than those recommended above should not be used. Higher doses should be used for stubborn or severe infections. In stubborn infections, therapy may be required for several weeks. It may be necessary to continue clinical and/or bacteriological follow-up for several months after cessation of therapy.

Urethritis in males due to *N. gonorrhoeae*.

**Adults** - Two doses of 500 mg each at an interval of 8 to 12 hours. Treatment may be repeated if necessary or extended if required.

In the treatment of complications of gonorrheal urethritis, such as prostatitis and epididymitis, prolonged and intensive therapy is recommended. Cases of gonorrhea with a suspected primary lesion of syphilis should have darkfield examinations before receiving treatment. In all other cases where concomitant syphilis is suspected, monthly serological tests should be made for a minimum of four months.

The doses for the preceding infections may be given by either the intramuscular or intravenous route. A change to oral ampicillin may be made when appropriate.

### **Bacterial Meningitis**

**Adults and children** - 150 to 200 mg/kg/day in equally divided doses every 3 to 4 hours. (Treatment may be initiated with intravenous drip therapy and continued with intramuscular injections.) The doses for other infections may be given by either the intravenous or intramuscular route.

**Neonates (less than or equal to 28 days of postnatal age)** - Dosage should be based on Gestational age and Postnatal age according to Table 3.

**Table 3: Dosage in Neonates (less than or equal to 28 days of postnatal age) for Bacterial Meningitis and Septicemia:**

Gestational Age (weeks)	Postnatal Age (days)	Dosage
less than or equal to 34	less than or equal to 7	100 mg/kg/day in equally divided doses every 12 hours
less than or equal to 34	greater than or equal to 8 and less than 28	150 mg/kg/day in equally divided doses every 12 hours
greater than 34	less than or equal to 28	150 mg/kg/day in equally divided doses every 8 hours

### **Septicemia**

**Adults and children** - 150 to 200 mg/kg/day. Start with intravenous administration for

at least three days and continue with the intramuscular route every 3 to 4 hours.

**Neonates (less than or equal to 28 days of postnatal age)** - Dosage should be based on Gestational age and Postnatal age according to Table 3 (above).

Treatment of all infections should be continued for a minimum of 48 to 72 hours beyond the time that the patient becomes asymptomatic or evidence of bacterial eradication has been obtained. A minimum of 10-days treatment is recommended for any infection caused by Group A beta-hemolytic streptococci to help prevent the occurrence of acute rheumatic fever or acute glomerulonephritis.

### **DIRECTIONS FOR USE**

Use only freshly prepared solutions. Intramuscular and intravenous injections should be administered within one hour after preparation since the potency may decrease significantly after this period.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

**For Intramuscular Use** - Dissolve contents of a vial with the amount of Sterile Water for Injection, USP, or Bacteriostatic Water for Injection, USP, listed in the table below:

<b>NDC</b>	<b>Label Claim</b>	<b>Recommended Amount of Diluent</b>	<b>Withdrawable Volume</b>	<b>Concentration (in mg/mL)</b>
67850-023-10	250 mg	1 mL	1 mL	250 mg
67850-024-10	500 mg	1.8 mL	2 mL	250 mg
67850-021-10	1 g	3.5 mL	4 mL	250 mg
67850-022-10	2 g	6.8 mL	8 mL	250 mg

While Ampicillin for Injection, USP, 1 g and 2 g, are primarily for intravenous use, they may be administered intramuscularly when the 250 mg or 500 mg vials are unavailable. In such instances, dissolve in 3.5 or 6.8 mL Sterile Water for Injection, USP, or Bacteriostatic Water for Injection, USP, respectively. The resulting solution will provide a concentration of 250 mg per mL.

Ampicillin for Injection, USP, 125 mg, is intended primarily for pediatric use. It also serves as a convenient dosage form when small parenteral doses of the antibiotic are required.

Bacteriostatic Water for Injection, USP is not to be used as a diluent when the product will be used in newborns.

**For Direct Intravenous Use** - Add 5 mL Sterile Water for Injection, USP, or Bacteriostatic Water for Injection, USP to the 250, and 500 mg vials and administer slowly over a 3- to 5- minute period.

Ampicillin for Injection, USP, 1 g or 2 g, may also be given by direct intravenous administration. Dissolve in 7.4 or 14.8 mL Sterile Water for Injection, USP, or Bacteriostatic Water for Injection, USP, respectively, and administer slowly over at least

10 to 15 minutes.

CAUTION: More rapid administration may result in convulsive seizures.

**For Administration by Intravenous Drip** - Reconstitute as directed above (For Direct Intravenous Use) prior to diluting with Intravenous Solution. Stability studies on ampicillin sodium at several concentrations in various intravenous solutions indicate the drug will lose less than 10% activity at the temperatures noted for the time periods stated.

### Room Temperature (25°C)

<b>Diluent</b>	<b>Concentrations</b>	<b>Stability Periods</b>
Sterile water for Injection	up to 30 mg/mL	8 hours
0.9% Sodium Chloride Injection, USP	up to 30 mg/mL	8 hours
5% Dextrose Injection, USP	10 to 20 mg/mL	1 hour
5% Dextrose Injection, USP	up to 2 mg/mL	2 hours
5% Dextrose and 0.45% Sodium Chloride Injection, USP	up to 2 mg/mL	2 hours
Lactated Ringer's Injection, USP	up to 30 mg/mL	8 hours

### Refrigerated (4°C)

<b>Diluent</b>	<b>Concentrations</b>	<b>Stability Periods</b>
Sterile Water for Injection	30 mg/mL	48 hours
Sterile Water for Injection	up to 20 mg/mL	72 hours
0.9% Sodium Chloride Injection, USP	30 mg/mL	24 hours
0.9% Sodium Chloride Injection, USP	up to 20 mg/mL	48 hours
Lactated Ringer's Injection, USP	up to 30 mg/mL	24 hours
5% Dextrose Injection, USP	up to 20 mg/mL	1 hour
5% Dextrose and 0.45% Sodium Chloride Injection, USP	up to 10 mg/mL	1 hour

Only those solutions listed above should be used for the intravenous infusion of Ampicillin for Injection, USP. The concentrations should fall within the range specified. The drug concentration and the rate and volume of the infusion should be adjusted so that the total dose of ampicillin is administered before the drug loses its stability in the solution in use.

### HOW SUPPLIED

Ampicillin for Injection, USP, is supplied as follows:

<b>NDC</b>	<b>Ampicillin for Injection, USP</b>	<b>Package Factor</b>
67850-023-10	250 mg per vial	10 vials per carton
67850-024-10	500 mg per vial	10 vials per carton

67850-021-10	1 gram per vial	10 vials per carton
67850-022-10	2 grams per vial	10 vials per carton

Each vial contains ampicillin sodium equivalent to 250 mg, 500 mg, 1 gram or 2 grams ampicillin per vial.

### Storage Conditions

Store dry powder at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]

### Sterile, Nonpyrogenic, Preservative-free.

**The container closure is not made with natural rubber latex.**

Clinitest is a registered trademark of Miles, Inc.

Clinistix is a registered trademark of Bayer Corporation.

Tes-Tape is a registered trademark of Eli Lilly Company.



Manufactured for:

Methapharm, Inc.  
Coral Springs, FL,  
33065 USA

### Made in Romania

Revised: February 2024

### Sample Distributor Labels:

Cartons and Labels will be included for listings under Private Label Distributors

NDC 67850-023-00

**Ampicillin for Injection, USP**

**250 mg\* per vial**

**For Intravenous or Intramuscular Use**

**Rx only**

LOT: \_\_\_\_\_

EXP.: \_\_\_\_\_

**\*This vial contains ampicillin sodium equivalent to 250 mg ampicillin.**

**For I.M. Use,** add 1 mL diluent (read accompanying insert). Resulting solution contains 250 mg ampicillin per mL.

**Use solution within 1 hour.**

**Usual Dosage:** Adults - 250 to 500 mg I.M. q6h. **READ ACCOMPANYING INSERT** for detailed indications, I.M. or I.V. dosage, and precautions. Store dry powder at 20° to 25°C (68° to 77°F) [See USP].

**Manufactured for Methapharm, Inc.**  
Coral Springs, FL, 33065 USA  
**Made in Romania**

(01)00367850023000

P011VUS4/00A

NDC 67850-024-00  
**Ampicillin for Injection, USP**

**500 mg\* per vial**

**For Intravenous or Intramuscular Use**  
 **Rx only**

**\*This vial contains ampicillin sodium equivalent to 500 mg ampicillin.**  
**For I.M. Use,** add 1.8 mL diluent (read accompanying insert). Resulting solution contains 250 mg ampicillin per mL.  
**Use solution within 1 hour.**  
**Usual Dosage:** Adults - 250 to 500 mg I.M. q6h. **READ ACCOMPANYING INSERT** for detailed indications, I.M. or I.V. dosage, and precautions. Store dry powder at 20° to 25°C (68° to 77°F) [See USP].  
**Manufactured for Methapharm, Inc.**  
 Coral Springs, FL, 33065 USA  
**Made in Romania**



P012VUS4/00A

LOT:  
 EXP.:

NDC 67850-021-00  
**Ampicillin for Injection, USP**

**1 gram\* per vial**

**For Intravenous or Intramuscular Use**  
 **Rx only**

**\*This vial contains ampicillin sodium equivalent to 1 gram ampicillin.**  
**For I.M. Use,** add 3.5 mL diluent (read accompanying insert). Resulting solution contains 250 mg ampicillin per mL.  
**Use solution within 1 hour.**  
**Usual Dosage:** Adults - 250 to 500 mg I.M. q6h. **READ ACCOMPANYING INSERT** for detailed indications, I.M. or I.V. dosage, and precautions. Store dry powder at 20° to 25°C (68° to 77°F) [See USP].  
**Manufactured for Methapharm, Inc.**  
 Coral Springs, FL, 33065 USA  
**Made in Romania**



P013VUS4/00A

LOT:  
 EXP.:

NDC 67850-022-00  
**Ampicillin for Injection, USP**

**2 grams\* per vial**

**For Intravenous Use**  
 **Rx only**

**\*This vial contains ampicillin sodium equivalent to 2 grams ampicillin.**  
**For Direct I.V. Use,** add 14.8 mL Sterile Water for Injection, USP.  
**I.V. injection: Use solution within 1 hour.**  
**I.V. infusion:** See accompanying insert.  
**Usual Dosage:** Adults - 250 to 500 mg I.V. q6h. **READ ACCOMPANYING INSERT** for detailed indications, dosage, and precautions. Store dry powder at 20° to 25°C (68° to 77°F) [See USP].  
**Manufactured for Methapharm, Inc.**  
 Coral Springs, FL, 33065 USA  
**Made in Romania**



P014VUS4/00A

LOT:  
 EXP.:



<p>Rx only</p> <p>UNVARNISH 60x30mm</p> <p>Ampicillin for Injection, USP</p> <p>500 mg* per vial</p> <p>Ten 500 mg* vials NDC 67850-024-10</p> <p>For Intravenous or Intramuscular use</p>			
<p>Sterile, Nonpyrogenic, Preservative-free. The container closure is not made with natural rubber latex. Read the leaflet before use.</p> <p>*Each vial contains ampicillin sodium equivalent to 500 mg ampicillin. For I.M. Use, add 1.8 mL diluent (read accompanying insert). Resulting solution contains 250 mg ampicillin per mL. Use solution within 1 hour.</p> <p>Usual Dosage: Adults - 250 to 500 mg I.M. q6h. READ ACCOMPANYING INSERT for detailed indications, I.M. or I.V. dosage, and precautions.</p> <p>Store dry powder at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].</p> <p>Manufactured for Methapharm, Inc., Coral Springs, FL, 33065 USA Made in Romania P013010USA00A</p> 	<p>Ten 500 mg* vials NDC 67850-024-10</p> <p><b>Ampicillin for Injection, USP</b></p> <p>500 mg* per vial</p> <p>For Intramuscular or Intravenous Use Rx only</p>	<p>Ten 500 mg* vials NDC 67850-024-10</p> <p><b>Ampicillin for Injection, USP</b></p> <p>500 mg* per vial</p> <p>For Intravenous or Intramuscular use</p> <p><b>Methapharm</b> Specialty Pharmaceuticals</p> <p>Rx only</p>	<p>Ten 500 mg* vials NDC 67850-024-10</p> <p><b>Ampicillin for Injection, USP</b></p> <p>500 mg* per vial</p> <p>For Intramuscular or Intravenous Use Rx only</p>

<p>Rx only</p> <p>UNVARNISH 60x30mm</p> <p>Ampicillin for Injection, USP</p> <p>1 gram* per vial</p> <p>Ten 1 gram* vials NDC 67850-021-10</p> <p>For Intravenous or Intramuscular use</p>			
<p>Sterile, Nonpyrogenic, Preservative-free. The container closure is not made with natural rubber latex. Read the leaflet before use.</p> <p>*Each vial contains ampicillin sodium equivalent to 1 gram ampicillin. For I.M. Use, add 3.5 mL diluent (read accompanying insert). Resulting solution contains 250 mg ampicillin per mL. Use solution within 1 hour.</p> <p>Usual Dosage: Adults - 250 to 500 mg I.M. q6h. READ ACCOMPANYING INSERT for detailed indications, I.M. or I.V. dosage, and precautions.</p> <p>Store dry powder at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].</p> <p>Manufactured for Methapharm, Inc., Coral Springs, FL, 33065 USA Made in Romania P013010USA00A</p> 	<p>Ten 1 gram* vials NDC 67850-021-10</p> <p><b>Ampicillin for Injection, USP</b></p> <p>1 gram* per vial</p> <p>For Intramuscular or Intravenous Use Rx only</p>	<p>Ten 1 gram* vials NDC 67850-021-10</p> <p><b>Ampicillin for Injection, USP</b></p> <p>1 gram* per vial</p> <p>For Intravenous or Intramuscular use</p> <p><b>Methapharm</b> Specialty Pharmaceuticals</p> <p>Rx only</p>	<p>Ten 1 gram* vials NDC 67850-021-10</p> <p><b>Ampicillin for Injection, USP</b></p> <p>1 gram* per vial</p> <p>For Intramuscular or Intravenous Use Rx only</p>

<p>Rx only</p> <p>UNVARNISH (60x30mm)</p> <p>Ampicillin for Injection, USP</p> <p>2 grams* per vial</p> <p>Ten 2 gram* vials NDC 67850-022-10</p> <p>For Intravenous use</p>			
<p>Sterile, Nonpyrogenic, Preservative-free. The container closure is not made with natural rubber latex. Read the leaflet before use.</p> <p>*Each vial contains ampicillin sodium equivalent to 2 grams of ampicillin. For Direct I.V. Use, add 14.5 mL Sterile Water for Injection, USP. IV Injection: Use solution within 1 hour. IV Infusion: See accompanying insert.</p> <p>Usual Dosage: Adults - 250 to 500 mg I.V. q6h. READ ACCOMPANYING INSERT for detailed indications, dosage, and precautions.</p> <p>Store dry powder at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].</p> <p>Manufactured for Methapharm, Inc., Coral Springs, FL, 33065 USA Made in Romania P01410USA00A</p> 	<p>Ten 2 gram* vials NDC 67850-022-10</p> <p><b>Ampicillin for Injection, USP</b></p> <p>2 grams* per vial</p> <p>For Intravenous use Rx only</p>	<p>Ten 2 gram* vials NDC 67850-022-10</p> <p><b>Ampicillin for Injection, USP</b></p> <p>2 grams* per vial</p> <p>For Intravenous use</p> <p><b>Methapharm</b> Specialty Pharmaceuticals</p> <p>Rx only</p>	<p>Ten 2 gram* vials NDC 67850-022-10</p> <p><b>Ampicillin for Injection, USP</b></p> <p>2 grams* per vial</p> <p>For Intravenous use Rx only</p>

## AMPICILLIN

ampicillin injection powder, for solution

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:67850-023
<b>Route of Administration</b>	INTRAMUSCULAR, INTRAVENOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
AMPICILLIN SODIUM (UNII: JFN36L5S8K) (AMPICILLIN - UNII:7C782967RD)	AMPICILLIN	250 mg in 250 mg

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67850-023-10	10 in 1 PACKAGE	03/06/2019	
1	NDC:67850-023-00	250 mg in 1 VIAL; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090354	03/06/2019	

## AMPICILLIN

ampicillin injection powder, for solution

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:67850-024
<b>Route of Administration</b>	INTRAMUSCULAR, INTRAVENOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
AMPICILLIN SODIUM (UNII: JFN36L5S8K) (AMPICILLIN - UNII:7C782967RD)	AMPICILLIN	500 mg in 500 mg

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67850-024-10	10 in 1 PACKAGE	03/06/2019	

<b>1</b>	NDC:67850-024-00	500 mg in 1 VIAL; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>		<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA090354		03/06/2019	

<b>AMPICILLIN</b>				
ampicillin injection powder, for solution				
<b>Product Information</b>				
<b>Product Type</b>	HUMAN PRESCRIPTION DRUG		<b>Item Code (Source)</b>	NDC:67850-021
<b>Route of Administration</b>	INTRAMUSCULAR, INTRAVENOUS			
<b>Active Ingredient/Active Moiety</b>				
<b>Ingredient Name</b>			<b>Basis of Strength</b>	<b>Strength</b>
AMPICILLIN SODIUM (UNII: JFN36L5S8K) (AMPICILLIN - UNII:7C782967RD)			AMPICILLIN	1 g in 1 g
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
<b>1</b>	NDC:67850-021-10	10 in 1 PACKAGE	03/06/2019	
<b>1</b>	NDC:67850-021-00	1 g in 1 VIAL; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>		<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA090354		03/06/2019	

<b>AMPICILLIN</b>				
ampicillin injection powder, for solution				
<b>Product Information</b>				
<b>Product Type</b>	HUMAN PRESCRIPTION DRUG		<b>Item Code (Source)</b>	NDC:67850-022
<b>Route of Administration</b>	INTRAMUSCULAR, INTRAVENOUS			

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>AMPICILLIN SODIUM</b> (UNII: JFN36L5S8K) (AMPICILLIN - UNII: 7C782967RD)	AMPICILLIN	2 g in 2 g

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
<b>1</b>	NDC:67850-022-10	10 in 1 PACKAGE	03/06/2019	
<b>1</b>	NDC:67850-022-00	2 g in 1 VIAL; Type 0: Not a Combination Product		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA090354	03/06/2019	

**Labeler** - Methapharm Inc. (066672887)

Revised: 5/2024

Methapharm Inc.