

**HEADACHE RELIEF- acetaminophen, aspirin, caffeine tablet, film coated**  
**Target Corporation**

-----  
**Target 44-334B**

***Active ingredients (in each caplet)***

Acetaminophen 250 mg

Aspirin 250 mg (NSAID)\*

Caffeine 65 mg

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever

Pain reliever

Pain reliever aid

***Uses***

- temporarily relieves minor aches and pains due to:
  - headache
  - arthritis
  - a cold
  - muscular aches
  - toothache
  - premenstrual and menstrual cramps

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- shock
- facial swelling
- hives
- asthma (wheezing)

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- 3 or more alcoholic drinks every day while using this product
- with other drugs containing acetaminophen

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

#### **Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

#### **Ask a doctor before use if**

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

#### **Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug or are under a doctor's care for any serious condition

#### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools

- have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- redness or swelling is present
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- new symptoms occur

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours. Do not take more than 8 caplets in 24 hours
- children under 12 years: ask a doctor

### **Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

### **Inactive ingredients**

corn starch, croscopvidone, hypromellose, microcrystalline cellulose, povidone, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

### **Questions?**

**Call 1-800-910-6874**

### **Principal display panel**

**Compare to active ingredients in  
Excedrin® Extra Strength Caplets\*\***

NDC 11673-433-59

extra strength  
**headache relief**  
**acetaminophen** 250 mg  
aspirin 250 mg (**NSAID**)  
caffeine 65 mg

pain reliever/pain reliever aid

**up&up™**

ACTUAL SIZE

**275 CAPLETS†**  
(†CAPSULE-SHAPED TABLETS)

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

50844          REV1119A33459

**100% satisfaction guaranteed  
or your money back.**

Distributed by Target Corporation  
Minneapolis, MN 55403  
TM & ©2021 Target Brands, Inc.

\*\*This product is not manufactured or distributed by  
GSK Consumer Healthcare S.A., owner of the  
registered trademark Excedrin® Extra Strength  
Caplets.

**Drug Facts (continued)**

- Take a blood thinning (anticoagulant) or steroid drug
- Take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- Have 3 or more alcoholic drinks every day while using this product
- Take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

**Do not use**

- If you have ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/fever reducer
- With any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- If you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- You have liver disease
- Stomach bleeding warning applies to you
- You have a history of stomach problems, such as heartburn
- You have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- You have asthma
- You are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- Taking a prescription drug for diabetes, gout, or arthritis
- Taking any other drug or are under a doctor's care for any serious condition

**Stop use and ask a doctor if**

- An allergic reaction occurs. Seek medical help right away.
- You experience any of the following signs of stomach bleeding:
  - Have bloody or black stools
  - Vomit blood
  - Feel faint
  - Have stomach pain that does not get better
  - Ringing in the ears or a loss of hearing occurs
  - Redness or swelling is present
- Pain gets worse or lasts for more than 10 days
- Fever gets worse or lasts for more than 3 days
- New symptoms occur
- These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- Do not take more than directed
- Drink a full glass of water with each dose
- Adults and children 12 years and over: take 2 caplets every 6 hours. Do not take more than 8 caplets in 24 hours.
- Children under 12 years: ask a doctor

**Other Information**

- Store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- Use by expiration date on package

**Inactive ingredients:** corn starch, croscopolone, hypromellose, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

**Questions? Call 1-800-910-8874**

STOP PEELING

NDC 11673-433-59

Compare to active ingredients in  
Excedrin® Extra Strength Caplets\*\*

**extra strength  
headache relief  
acetaminophen 250 mg  
aspirin 250 mg (NSAID)  
caffeine 65 mg**

pain reliever/pain reliever aid




**275**  
CAPLETS

275 CAPLETS† (†CAPSULE-SHAPED TABLETS)

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**Drug Facts**

**Active ingredients (in each caplet)**

Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID)*	Pain reliever
Caffeine 65 mg	Pain reliever aid

\*nonsteroidal anti-inflammatory drug

**Uses**

- Temporarily relieves minor aches and pains due to:
  - Headache
  - Cold
  - Arthritis
  - Toothache
  - Muscular aches
  - Premenstrual and menstrual cramps

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:
 

- Hives
- Facial swelling
- Shock
- Asthma (wheezing)

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
 

- Skin redness
- Blister
- Rash

If a skin reaction occurs, stop use and seek medical help right away.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours

- With other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
 


- Are age 60 or older
- Have had stomach ulcers or bleeding problems

50844 REY119A33459  
094 01 0124 R00 C-001472-01-116

**100% satisfaction guaranteed or your money back.**

Distributed by Target Corporation  
Minneapolis, MN 55403  
TM & ©2021 Target Brands, Inc.

\*\*This product is not manufactured or distributed by GSK Consumer Healthcare S.A., owner of the registered trademark Excedrin® Extra Strength Caplets.



1167343359 1  
3

**PEEL HERE FOR MORE DRUG FACTS**

Target 44-334B

HEADACHE RELIEF			
acetaminophen, aspirin, caffeine tablet, film coated			
<b>Product Information</b>			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-433
Route of Administration	ORAL		
<b>Active Ingredient/Active Moiety</b>			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg	
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg	

## Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSPVIDONE</b> (UNII: 2S7830E561)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	44;334
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-433-12	1 in 1 CARTON	04/01/2019	
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:11673-433-59	275 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/01/2019	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	04/01/2019	

**Labeler** - Target Corporation (006961700)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(11673-433)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(11673-433)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(11673-433)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(11673-433)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(11673-433)

Revised: 12/2023

Target Corporation