

**COUGH DM- dextromethorphan polistirex suspension**  
**Meijer Distribution Inc**

-----  
**Meijer Distribution, Inc. Cough DM Drug Facts**

**Active ingredient (in each 5 mL)**

Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide

**Purpose**

Cough suppressant

**Uses**

temporarily relieves

- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the impulse to cough to help you get to sleep

**Warnings**

**Do not use**

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Allergy Alert:** Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions.

**Ask a doctor before use if you have**

- chronic cough that lasts as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

**Stop use and ask a doctor if**

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222.

**Directions**

- **shake bottle well before use**
- measure only with dosing cup provided. Do not use dosing cup with other products.
- dose as follows or as directed by doctor
- mL = milliliter

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

### Other information

- **each 5 mL contains:** sodium 5 mg
- store at 20° to 25°C (68° to 77°F)
- dosing cup provided

### Inactive Ingredients

D&C Red #30 aluminum lake, D&C Yellow #10 aluminum lake, flavor, glycerin, high fructose corn syrup, methylparaben, polysorbate 80, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, tartaric acid, tragacanth gum, triacetin, xanthan gum

### Questions or comments?

1-800-719-9260

### Package/Label Principal Display Panel

Compare to Children's Delsym® active ingredient

CHILDREN'S cough DM

Dextromethorphan Polistirex

Extended-Release Oral Suspension

Cough Suppressant

AGE 4+

12 HOUR Cough Relief

Orange-Flavored Liquid

Day or Night | Alcohol Free For Children & Adults

Dosing Cup Included

Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions

5 FL OZ (148 mL)





**meijer**  
CHILDREN'S  
**cough DM**  
Dextromethorphan Polistirex  
Extended-Release Oral Suspension  
Cough Suppressant

**AGE 4+ 12 HOUR Cough Relief**  
Orange-Flavored Liquid

Day or Night | Alcohol Free  
For Children & Adults

5 FL OZ (148 mL)

actual size  
Also Available in Grape Flavor



**meijer**  
NDC 41250-958-28

Compare to  
Children's Delsym®  
active ingredient\*

**CHILDREN'S  
cough DM**  
Dextromethorphan Polistirex  
Extended-Release Oral Suspension  
Cough Suppressant

**AGE 4+ 12 HOUR Cough Relief**  
Orange-Flavored Liquid

Day or Night | Alcohol Free  
For Children & Adults

5 FL OZ (148 mL)

Contains sodium metabisulfite, a sulfite  
that may cause allergic-type reactions

**Drug Facts**

**Active ingredient (in each 5 mL)** Purpose  
Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide.....Cough suppressant

**Uses** temporarily relieves  
 ■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants  
 ■ the impulse to cough to help you get to sleep

**Warnings**  
 Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
**Allergy Alert:** Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions.

**Ask a doctor before use if you have**  
 ■ chronic cough that lasts as occurs with smoking, asthma or emphysema  
 ■ cough that occurs with too much phlegm (mucus)

**Stop use and ask a doctor if**  
 ■ side effects occur: You may report side effects to FDA at 1-800-FDA-1088.  
 ■ cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222.**

**Directions** ■ shake bottle well before use  
 ■ measure only with dosing cup provided. Do not use dosing cup with other products.  
 ■ dose as follows or as directed by doctor  
 ■ mL = milliliter

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

**Other information** ■ each 5 mL contains: sodium 5 mg  
 ■ store at 20° to 25°C (68° to 77°F) ■ dosing cup provided

**Inactive ingredients** D&C Red #30 aluminum lake, D&C Yellow #10 aluminum lake, flavor glycerin, high fructose corn syrup, methylparaben, polysorbate 80, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, tartaric acid, tragacanth gum, triacetin, xanthan gum

**Questions or comments?** 1-800-719-9260

\*This product is not manufactured or distributed by Reckitt Benckiser LLC, owner of the registered trademark Delsym®.

**DOSING**  
 SHAKE WELL BEFORE USE.  
 Measure only with dosing cup provided.  
 Do not use dosing cup with other products.  
 mL = milliliter

Age (yr)	Dose
12 years to adult	10 mL EVERY 12 HOURS
6 to under 12	5 mL EVERY 12 HOURS
4 to under 6	2.5 mL EVERY 12 HOURS
Under 4	Do not use

See back panel for full dosing directions.



Dosing cup included

**PARENTS:**  
 Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

DIST. BY MEIJER DISTRIBUTION, INC.  
 GRAND RAPIDS, MI 49544  
[www.meijer.com](http://www.meijer.com)




**TAMPER EVIDENT:**  
 Do not use if carton is opened or neckband imprinted with "sealed for your protection" is broken or missing.

PID 3269528



7 13733 31412 1

LOT NO.

EXP.

1 95826 AE C3

## COUGH DM

dextromethorphan polistirex suspension

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41250-958
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 5 mL

## Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
METHYL PARABEN (UNII: A2I8C7HI9T)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POVIDONE (UNII: FZ989GH94E)	
PROPYL PARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
SODIUM POLYSTYRENE SULFONATE (UNII: 1699G8679Z)	
SUCROSE (UNII: C151H8M554)	
TARTARIC ACID (UNII: W4888I119H)	
TRAGACANTH (UNII: 2944357O2O)	
TRIACETIN (UNII: XHX3C3X673)	
XANTHAN GUM (UNII: TTV12P4NEE)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
D&C RED NO. 30 (UNII: 2S42T2808B)	
POLYVINYL ACETATE (UNII: 32K497ZK2U)	

## Product Characteristics

Color	ORANGE	Score	
Shape		Size	
Flavor	ORANGE	Imprint Code	
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-958-21	1 in 1 CARTON	07/08/2013	09/01/2015
1		89 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:41250-958-28	1 in 1 CARTON	07/31/2013	
2		148 mL in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091135	07/08/2013	

**Labeler** - Meijer Distribution Inc (006959555)