

TUSNEL C- codeine phosphate, guaifenesin, pseudoephedrine hcl liquid
Llorens Pharmaceutical International Division

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active Ingredients (in each 5 ml tsp.)

Purpose

Codeine Phosphate	10 mg	Cough Suppressant
Guaifenesin	100 mg	Expectorant
Pseudoephedrine HCl	30 mg	Nasal Decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily relieves: nasal congestion due to the common cold, hay fever or other upper respiratory allergies and nasal congestion associated with sinusitis, cough due to minor bronchial irritation as may occur with the common cold
- temporarily restores freer breathing through the nose
- calms the cough control center and relieves coughing

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

When using this product

- do not exceed recommended dosage
- may cause or aggravate constipation

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem or persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema. chronic pulmonary disease or shortness of breath, or who are taking other drugs

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless. symptoms do not get better within 7 days or are accompanied by fever. cough lasts more than 7 days, comes back, or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

If pregnant or breast-feeding, ask a health professional before use.

Directions: Follow dosage chart Dosage may be repeated every 6 hours. Not to exceed 4 doses in 24-hours. Giving a higher dose than recommended by a doctor could result in serious side effects for your child. A special measuring device should be used to give an accurate dose of this product to children under 6 years of age.

Age	Weight	Dose
children 12 years of age and over	95 lbs and up	Take 2 teaspoonfuls (10 mL) every 6 hours
Children 6 to under 12 years of age	45 lbs to under 95 lbs	Take 1 teaspoonful (5mL) every 6 hours
Children under 6 years of age	under 45 lbs	ask a doctor

Other information

- store at controlled room temperature 20°- 25°C (68°- 77°F)
- Each teaspoonful (5 ml) contains: sodium 2mg
- Tamper evident by imprinted heat seal under cap. Do not use if there is evidence of tampering.

Inactive ingredients aspartame, citric acid, glycerine, methylparaben, natural and artificial flavors, potassium citrate, potassium sorbate, propylene glycol, propylparaben, purified water.

Questions or Comments? 1-866-595-5598

NDC 54859-520-16
TAMPER-EVIDENT

TUSNEL C 
SYRUP

**EXPECTORANT
NASAL DECONGESTANT
COUGH SUPPRESSANT
SUGAR-FREE
ALCOHOL-FREE**

*Under Federal law Tusnel C is available without a prescription. Certain State laws may differ.

BULK CONTAINER - NOT FOR HOUSEHOLD USE.

Dispense in a tight, light-resistant container as defined in the USP.

MANUFACTURED FOR:

LLORENS 
PHARMACEUTICAL CORP.
San Juan, P.R. 00926
www.llorenspfarm.com

ONE PINT (473 mL)

Drug Facts

Active Ingredients (in each 5 ml tsp.)	Purpose
Codeine Phosphate..... 10 mg.....	Cough Suppressant
Guafenesin..... 100 mg.....	Expectorant
Pseudoephedrine HCl..... 30 mg.....	Nasal Decongestant

Uses • helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive • temporarily relieves: nasal congestion due to the common cold, hay fever or other upper respiratory allergies and nasal congestion associated with sinusitis, cough due to minor bronchial irritation as may occur with the common cold • temporarily restores freer breathing through the nose • calms the cough control center and relieves coughing

Warnings Do not use this product if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to an enlarged prostate gland • cough that occurs with too much phlegm (mucus) • a breathing problem or persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema • chronic pulmonary disease or shortness of breath, or who are taking other drugs

When using this product • do not exceed recommended dosage
• may cause or aggravate constipation

Stop use and ask a doctor if • you get nervous, dizzy or sleepless • symptoms do not get better within 7 days or are accompanied by fever • cough lasts more than 7 days, comes back, or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

If you are pregnant or breast-feeding ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact Poison Control Center right away.

Directions: Follow dosage chart
• dosage may be repeated every 6 hours. Not to exceed 4 doses in 24-hours. Giving a higher dose than recommended by a doctor could result in serious side effects for your child. A special measuring device should be used to give an accurate dose of this product to children under 6 years of age.

Age	Weight	Dose
children 12 years and over	95 lbs and up	2 teaspoonfuls every 4 to 6 hours
children 6 to under 12 years of age	45 lbs to under 95 lbs	1 teaspoonful every 4 to 6 hours
children under 6 years of age	under 45 lbs	Consult a doctor

Other information • store at controlled room temperature 20°–25°C (68°–77°F)

• Each teaspoonful (5 ml) contains: sodium 2mg

• **Tamper Evident:** Do not use this product if breakaway band on cap seal is broken or missing.

Inactive ingredients: citric acid, FD&C Red #40, flavor, propylene glycol, sodium benzoate, sodium saccharin, sorbitol and water.

Questions or comments: 1-866-595-5598

Distributed By: Llorens Pharmaceutical Corp, San Juan, P.R. 00926

Rev 11/08


N 3 54859-520-16 6

NDC 54859-520-16
TAMPER-EVIDENT

TUSNEL C

SYRUP

**EXPECTORANT
NASAL DECONGESTANT
COUGH SUPPRESSANT
SUGAR-FREE
ALCOHOL-FREE**

*Under Federal law Tusnel C is available without a prescription. Certain State laws may differ.

BULK CONTAINER - NOT FOR HOUSEHOLD USE.

Dispense in a tight, light-resistant container as defined in the USP.

MANUFACTURED FOR:
LLORENS
PHARMACEUTICAL CORP.
San Juan, P.R. 00926
www.llorenspfarm.com

ONE PINT (473 mL)

Drug Facts

Active Ingredients (in each 5 ml tsp.)	Purpose
Codeine Phosphate..... 10 mg.....	Cough Suppressant
Guafenesin..... 100 mg.....	Expectorant
Pseudoephedrine HCl..... 30 mg.....	Nasal Decongestant

Uses • helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive • temporarily relieves: nasal congestion due to the common cold, hay fever or other upper respiratory allergies and nasal congestion associated with sinusitis, cough due to minor bronchial irritation as may occur with the common cold • temporarily restores freer breathing through the nose • calms the cough control center and relieves coughing

Warnings Do not use this product if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to an enlarged prostate gland • cough that occurs with too much phlegm (mucus) • a breathing problem or persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema • chronic pulmonary disease or shortness of breath, or who are taking other drugs

When using this product • do not exceed recommended dosage
• may cause or aggravate constipation

Stop use and ask a doctor if • you get nervous, dizzy or sleepless • symptoms do not get better within 7 days or are accompanied by fever • cough lasts more than 7 days, comes back, or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

If you are pregnant or breast-feeding ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact Poison Control Center right away.

Directions: Follow dosage chart
• dosage may be repeated every 6 hours. Not to exceed 4 doses in 24-hours. Giving a higher dose than recommended by a doctor could result in serious side effects for your child. A special measuring device should be used to give an accurate dose of this product to children under 6 years of age.

Age	Weight	Dose
children 12 years and over	95 lbs and up	2 teaspoonfuls every 4 to 6 hours
children 6 to under 12 years of age	45 lbs to under 95 lbs	1 teaspoonful every 4 to 6 hours
children under 6 years of age	under 45 lbs	Consult a doctor

Other information • store at controlled room temperature 20°–25°C (68°–77°F)
• Each teaspoonful (5 ml) contains: sodium 2mg
• **Tamper Evident:** Do not use this product if breakaway band on cap seal is broken or missing.

Inactive ingredients: citric acid, FD&C Red #40, flavor, propylene glycol, sodium benzoate, sodium saccharin, sorbitol and water.

Questions or comments: 1-866-595-5598

Distributed By: Llorens Pharmaceutical Corp, San Juan, P.R. 00926

Rev 11/08



TUSNEL C

codeine phosphate, guaifenesin, pseudoephedrine hcl liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:54859-520
Route of Administration	ORAL	DEA Schedule	CV

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CODEINE PHOSPHATE (UNII: GSL05Y1MN6) (CODEINE ANHYDROUS - UNII:UX6OWY2V7J)	CODEINE PHOSPHATE	10 mg in 5 mL
Guaifenesin (UNII: 495W7451VQ) (Guaifenesin - UNII:495W7451VQ)	Guaifenesin	100 mg in 5 mL
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL (UNII: 506T60A25R)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54859-520-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/01/1995	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	01/01/1995	

Labeler - Llorens Pharmaceutical International Division (037342305)

Revised: 12/2020

Llorens Pharmaceutical International Division