

IBUPROFEN - ibuprofen capsule, liquid filled

MARKSANS PHARMA LIMITED

Active ingredient (in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*
(present as the free acid and potassium salt)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - o headache
 - o toothache
 - o backache
 - o menstrual cramps
 - o the common cold
 - o muscular aches
 - o minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:

- chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
-
- pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains: **potassium 20 mg**
- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F).

Inactive ingredients

ammonium hydroxide, FD&C Green No. 3, gelatin, iron oxide black, medium-chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, shellac, sorbitan monooleate, sorbitol sorbitan

Questions or comments?

call toll free **1-877-376-4271**

PRINCIPAL DISPLAY PANEL

Ibuprofen capsules 200 mg (Liquid Filled Capsules)

NDC 25000-133-43

Ibuprofen capsules 200 mg (Liquid filled capsules)

10's count bottle label

**Ibuprofen Capsules,
200 mg**

Pain Reliever / Fever Reducer (NSAID)

****Liquid Filled Capsules****10 Softgels******Marksans****READ AND KEEP CARTON
FOR COMPLETE WARNINGS
AND INFORMATION**

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Uses

- temporarily relieves minor aches and pains due to: headache, toothache, backache, menstrual cramps, the common cold, muscular aches, minor pain of arthritis.
- temporarily reduces fever

LOT :

EXP :

R604/23

Warnings

■ **Ask your doctor before use** if you are pregnant, under a doctor's care for a serious condition, age 60 or over, taking any other drug or have stomach problems. ■ This product may cause a **severe allergic reaction**, especially in

people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Do not use** this product if you have ever had an

allergic reaction to any pain reliever/fever reducer. ■ **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have

had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day

while using this product ■ take more or for a longer time than directed. ■ **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for

longer than directed.

Directions ■ **do not take more than directed** ■ **the smallest effective dose should be used** ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1

capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor. ■ children under 12 years: ask a doctor.

Each capsule contains: **potassium 20 mg**

Questions or comments?call toll free **1-877-376-4271**

Manufactured for:
Time-Cap Labs, Inc.
7 Michael Avenue, Farmingdale,
NY 11735, USA

Manufactured by:
Marksans Pharma Ltd.
Plot No. L-82, L-83,
Verna Indl. Estate,
Verna, GOA - 403 722, India.
Mfg. Lic. No. : GO/DRUGS/515

Ibuprofen capsules 200 mg (Liquid filled capsules) 10's count carton label



NDC 25000-133-01

Ibuprofen capsules 200 mg (Liquid filled capsules) 20's count bottle label

NDC 25000-133-01

Ibuprofen Capsules, 200 mg

Pain Reliever / Fever Reducer (NSAID)

****Liquid Filled Capsules**

20 Softgels**



Marksans

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Uses

- temporarily relieves minor aches and pains due to: headache, toothache, backache, menstrual cramps, the common cold, muscular aches, minor pain of arthritis.
- temporarily reduces fever

LOT :

EXP :

R604/23

Warnings

■ **Ask your doctor before use** if you are pregnant, under a doctor's care for a serious condition, age 60 or over, taking any other drug or have stomach problems. ■ This product may cause a **severe allergic reaction**, especially in

people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Do not use** this product if you have ever had an

allergic reaction to any pain reliever/fever reducer. ■ **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have

had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day

while using this product ■ take more or for a longer time than directed. ■ **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for

longer than directed.

Directions ■ **do not take more than directed** ■ **the smallest effective dose should be used** ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1

capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor. ■ children under 12 years: ask a doctor.

Each capsule contains: **potassium 20 mg**

Questions or comments?

call toll free **1-877-376-4271**

Manufactured for:
Time-Cap Labs, Inc.
7 Michael Avenue, Farmingdale,
NY 11735, USA

Manufactured by:
Marksans Pharma Ltd.
Plot No. L-82, L-83,
Verna Indl. Estate,
Verna, GOA - 403 722, India.
Mfg. Lic. No. : GO/DRUGS/515

Ibuprofen capsules 200 mg (Liquid filled capsules)
32's count bottle label

Drug Facts (continued)

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Drug Facts (continued)

Other information

- each capsule contains: **potassium 20 mg**
- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F).
- avoid excessive heat above 40°C (104°F)

Inactive ingredients Ammonium hydroxide, FD&G green no. 3, Gelatin, iron oxide black, Medium chain triglycerides, Polyethylene glycol, Potassium hydroxide, Propylene glycol, Purified water, Shellac, Sorbitan monooleate, Sorbitol sorbitan

Questions or comments?
call toll free 1-877-376-4271

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Manufactured by:
Time-Cap Labs, Inc.
7 Michael Avenue, Farmingdale,
NY 11735, USA

Manufactured by:
Marksans Pharma Ltd.
Plot No. L-82, L-83, Verna Indl. Estate,
Verna, GOA - 403 722, India.
Mfg. Lic. No.: GO/DRUGS/515

20 Softgels**

Pain Reliever / Fever Reducer (NSAID)

Ibuprofen Capsules, 200 mg

NDC 25000-133-01

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION.

Drug Facts

Active ingredient (in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*

Purpose

Pain reliever/
Fever reducer
(present as the free acid and potassium salt)
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older

**Ibuprofen Capsules,
200 mg**

Pain Reliever / Fever Reducer (NSAID)

****Liquid Filled Capsules****32 Softgels******Marksans****READ AND KEEP CARTON
FOR COMPLETE WARNINGS
AND INFORMATION**

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Uses

- temporarily relieves minor aches and pains due to: headache, toothache, backache, menstrual cramps, the common cold, muscular aches, minor pain of arthritis.
- temporarily reduces fever

LOT :

EXP :

R1003/22

Warnings

■ **Ask your doctor before use** if you are pregnant, under a doctor's care for a serious condition, age 60 or over, taking any other drug or have stomach problems. ■ This product may cause a **severe allergic reaction**, especially in

people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Do not use** this product if you have ever had an

allergic reaction to any pain reliever/fever reducer. ■ **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have

had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day

while using this product ■ take more or for a longer time than directed. ■ **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for

longer than directed.

Directions ■ **do not take more than directed** ■ **the smallest effective dose should be used** ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1

capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor. ■ children under 12 years: ask a doctor.

Each capsule contains: **potassium 20 mg**

Questions or comments?
call toll free **1-877-376-4271**

Manufactured for:
Time-Cap Labs, Inc.
7 Michael Avenue, Farmingdale,
NY 11735, USA

Manufactured by:
Marksans Pharma Ltd.
Plot No. L-82, L-83,
Verna Indl. Estate,
Verna, GOA - 403 722, India.
Mfg. Lic. No. : GO/DRUGS/515

Ibuprofen capsules 200 mg (Liquid filled capsules)
32's count carton label



NDC 25000-133-69

Ibuprofen capsules 200 mg (Liquid filled capsules)
300's count bottle label

NDC 25000-133-69

Ibuprofen Capsules, 200 mg

Pain Reliever / Fever Reducer (NSAID)

****Liquid Filled Capsules**

300 Softgels**

Marksans



Drug Facts

Active ingredient (in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)* Pain reliever/
(present as the free acid and potassium salt) Fever reducer
*nonsteroidal anti-inflammatory drug

Purpose

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

LOT :
EXP :



Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you

Drug Facts (continued)

- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or

Drug Facts (continued)

- side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a

Drug Facts (continued)

doctor.

- children under 12 years: ask a doctor

Other information

- each capsule contains: potassium 20 mg
- read all warnings and directions before use.
- store at 20-25°C (68-77°F).
- avoid excessive heat above 40°C (104°F)

Inactive ingredients Ammonium hydroxide, FD&C green no. 3, Gelatin, Iron oxide black, Medium chain triglycerides, Polyethylene glycol, Potassium hydroxide, Propylene glycol, Purified water, Shellac, Sorbitan monooleate, Sorbitol sorbitan

Questions or comments?

call toll free 1-877-376-4271

Drug Facts (continued)

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Manufactured for:

Time-Cap Labs, Inc.
7 Michael Avenue, Farmingdale,
NY 11735, USA

Manufactured by:

Marksans Pharma Ltd.
Plot No. L-82, L-83,
Verna Indl. Estate,
Verna, GOA - 403 722, India.

Mfg. Lic. No. : GO/DRUGS/515

R10322

NDC 25000-133-13

Ibuprofen capsules 200 mg (Liquid filled capsules)
600's count bottle label

NDC 25000-133-13

Ibuprofen Capsules, 200 mg

Pain Reliever / Fever Reducer (NSAID)

****Liquid Filled Capsules**

600 Softgels**



Marksans

Drug Facts

Active ingredient (in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*Pain reliever/
(present as the free acid and potassium salt) Fever reducer
*nonsteroidal anti-inflammatory drug

Purpose

Uses

- temporarily relieves minor aches and pains due to:
 - headache ■ toothache ■ backache
 - menstrual cramps ■ the common cold
 - muscular aches ■ minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include :

- hives ■ facial swelling ■ asthma (wheezing)
- shock ■ skin reddening ■ rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

LOT :

EXP :



Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you :

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke.

These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney

Drug Facts (continued)

disease, asthma or had a stroke

- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint ■ vomit blood ■ have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain ■ trouble breathing ■ weakness in one part or side of body
 - slurred speech ■ leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy

Drug Facts (continued)

Drug Facts (continued)

unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor.
- children under 12 years: ask a doctor

Other information

- each capsule contains: **potassium 20 mg**
- read all warnings and directions before use.
- store at 20-25°C (68-77°F).
- avoid excessive heat above 40°C (104°F)

Inactive ingredients Ammonium hydroxide, FD&C green no. 3, Gelatin, Iron oxide black, Medium chain triglycerides, Polyethylene glycol, Potassium hydroxide, Propylene glycol, Purified water, Shellac, Sorbitan monooleate, Sorbitol sorbitan

Questions or comments?

call toll free **1-877-376-4271**

Do Not Use if seal under bottle cap imprinted with
"SEALED for YOUR PROTECTION" is broken or missing.

Manufactured for:
Time-Cap Labs, Inc.
7 Michael Avenue, Farmingdale,
NY 11735, USA

Manufactured by:
Marksans Pharma Ltd.
Plot No. L-82, L-83,
Verna Indl. Estate,
Verna, GOA - 403 722, India.




Mfg. Lic. No. : GO/DRUGS/515


R1103/22


Ibuprofen capsules 200 mg (Mini liquid filled capsules)

NDC 25000-062-43

Ibuprofen capsules 200 mg (Mini liquid filled capsules)
10's count bottle label


<div>  Marksans NDC25000-062-43 </div> <div> Ibuprofen Capsules 200 mg <i>(Solubilized Ibuprofen Capsules, 200 mg)</i> Pain Reliever/Fever Reducer (NSAID) </div> <div>  </div> <div> 10 Liquid Filled Capsules </div>	READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION	<div>Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.</div>	<div> <div>LOT :</div> <div>EXP :</div> <div>R20322</div> </div>
	<div> Uses ■ temporarily relieves minor aches and pains due to: headache, toothache, backache, menstrual cramps, the common cold, muscular aches, minor pain of arthritis ■ temporarily reduces fever </div>	<div> Warnings ■ Ask your doctor before use if you are pregnant, under a doctor's care for a serious condition, age 60 or over, taking any other drug or have stomach problems. </div> <div>  LIFT HERE <i>for More Drug Facts</i> </div>	

	<p>Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p>Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor</p> <p>Each capsule contains: potassium 20 mg</p> <p>Questions or comments? call toll free 1-877-376-4271</p> <p>Manufactured for: Time-Cap-Labs, Inc., 7 Michael Avenue, Farmingdale, NY 11735, USA</p> <p>Manufactured by: Marksans Pharma Ltd. Plot No. L-82, L-83, Verna Indl. Estate, Verna, Goa-403722, India</p>	Mfg. Lic. No. : GO/DRUGS/515
---	---	------------------------------

<p>■ This product may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away. Do not use this product if you have ever had an allergic reaction to any pain reliever/fever reducer.</p> <p>■ Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. ➡</p>	
---	---

NDC 25000-062-43

Ibuprofen capsules 200 mg (Mini liquid filled capsules)
 10's count carton label



READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredient (in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*
(present as the free acid and potassium salt)
nonsteroidal anti-inflammatory drug

Purpose

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.
Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- stomach pain
- skin redness
- itch
- diarrhea

If an allergic reaction occurs, stop use and seek medical help right away.

202022

Mfg. Lot. No. : G00RUGS515

EXP :
LOT :

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer right before or after heart surgery.

Ask a doctor before use if you have problems or serious side effects from taking pain relievers or fever reducers or you have a history of stomach problems, such as heartburn, you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke or you are taking a diuretic.
Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition, taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin, taking any other drug.
When using this product take with food or milk if stomach upset occurs.

Drug Facts (continued)

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

Do not take more than directed the smallest effective dose should be used.

- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information each capsule contains:

- potassium 20 mg
- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F).
- avoid excessive heat above 40°C (104°F).

Inactive Ingredients ammonium hydroxide, FD&C green no. 3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, shellac, sorbitan monoleate, sorbitol sorbitan.

Questions or comments?

Do Not Use if seal under bottle cap imprinted with "SEALED FOR YOUR PROTECTION" is broken or missing.

Manufactured for: Time-Cap-Labs, Inc., 7 Mitchell Avenue, Farmingdale, NY 11735 USA
Manufactured by: Marksans Pharma Ltd., Plot No. L-82, L-83, Venna Indl. Estate, Venna, Goa-403722, India

M Marksans

NDC25000-062-43


Ibuprofen
Capsules 200 mg
(Solubilized Ibuprofen Capsules, 200 mg)
Pain Reliever/Fever Reducer (NSAID)


10 Liquid Filled Capsules




NEW
Smaller Capsule
Same Strength

Ibuprofen capsules 200 mg (Mini liquid filled capsules)
20's count bottle label


	<p>READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION</p> <p>Uses</p> <ul style="list-style-type: none"> temporarily relieves minor aches and pains due to: headache, toothache, backache, menstrual cramps, the common cold, muscular aches, minor pain of arthritis temporarily reduces fever 	<p>Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.</p> <p>Warnings</p> <ul style="list-style-type: none"> Ask your doctor before use if you are pregnant, under a doctor's care for a serious condition, age 60 or over, taking any other drug or have stomach problems. <p>LIFT HERE for More Drug Facts</p>	<p>LOT : EXP : R20322</p>
---	---	--	-----------------------------------

	<p>Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p>Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor</p> <p>Each capsule contains: potassium 20 mg</p> <p>Questions or comments? call toll free 1-877-376-4271</p> <p>Manufactured for: Time-Cap-Labs, Inc., 7 Michael Avenue, Farmingdale, NY 11735, USA</p> <p>Manufactured by: Marksans Pharma Ltd. Plot No. L-82, L-83, Verna Indl. Estate, Verna, Goa-403722, India</p>	<p>Mfg. Lic. No. : GO/DRUGS/515</p>
--	---	-------------------------------------

<p>■ This product may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away. Do not use this product if you have ever had an allergic reaction to any pain reliever/fever reducer.</p> <p>■ Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. ➡</p>	
---	---

NDC 25000-062-01

Ibuprofen capsules 200 mg (Mini liquid filled capsules)
20's count carton label



READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredient (in each capsule)

Solubilized ibuprofen (equal to 200 mg ibuprofen (NSAID)*)
(present as the free acid and potassium salt)
nonsteroidal anti-inflammatory drug

Purpose

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Warning: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- stomach pain
- skin redness
- itching
- blistering

If an allergic reaction occurs, stop use and seek medical help right away.

R20322

Mfg. Lic. No.: G0DRUGS515

LOT :
EXP :

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing naproxen, or others
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer or right before or after heart surgery.

Ask a doctor before use if you have problems or serious side effects from taking pain relievers or fever reducers or you have a history of stomach problems, such as heartburn, you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke or you are taking a diuretic. **Ask a doctor or pharmacist before use if you are** under a doctor's care for any serious condition, taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin, taking any other drug.

When using this product take with food or milk if stomach upset occurs.

Drug Facts (continued)

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

Do not take more than directed the smallest effective dose should be used. **Adults and children 12 years and over:** take 1 capsule every 4 to 6 hours while symptoms persist. **Children 6 to 11 years:** take 1 capsule every 6 to 8 hours while symptoms persist. **Children under 6 years:** ask a doctor.

Other information each capsule contains:

- potassium 20 mg
- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F).
- avoid excessive heat above 40°C (104°F).

Inactive Ingredients ammonium hydroxide, FD&C green no. 3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, shellac, sorbitan monoleate, sorbitol sorbitan.

Questions or comments? call toll free 1-877-376-4271. **Do Not Use if seal under bottle cap imprinted with "SEALED FOR YOUR PROTECTION" is broken or missing.**

Manufactured for: Time-Cap-Labs, Inc., 7 Michael Avenue, Farmingdale, NY 11735 USA
Manufactured by: Marksans Pharma Ltd., Plot No. L-82, L-83, Verna Indl. Estate, Verna, Goa-403222, India

M Marksans

NDC25000-062-01


Ibuprofen
Capsules 200 mg
(Solubilized Ibuprofen Capsules, 200 mg)
Pain Reliever/Fever Reducer (NSAID)


20 Liquid Filled Capsules



NEW
Smaller Capsule
Same Strength

Ibuprofen capsules 200 mg (Mini liquid filled capsules)
32's count carton label





P03022

Mfg. Lic. No.: G0DRUGS515

EXP :
LOT : 107

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredient (in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*
(present as the free acid and potassium salt)
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever
Fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Major alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.
Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- stomach pain
- skin redness
- itching
- diarrhea

If an allergic reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer or right before or after heart surgery.

Ask a doctor before use if you have problems or serious side effects from taking pain relievers or fever reducers or you have a history of stomach problems, such as heartburn, you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke or you are taking a diuretic.
Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition or taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin or taking any other drug.

When using this product take with food or milk if stomach upset occurs.

Drug Facts (continued)

Other information each capsule contains:

- potassium 20 mg
- read all warnings and directions before use
- Keep carton
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

ammonium hydroxide, FD&C green no. 3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydride, sorbitan monolaurate, sorbitol, sorbitan

Questions or comments? call toll free 1-877-376-4271
Do Not Use if seal under bottle cap imprinted with "SEALED FOR YOUR PROTECTION" is broken or missing.

Manufactured for: Time-Cap-Labs, Inc., 7 Michael Avenue, Farmingdale, NY 11735 USA
Verna Int'l. Estate, Verna, Goa-403722, India

Drug Facts (continued)

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding:

- feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - sudden speech
 - leg swelling
 - pain gets worse or lasts more than 10 days
 - tooth gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear
- If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
- Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

Do not take more than directed the smallest effective dose should be used adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist if pain or fever does not respond to 1 capsule, 2 capsules may be used do not exceed 6 capsules in 24 hours, unless directed by a doctor children under 12 years: ask a doctor



NDC25000-062-04

Ibuprofen
Capsules 200 mg
(Solubilized Ibuprofen Capsules, 200 mg)
Pain Reliever/Fever Reducer (NSAID)

32 Liquid Filled Capsules



NEW
Smaller
Capsule


Same
Strength

Ibuprofen capsules 200 mg (Mini liquid filled capsules)
80's count bottle label

<p>M Marksans NDC25000-062-67</p> <p>Ibuprofen Capsules 200 mg (Solubilized Ibuprofen Capsules, 200 mg)</p> <p>Pain Reliever/Fever Reducer (NSAID)</p> <p>80 Liquid Filled Capsules</p> <p>minis</p> <p>READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION</p>	<p>Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.</p> <p>Uses ■ temporarily relieves minor aches and pains due to: headache, toothache, backache, menstrual cramps, the common cold, muscular aches, minor pain of arthritis ■ temporarily reduces fever</p> <p>Warnings ■ Ask your doctor before use if you are pregnant, under a doctor's care for a serious condition, age 60 or over, taking any other drug or have stomach problems. ■ This product may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away. Do not use this product if you have ever had an allergic reaction to any pain reliever/fever reducer.</p> <p>■ Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning</p>	<p>(anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p>Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor</p> <p>Each capsule contains: potassium 20 mg</p> <p>Questions or comments? call toll free 1-877-376-4271</p> <p>Manufactured for: Time-Cap-Labs, Inc., 7 Michael Avenue, Farmingdale, NY 11735, USA</p> <p>Manufactured by: Marksans Pharma Ltd. Plot No. L-82, L-83, Verna Indl. Estate, Verna, Goa-403722, India Mfg. Lic. No.: GO/DRUGS/515</p>	<p>LOT : EXP : R20322</p>
---	---	---	-----------------------------------

NDC 25000-062-67

Ibuprofen capsules 200 mg (Mini liquid filled capsules)
80's count carton label



NDC 25000-062-67

Mfg. Lic. No. : G0DRUGS615

P20322

LOT :
EXP :

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredient (in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID) (present as the free acid and potassium salt), nonsteroidal anti-inflammatory drug

Purpose

Pain reliever
Fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - backache
 - muscular aches
 - the common cold
 - menstrual cramps
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- stomach cramps
- skin redness
- itching
- blistering

If an allergic reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer. **Use right before or after heart surgery.**

Ask a doctor before use if:

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are:

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product:

- take with food or milk if stomach upset occurs

Drug Facts (continued)

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - leg faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

Do not take more than directed. Use the smallest effective dose should be used. Adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 capsule, 2 capsules may be used. Do not exceed 6 capsules in 24 hours, unless directed by a doctor. Children under 12 years: ask a doctor.

Drug Facts (continued)

Other information each capsule contains:

- potassium 20 mg
- read all warnings and directions before use
- keep carton
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

ammonium hydroxide, FD&C green no. 3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, shellac, sorbitan monooleate, sorbitol sorbitan

Questions or comments? call toll free 1-877-376-4271

Do Not Use if seal under bottle cap imprinted with "SEALED FOR YOUR PROTECTION" is broken or missing.

Manufactured for: Time-Cap-Labs, Inc., 7 Michael Avenue, Farmingdale, NY 11735, USA
Manufactured by: Marksans Pharma Ltd., Plot No. L-92, L-93, Verna Indl. Estate, Verna, Goa-403722, India



NDC25000-062-67

Ibuprofen Capsules 200 mg

(Solubilized Ibuprofen Capsules, 200 mg)
Pain Reliever/Fever Reducer (NSAID)

80 Liquid Filled Capsules



NEW
Smaller
Capsule

Same
Strength

Ibuprofen capsules 200 mg (Mini liquid filled capsules)
300's count PET bottle label

Ibuprofen

Capsules 200 mg

(Solubilized Ibuprofen Capsules, 200 mg)
Pain Reliever/Fever Reducer (NSAID)

300 Liquid Filled Capsules



Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.



LIFT HERE
for More Drug Facts

Drug Facts

Active ingredient (in each capsule)	Purpose
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*Pain reliever/Fever reducer (present as the free acid and potassium salt)	
*nonsteroidal anti-inflammatory drug	

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

LOT :
EXP :
P20322

Drug Facts (continued)**Warnings**

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older

- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product

- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or

Drug Facts (continued)

- had a stroke

- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin

- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:

- feel faint

- vomit blood

- have bloody or black stools

- have stomach pain that does not get better

- you have symptoms of heart problems or stroke:

- chest pain

- trouble breathing

- weakness in one part or side of body

- slurred speech

- leg swelling

- pain gets worse or lasts more than 10 days

- fever gets worse or lasts more than 3 days

- redness or swelling is present in the painful area

- any new symptoms appear

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed

- the smallest effective dose should be used

- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist

- if pain or fever does not respond to 1 capsule, 2 capsules may be used

- do not exceed 6 capsules in 24 hours, unless directed by a doctor

- children under 12 years: ask a doctor

Other information

- each capsule contains: potassium 20 mg

- read all warnings and directions before use. Keep carton.

- store at 20-25°C (68-77°F)

- avoid excessive heat above 40°C (104°F).

Inactive ingredients

ammonium hydroxide, FD&C green no.3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, shellac, sorbitan monooleate, sorbitol sorbitan

Drug Facts (continued)**Questions or comments?** call toll free 1-877-376-4271

Manufactured for:

Time-Cap-Labs, Inc.

7 Michael Avenue,

Farmingdale, NY 11735, USA

Manufactured by:

Marksans Pharma Ltd.

Plot No. L-82, L-83,

Verna Indl. Estate, Verna,

Goa-403722, India

Mfg. Lic. No.:GO/DRUGS/515

NDC 25000-062-13

Ibuprofen capsules 200 mg (Mini liquid filled capsules)
600's count bottle label

Pannel 1

Marksans

NDC25000-062-13

NEW
Smaller Capsule

Same Strength

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

LIFT HERE
for More Drug Facts

Ibuprofen

Capsules 200 mg

(Solubilized Ibuprofen Capsules, 200 mg)
Pain Reliever/Fever Reducer (NSAID)

600 Liquid Filled Capsules

N 3112500010621312

Drug Facts	
Active ingredient (in each capsule) Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*Pain reliever/Fever reducer (present as the free acid and potassium salt) *nonsteroidal anti-inflammatory drug	Purpose Pain reliever/Fever reducer
Uses ■ temporarily relieves minor aches and pains due to: ■ headache ■ toothache ■ backache ■ menstrual cramps ■ the common cold ■ muscular aches ■ minor pain of arthritis ■ temporarily reduces fever	

LOT :

EXP :

R20322

Pannel 2

Drug Facts (continued)	Drug Facts (continued)
Warnings Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or	■ had a stroke ■ you are taking a diuretic Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug When using this product ■ take with food or milk if stomach upset occurs Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

Pannel 3

Drug Facts (continued)	Drug Facts (continued)
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor Other information ■ each capsule contains: potassium 20 mg ■ read all warnings and directions before use. Keep carton. ■ store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F). Inactive ingredients ammonium hydroxide, FD&C green no.3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water,	Questions or comments? call toll free 1-877-376-4271 Manufactured for: Time-Cap-Labs, Inc., 7 Michael Avenue, Farmingdale, NY 11735, USA Manufactured by: Marksans Pharma Ltd. Plot No. L-82, L-83, Verna Indl. Estate, Verna, Goa-403722, India Mfg. Lic. No.: GO/DRUGS/515

IBUPROFEN

ibuprofen capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:25000-133
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
AMMONIA (UNII: 5138Q19F1X)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
SORBITOL (UNII: 506T60A25R)	
WATER (UNII: 059QF0KO0R)	
SHELLAC (UNII: 46N107B71O)	
SORBITAN MONOOLEATE (UNII: 06XEA2VD56)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
FERROUS OXIDE (UNII: G7036X8B5H)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	

Product Characteristics

Color	GREEN	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	133
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:25000-133-23	5 in 1 DRUM	09/01/2017	
1		2000 in 1 BAG; Type 0: Not a Combination Product		
2	NDC:25000-133-18	4 in 1 BOX	09/01/2017	
2		2500 in 1 BAG; Type 0: Not a Combination Product		
3	NDC:25000-133-04	1 in 1 CARTON	09/01/2017	
3		32 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:25000-133-13	1 in 1 BOX	09/01/2017	
4		600 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:25000-133-43	1 in 1 CARTON	09/01/2017	
5		10 in 1 BOTTLE; Type 0: Not a Combination Product		
6	NDC:25000-133-01	1 in 1 CARTON	09/01/2017	
6		20 in 1 BOTTLE; Type 0: Not a Combination Product		
7	NDC:25000-133-69	300 in 1 BOTTLE; Type 0: Not a Combination Product	09/01/2017	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA079205	09/01/2017	

IBUPROFEN			
ibuprofen capsule, liquid filled			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:25000-062
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)		IBUPROFEN	200 mg
Inactive Ingredients			

Ingredient Name	Strength
AMMONIA (UNII: 5138Q19F1X)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
SORBITOL (UNII: 506T60A25R)	
WATER (UNII: 059QF0KO0R)	
SHELLAC (UNII: 46N107B71O)	
SORBITAN MONOOLEATE (UNII: 06XEA2VD56)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
FERROUS OXIDE (UNII: G7036X8B5H)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	

Product Characteristics

Color	GREEN	Score	no score
Shape	CAPSULE	Size	6mm
Flavor		Imprint Code	62
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:25000-062-43	1 in 1 CARTON	08/18/2020	
1		10 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:25000-062-01	01 in 1 CARTON	08/18/2020	
2		20 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:25000-062-04	1 in 1 CARTON	08/18/2020	
3		32 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:25000-062-67	1 in 1 CARTON	08/18/2020	
4		80 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:25000-062-13	600 in 1 BOTTLE; Type 0: Not a Combination Product	08/18/2020	
6	NDC:25000-062-87	1 in 1 BOX	08/18/2020	
6		2500 in 1 BAG; Type 0: Not a Combination Product		
7	NDC:25000-062-69	300 in 1 BOTTLE; Type 0: Not a Combination Product	11/25/2021	

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
-----------	---------------------------------	-----------------	---------------

Category	Citation	Date	Date
ANDA	ANDA079205	08/18/2020	

Labeler - MARKSANS PHARMA LIMITED (925822975)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	MANUFACTURE(25000-062, 25000-133)

Revised: 4/2023 MARKSANS PHARMA LIMITED