

**HEAD CONGESTION FLU SEVERE PE- acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated**  
**CHAIN DRUG MARKETING ASSOCIATION INC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**1174-QCH-2022-0803**

**Drug Facts**

<b>Active ingredients (in each caplet)</b>	<b>Purposes</b>
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 100 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses**

- temporarily relieves these symptoms due to the common cold:
  - nasal congestion
  - headache
  - minor aches and pains
  - cough
  - sore throat
  - sinus congestion and pressure
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive
- temporarily reduces fever

**Warnings**

**Liver warning**

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening

- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### **Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

### **When using this product do not exceed recommended dosage**

#### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

### **Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"><li>▪ take 2 caplets every 4 hours</li><li>▪ do not take more than 10 caplets in 24 hours</li></ul>
children under 12 years	<ul style="list-style-type: none"><li>▪ ask a doctor</li></ul>

### Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information and warnings

### Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

### PRINCIPAL DISPLAY PANEL

QUALITY CHOICE®

NDC 63868-102-24

†Compare to Active Ingredients in SUDAFED PE® Head Congestion + Flu Severe  
Head Congestion + Flu Severe

Acetaminophen, Dextromethorphan HBr, Guaifenesin, Phenylephrine HCl

Pain Reliever | Fever Reducer, Cough Suppressant, Expectorant, Nasal Decongestant

For Relief of:

Sinus Pressure

Headache

Sore Throat

Cough

Chest Congestion

Actual Size

**DO NOT USE IF BLISTER UNITS  
ARE TORN OR BROKEN**



Distributed by CDMA, Inc.  
Novi, MI 48375  
www.qualitychoice.com  
Questions: 800-935-2362

**†** This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Sudafed PE®. Head Congestion + Flu Severe.

<p><b>Drug Facts</b> (continued)</p> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ store between 20-25°C (68-77°F) in a dry place</li> <li>■ retain carton for complete product information and warnings</li> </ul> <p><b>Inactive ingredients</b></p> <p>colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone,</p>	<p>◀</p>
---	----------

FD&C yellow #6 aluminum lake, magnesium stearate, malbiodexin, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide,

**Warning Facts (continued)**

- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)
- ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin
- When using this product do not exceed recommended dosage
- When use and ask a doctor if:
  - nervousness, dizziness, or sleeplessness occur
  - liver, gas, nasal congestion, or cough gets worse or lasts more than 7 days
  - fever gets worse or lasts more than 3 days
  - pain or tenderness or swelling is present
  - cough comes back or occurs with rash or headache that lasts
- These could be signs of a serious condition.
- Keep out of reach of children.
- In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed (see overdose warning)
- adults and children 12 years and over:
  - take 2 capsules every 4 hours
  - do not take more than 10 capsules in 24 hours
- ask a doctor

12 years

**Warnings**

**Drug Facts** (continued)

**Other warnings:** This product contains acetaminophen. The maximum daily dose of this product is 10 capsules (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours or with other drugs containing acetaminophen.

**3 or more alcoholic drinks every day while using this product.**

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Severe throat warning:** If your throat is severe, persists for more than 2 days, is accompanied by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- liver disease ■ heart disease ■ high blood pressure
- thyroid disease ■ diabetes

F117402QCH\_R1



Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guafenesin 100 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

  

Uses	Drug Facts (continued)
Temporarily relieves these symptoms due to the common cold: <ul style="list-style-type: none"> <li>nasal congestion</li> <li>headache</li> <li>minor aches and pains</li> <li>cough</li> <li>sore throat</li> </ul>	<ul style="list-style-type: none"> <li>Temporarily relieves these symptoms due to the common cold:               <ul style="list-style-type: none"> <li>nasal congestion</li> <li>headache</li> <li>minor aches and pains</li> <li>cough</li> <li>sore throat</li> </ul> </li> <li>helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive</li> <li>temporarily reduces fever</li> </ul>



NDC 63868-102-24

**†Compare to Active  
Ingredients in SUDAFED PE®  
Head Congestion + Flu Severe**

# Head Congestion + Flu Severe PE

**Acetaminophen, Dextromethorphan HBr, Guaifenesin, Phenylephrine HCl**

**Pain Reliever | Fever Reducer, Cough Suppressant, Expectorant, Nasal Decongestant**

**For Relief of:**

### Sinus Pressure

## Sore Throat

## Chest Congestion

Headache

## Cough

Actual Size

## 24 CAPLETS



HEAD CONGESTION FLU SEVERE PE

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-102
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients	
Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (UNII: 2S7830E561)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	orange	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	AAA;1134
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date

<b>1</b>	NDC:63868-102-24	2 in 1 CARTON	08/03/2022	
<b>1</b>		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>		<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph final		part341	08/03/2022	

**Labeler -** CHAIN DRUG MARKETING ASSOCIATION INC (011920774)

Revised: 8/2022

CHAIN DRUG MARKETING ASSOCIATION INC